

Chem. 1987, 220, 295.

6. Itaya, K.; Shibayama, H.; Akahoshi, H.; Toshima, S. *J. Appl. Phys.* 1982, 53, 804.
7. Neff, V. D. *J. Electrochem. Soc.* 1985, 132, 1382.
8. (a) Bharathi, S.; Yegnaraman, V.; Rao, G. P. *Langmuir* 1995, 11, 666. (b) Upadhyay, D. N.; Yegnaraman, V.; Rao, G. P. *Langmuir* 1996, 12, 4249.
9. Ulman, A. *An Introduction to Ultrathin Organic Films:*

From Langmuir-Blodgett to Self-Assembly; Academic Press: Boston, 1991.

10. Hong, H.-G.; Mallouk, T. E. *Langmuir* 1991, 7, 2362.
11. Kiss, T.; Balla, J.; Nagy, G.; Kozłowski, H.; Kowalik, J. *Inorg. Chim. Acta* 1987, 138, 25.
12. Herren, F.; Fischer, P.; Ludi, A.; Halg, W. *Inorg. Chem.* 1980, 19, 956.

Dehydrogenation of Ketones by Palladium(II) trifluoroacetate-Phosphine and -Sulfide Systems

Young-ae Whang Park* and Hyun Hee Oh

Department of Chemistry, Sang Myung University, Seoul 110-743, Korea
Received June 10, 1997

Activation of carbon-hydrogen bonds is an important subject in organometallic chemistry. We have investigated the activation of aromatic carbon-hydrogen bond by Pd(OAc)₂-phosphine system and have showed the system to be more effective than Pd(OAc)₂-sulfide system.¹ Herein, we wish to report the activation of aliphatic carbon-hydrogen bond of cyclic ketones (cyclopentanone and cyclohexanone) using Pd(CF₃CO₂)₂-phosphine and -sulfide systems. Although many papers on the activation of aliphatic C-H bond have been published previously²⁻⁵ and dehydrogenation reactions of cyclic ketones were studied by Fuchita *et al.* using Pd(OAc)₂-SBU₂ system,⁶ trifluoroacetate as an anion is an interesting choice for the following reasons: 1) the CF₃CO₂⁻ ion is relatively weaker than CH₃CO₂⁻ ion in basicity and M-O₂CCF₃ bond is quite labile,^{2,7} which can be dissociated more easily to give a vacant site of the metal complexes, 2) however, trifluoroacetate ligand might reduce the catalytic activity due to the poor basicity as well, and 3) trifluoroacetate does not contain C-H bond itself. Also, phosphine has been selected to compare the catalytic activity with the sulfide system. The reaction of cyclic ketone has been carried out for 5 hours in the presence of Pd(CF₃CO₂)₂ with phosphine or sulfide (PR₃=P(*p*-C₆H₄CH₃)₃, P(*p*-C₆H₄F)₃, PPh₃, P(OBuⁿ)₃ and SR₂=S(CH₂CH₂CH₃)₂, S{CH(CH₃)₂}₂, S{CH(CH₃)(C₂H₅)₂}, S(CH₂Ph)₂, S{C(CH₃)₃}₂) at 75 °C in this work. The mechanism of dehydrogenation of cyclic ketones is considered to be the formation of α-palladated cyclic ketone followed by the formation of dehydrogenation

product from β-hydride elimination.⁶

The yield of 2-cyclohexen-1-one produced from cyclohexanone is highest (589%) when the molar ratio of Pd(CF₃CO₂)₂ to P(*p*-C₆H₄CH₃)₃ is 1:2, while the highest yield was obtained for the molar ratio of 1:2/3 in Pd(OAc)₂-SBU₂ system. The yields based on the amount of Pd(CF₃CO₂)₂ used are the average value of three or four trials and have been measured on GC with hexadecane as an internal standard. Lower yields were obtained for the ratio of 1:2/3 (417%), 1:1 (513%) and others. Therefore, the molar ratio has been kept 1:2 for all of the reactions in this work to maintain the identical reaction condition. Although it is not clear for now, the ratio of 1:2 may suggest the structure of the intermediate involves two phosphine ligands coordinated to a palladium metal. Table 1 shows the yields of cycloalkenones catalyzed by Pd(CF₃CO₂)₂-phosphine and -sulfide systems. In dehydrogenation reaction of cyclohexanone, the highest yield (589%) of 2-cyclohexen-1-one has been afforded by P(*p*-C₆H₄CH₃)₃, which includes the most basic substituent among the series of phosphines. It is interesting to point out P(*p*-C₆H₄F)₃ gives more yield (363%) than PPh₃ (307%). The *para*-fluorine of the phenyl group seems to be rather electron donating than electron withdrawing due to the resonance effect. Yield is 275% in the absence of phosphine, which means Pd(CF₃CO₂)₂ itself can activate the aliphatic C-H bond of ketone very well. For reference, the same reaction using Pd(OAc)₂ was carried out under the identical condition, and the yield is extremely low (28%), therefore, trifluoroacetate anion reinforces catalytic activity more than acetate does. On the other hand, P(OBuⁿ)₃ containing an electronegative and bulky substituent gives much lower yield (39%) than Pd(CF₃CO₂)₂ itself. Therefore, more basic and less hindered phosphines⁸ afford more products.

In case of sulfide series, S(CH₂CH₂CH₃)₂ affords the highest yield (422%) and the second yield (320%) is obtained for S{CH(CH₃)₂}₂ and SPh₂ gives 223% of the yield. It seems the sulfide with more basic substituent gives more yield than phosphine series. However, steric factor must be




Table 1. Yields of Cycloalkanones by Pd(CF₃CO₂)₂-Phosphine and -Sulfide Systems^a

Product	Phosphine	Yield (%) ^b	Sulfide	Yield (%) ^b
2-cyclohexen-1-one	P(<i>p</i> -C ₆ H ₄ CH ₃) ₃	589	S(CH ₂ CH ₂ CH ₃) ₂	422
	P(<i>p</i> -C ₆ H ₄ F) ₃	363	S{CH(CH ₃) ₂ } ₂	320
	PPh ₃	307	SPh ₂	223
	P(OBu ⁿ) ₃	39	S(CH ₂ Ph) ₂	57
	None	275	S{C(CH ₃) ₃ } ₂	51
2-cyclopenten-1-one	P(<i>p</i> -C ₆ H ₄ CH ₃) ₃	200	S(CH ₂ CH ₂ CH ₃) ₂	133
	P(<i>p</i> -C ₆ H ₄ F) ₃	195	S{CH(CH ₃) ₂ } ₂	134
	PPh ₃	164	S{CH(CH ₃)(C ₂ H ₅)} ₂	93
	P(OBu ⁿ) ₃	85	S(CH ₂ Ph) ₂	57
	None	161	S{C(CH ₃) ₃ } ₂	45

^a Reaction conditions: Pd(CF₃CO₂)₂, 50 mg (0.163 mmol); phosphines and sulfides, 2 molar equivalents to Pd(CF₃CO₂)₂; cycloalkanone 5.0 mL; 70 °C for 5 h. ^b Calculated based on Pd(CF₃CO₂)₂.

also important since the sulfides with bulky substituents such as S{C(CH₃)₃}₂ and S(CH₂Ph)₂ give very low yield, 51% and 57% respectively. Although direct comparison between the effect of phosphine and sulfide is not available from this data due to the different substituents, phosphines seems to show better catalytic activity than sulfide from the results of PPh₃ (307%) and SPh₂ (223%), containing the same phenyl substituent.

The effect of the substituents of both phosphines and sulfides on the yield of 2-cyclopenten-1-one is similar to the case of 2-cyclohexen-1-one except the yield of 2-cyclohexen-1-one being higher for the same catalytic system. In the phosphine series, the highest yield of 2-cyclopenten-1-one has been obtained for P(*p*-C₆H₄CH₃)₃ (200%) and P(*p*-C₆H₄F)₃ (195%) gives more yield than PPh₃ (164%), again. The product has been formed in good yield (161%) for Pd(CF₃CO₂)₂ only. Addition of sulfide does not seem to increase the catalytic activity of Pd(CF₃CO₂)₂, in fact, decrease the activity in this case. However, S(CH₂CH₂CH₃)₂ and S{CH(CH₃)₂}₂ systems act as catalysts since the yield of the product is over 100%. Pd(OAc)₂-SBU₂^{1,6} system affords 412% yield of 2-cyclopenten-1-one and 371% of 2-cyclohexen-1-one, but the yields were so low for other sulfides that data was available only for SBU₂.⁶

In conclusion, 1) trifluoroacetate is a good choice as an

anion since Pd(CF₃CO₂)₂ itself catalyzes the C-H bond activation very well, and both Pd(CF₃CO₂)₂-phosphine and Pd(CF₃CO₂)₂-sulfide systems act as reasonably good catalysts compared with Pd(OAc)₂-sulfide system, 2) phosphines or sulfides containing more basic and less hindered substituents increase the yield of the product and 3) Pd(CF₃CO₂)₂-phosphine system seems to catalyze more effectively than Pd(CF₃CO₂)₂-sulfide system. 4) 2-cyclohexen-1-one is obtained in more yield than 2-cyclopenten-1-one.

Experimental

Materials. Palladium trifluoroacetate, phosphines, sulfides, 1-cyclopentanone and 2-cyclohexanone purchased from Aldrich were all reagent-grade and were used without further purification. Hexane was distilled over sodium at reflux under N₂ just prior to use.

Measurements. Reactions were carried out using 50 mg (0.163 mmol) of Pd(CF₃CO₂)₂, 0.326 mmol of phosphine or sulfide in 5 mL of cycloalkanone at 70 °C under O₂ for 5 h. After decanting the solution, hexane was added to remove the palladium metal by precipitation and the residue was filtered off. The collected solution was analyzed by Hewlett Packard 5890 Series II spectrometer on a HP-5 column.

Acknowledgment. This work was supported by Korea Science and Engineering Foundation (951-0303-082-2).

References

- Park, Y. W.; Hwang, S. W. *Bull. Korean Chem. Soc.* 1997, 18, 218.
- Gretz, E.; Oliver, T. F.; Sen, A. *J. Am. Chem. Soc.* 1987, 109, 8109.
- Bennett, M. A.; Robertson, G. B.; Whimp, P. O.; Yoshida, T. *J. Am. Chem. Soc.* 1973, 95, 3028.
- Shul'pin, G. B.; Nizova, G. V.; Shilov, A. E. *J. Chem. Soc., Chem. Commun.* 1983, 671.
- Ittel, S. D.; Tolman, C. A.; English, D.; Jesson, J. P. *J. Am. Chem. Soc.* 1978, 100, 7577.
- Fuchita, Y.; Harada, Y. *Inorg. Chim. Acta* 1993, 208, 43.
- Garner, C. D.; Hughes, B. *Adv. Inorg. Chem. Radiochem.* 1975, 17, 1.
- Tolman, C. A. *Chem. Rev.* 1977, 77, 313.