

Characteristics of Spectral Reflectance for Corns and Legumes at OSMI(Ocean Scanning Multi-spectral Imager) Bands

Suk-Young Hong, Sang-Kyu Rim, Sun-Joo Hwang, and Sun-Oh Kim

Remote Sensing Lab., Soil Management Division,
National Institute of Agricultural Science and Technology(NIAST)

OSMI 파장영역에서 옥수수과 두류작물의 분광반사특성

홍석영 · 임상규 · 황전주 · 김선오

농업과학기술원 토양관리과 원격탐사실

Abstract

Spectral reflectance data of upland crops at OSMI bands were collected and evaluated for the feasibility of crop discrimination knowledge-based on crop calendar. Effective bands and their ratio values for discriminating corn from two other legumes were defined with OSMI equivalent bands and their ratio values. For corn discrimination from two other legumes, peanut and soybean, June 22 among measurements dates was the best since all OSMI equivalent bands and their ratio values in June 22 were highly significant for corn separability.

Phenological growth stage of a silage corn (rs510) could be estimated as a function of spectral reflectance in vegetative stage. Five growth stage prediction models were generated by the SAS procedures REG and STEPWISE with OSMI equivalent bands and their ratio values in vegetative stage.

Keywords : OSMI bands, spectral reflectance, upland crops, crop separability

요 약

주요 발작물인 옥수수, 땅콩, 콩에 대한 시계열 분광반사율 자료를 수집하여, 작물의 재배력과 분광 반사 특성 차이에 따른 작물간 구분여부를 살펴보았다. 분광방사계를 이용하여 각 작물의 생육기간 동안 0.33~1.10 μm 범위의 파장영역에서 약 12회 분광반사율을 측정하였다. OSMI(Ocean Scanning Multi-spectral Imager)의 중심 파장역 또는 중심 파장역과 가장 가까운 채널인 445nm, 490nm, 510nm, 555nm, 690nm, 865nm를 각각 OSMI 동등 밴드 1, 2, 3, 4, 5, 6으로 하였다. OSMI 동등 밴드와 밴드간 비율을 이용하여 일반선형모형(GLM) 분석한 결과, 분광반사율 측정일 중 옥수수와 두류작물을 구분 하기에 가장 적합한 시기는 6월 22일인 것으로 나타났다. 또한 옥수수의 생육에 따른 분광반사율과 밴드간 비율을 이용하여 청예용 옥수수인 rs510의 영양생장기 동안 생육단계를 추정하는 다중회귀식을 만들었다.

1. Introduction

Image interpretation for land surface cover is basic technology in remote sensing. The same may be said of crop discrimination in agricultural remote sensing(秋山 *et al.*, 1996). In general, spectral reflectance signatures in agricultural fields are shown to be different from crop species, developmental stage, cropping pattern, and cropping environment. For crop classification with remotely sensed data, spectral separability assessment based on time-series spectral reflectance and crop calendar has to be carried out as a basic study.

Panigrahy and Chakraborty(1998) used temporal remote sensing data, along with soil, physiographic rainfall and temperature information on model to derive a potato-growing environment index. Jensen and Middelkoop(1992) revealed that the accuracy of the knowledge-based classification was 6 to 20 per cent better compared with a maximum likelihood classification, depending on the spectral class discrimination.

On the other hand, population of Korea has been concentrated in the specific area and the landuse is subdivided extremely. Especially agricultural unit of upland crop is very small. Crop inventory with satellite data would be hindered by the small, irregularly shaped fields, and the lack of continuous crop canopy. With the spatial resolution of OSMI, it is still impossible to discriminate crops in Korea. But there are many schedules to launch commercial satellite such as IKONOS, Quick Bird, Orbview loaded with a high resolution and multispectral sensor within a or two years(Yeo, 1998). And also, many countries including America, Russia, Japan, Brazil, China, India, and Korea are involved in launching new satellites.

The study described herein evaluated for the feasibility of crop discrimination knowledge-

throughout each growing season at approximately weekly intervals. The spectroradiometer was elevated 50 cm above the crop canopies. Data were taken only when there were no clouds in the vicinity of the sun and when the solar elevation was at least 51°. Measurements of incident solar radiance and reflective radiance from canopies were made after the instrument was leveled for a nadir view angle. The spectroradiometer has a 60° field of view. Per cent spectral reflectance was calculated as the ratio of canopy radiance to the incident solar radiance.

3) Growth index induction of corn and data analysis

The index of phenological growth stage by Hanway(1971) was used for estimating developmental stage of a corn by using OSMI equivalent bands and their ratio values in vegetative stage. SAS statistics procedures were used for general linear models and regression models.

3. Results and Discussion

1) OSMI bands preparation with spectroradiometer channels

OSMI equivalent bands were selected in spectral wavebands obtained with spectroradiometer ranged from 330 to 1100 nm(5 nm interval), those are 445 nm, 490 nm, 510 nm, 555 nm, 670 nm, and 865 nm as bands 1, 2, 3, 4, 5, and 6, respectively. Band width of OSMI is 20 nm in bands 1, 2, 3, 4, and 5, and 40 nm in band 6. In this study, however, single bands were used for data analysis because data were taken with 5 nm interval spectroradiometer not an OSMI simulated sensor. Unpublished results showed no critical differences between single bands and averaged(20 nm) multibands(no data were shown in this paper). The values of averaged multibands were just smoother than those of single bands.

Ratio value(RV) 1, 2, 3, 4, and 5 were derived from the OSMI equivalent band ratios as band 6 : band 1, band 6 : band 2, band 6 : band 3, band 6 : band 4, and band 6 : band 5, respectively.

2) Time-series characteristics of spectral reflectance in corn, peanut, and soybean

Spectral reflectance profiles of corns, peanuts, and soybean based on day of year(DOY) are shown in Figure 2. In corns, since vegetative and reproductive stages are distinct and flowering period is short, corn stops vegetative growth soon after flowering. Spectral reflectances of OSMI

Characteristics of Spectral Reflectance for Corns and Legumes at OSMI(Ocean Scanning Multi-spectral Imager) Bands




Fig. 2. Spectral reflectance profiles at OSMI equivalent bands of corns, peanuts, and soybean based on day of year (DOY).

equivalent bands 1, 2, 3, 4, and 5 at visible wavelengths had a trend to decrease continually because canopy maintained greenness until harvest. But spectral reflectance of OSMI equivalent band 6 had a maximum value around flowering stage. And then it was decreased on maturing.

On the other hand, spectral reflectances of OSMI equivalent bands 1, 2, 3, 4, and 5 at visible wavelengths in peanuts and soybean were decreased as well. However, per cent reflectance values of OSMI equivalent band 6 in peanuts and soybean were increased even after flowering. That is unique signature of peanuts and soybean, although they have different growth type. Peanuts are indeterminate growth type while soybeans are determinate growth type.

In general, per cent reflectance value of OSMI equivalent bands 1, 2, 3, 4, and 5 were in upward order at the early stage of vegetative growth in corns, peanuts, and soybean. As crop grows, per cent reflectance value of OSMI equivalent band 5 was lower and lower to the lowest. What was interesting was that crossing point (per cent reflectance value of OSMI band 5 is lower than that of band 4, 3, 2, and 1 between bands of every crop) were different from their developmental stages. While this crossing point of band 4 and 5 was at 7 to 9 leaf stage far before flowering in corns, it was at flowering stage in soybean and after flowering stage in peanuts. This phenomenon is concerned with canopy cover and development, because chlorophyll has absorption peak around at 670 nm. And also, time difference of crossing points between spectral reflectance curves could be a parameter to discriminate crops knowledge-based on developmental stages.

3) Crop separability with OSMI equivalent bands at different dates

General linear model(GLM) procedures of SAS statistics program were carried out to find out the appropriate date for crop discrimination by using OSMI equivalent bands(Table 1) and their ratio values(Table 2). Corn and two other legumes were discriminated from band 1 and 5 in June 9, band 1, 2, 3, 5, and 6 in June 15, band 1, 2, 3, 4, 5, and 6 in June 22, and band 4 and 5 in June 29(Table 1). Peanut and soybean could not be separable from each other only by using spectral reflectance signatures measured during May to July.

Effective ratio values for discriminating corn from 2 other legumes were RV 1, 2, 3, and 4 in June 9, RV 2, 3, and 4 in June 15, RV 1, 2, 3, 4, 5, and 6 in June 22, and RV 4 in June 29(Table 2). Peanut and soybean could not be separable with each other by ratio values as well. We need more spectral data collected during flowering to maturing stage and another parameter to discriminate peanut and soybean including vinyl mulching and cropping pattern. Around June 22 was the best date for corn discrimination from 2 other legumes(no significances were shown in measurement dates other than June) because all OSMI equivalent bands and ratio values in June 22 were highly significant for corn separability.

Table 1. Crop separabilities by using spectral reflectances of OSMI equivalent bands.

June 9	B1 ¹⁾	B2	B3	B4	B5	B6	June 15	B1	B2	B3	B4	B5	B6
<i>Corn</i>	4.03 ^{b2)}	4.53 ^b	5.44 ^a	9.22 ^a	6.27 ^b	40.14 ^a	<i>Corn</i>	3.27 ^b	3.62 ^b	4.51 ^b	7.83 ^b	4.20 ^b	43.94 ^a
<i>Peanut</i>	8.16 ^a	9.24 ^a	10.44 ^a	14.30 ^a	17.28 ^a	27.67 ^a	<i>Peanut</i>	6.61 ^a	7.31 ^a	8.33 ^a	11.88 ^a	12.63 ^a	28.08 ^b
<i>Soybean</i>	6.31 ^{ab}	7.60 ^{ab}	9.07 ^a	12.67 ^a	16.96 ^a	24.77 ^a	<i>Soybean</i>	5.60 ^{ab}	6.61 ^a	7.65 ^a	11.38 ^{ab}	14.05 ^a	23.21 ^b
Mean	5.79	6.62	7.71	11.49	11.72	33.42	Mean	4.77	5.35	6.31	9.77	8.65	35.20
C.V.(%)	17.33	18.71	18.93	20.30	19.53	15.69	C.V.(%)	14.10	8.83	8.75	11.50	12.73	4.70
F value	10.34*	9.06	7.56	3.00	17.02*	5.02	F value	15.74*	40.94**	32.41**	8.99	49.52**	86.74**
June 22	B1	B2	B3	B4	B5	B6	June 29	B1	B2	B3	B4	B5	B6
<i>Corn</i>	3.59 ^b	3.48 ^b	4.26 ^b	7.03 ^b	3.46 ^b	50.46 ^a	<i>Corn</i>	3.07 ^b	2.93 ^b	3.81 ^b	6.42 ^b	3.35 ^b	50.70 ^a
<i>Peanut</i>	5.77 ^a	6.69 ^a	7.71 ^a	11.53 ^a	12.97 ^a	31.11 ^b	<i>Peanut</i>	4.60 ^{ab}	4.85 ^{ab}	5.92 ^{ab}	9.89 ^{ab}	8.33 ^a	42.42 ^a
<i>Soybean</i>	5.12 ^a	6.18 ^a	7.41 ^a	11.21 ^a	13.80 ^a	27.22 ^b	<i>Soybean</i>	4.96 ^a	5.28 ^a	6.75 ^a	11.46 ^a	10.69 ^a	36.51 ^a
Mean	4.57	5.00	5.94	9.23	8.35	40.14	Mean	3.90	3.96	5.00	8.42	6.23	45.57
C.V.(%)	5.20	7.36	3.99	4.21	9.67	5.47	C.V.(%)	12.92	16.21	13.73	11.95	13.78	15.54
F value	53.63**	51.83**	150.48**	95.94**	110.63**	67.41**	F value	8.20	7.84	9.53	12.62*	36.37**	1.80

1) : OSMI equivalent band

2) : Duncan's multiple range test ($\alpha=0.05$)

* 0.05>p>0.01, ** 0.01>p>0.001, *** p<0.001

Table 2. Crop separabilities by using ratio values between OSMI equivalent bands.

June 9	RV1 ¹⁾	RV2	RV3	RV4	RV5	June 15	RV1	RV2	RV3	RV4	RV5
<i>Corn</i>	10.06 ^{a2)}	9.07 ^a	7.51 ^a	4.43 ^a	6.76 ^a	<i>Corn</i>	14.15 ^a	12.36 ^a	9.91 ^a	5.74 ^a	10.98 ^a
<i>Peanut</i>	3.41 ^b	3.01 ^b	2.67 ^b	1.95 ^b	1.61 ^b	<i>Peanut</i>	4.25 ^a	3.84 ^b	3.37 ^b	2.37 ^b	2.24 ^a
<i>Soybean</i>	3.92 ^b	3.26 ^b	2.73 ^b	1.95 ^b	1.46 ^b	<i>Soybean</i>	4.14 ^a	3.51 ^b	3.03 ^b	2.04 ^b	1.65 ^a
Mean	6.82	6.08	5.10	3.19	4.16	Mean	9.18	8.05	6.59	4.00	6.51
C.V.(%)	12.74	21.42	17.12	14.28	35.23	C.V.(%)	38.85	21.58	22.18	23.65	41.10
F value	41.87**	15.84*	22.92*	22.35*	9.46	F value	5.82	18.55*	15.60*	10.20*	8.38
June 22	RV1	RV2	RV3	RV4	RV5	June 29	RV1	RV2	RV3	RV4	RV5
<i>Corn</i>	14.09 ^a	14.62 ^a	11.86 ^a	7.19 ^a	14.83 ^a	<i>Corn</i>	16.75 ^a	17.90 ^a	13.57 ^a	8.01 ^a	16.02 ^a
<i>Peanut</i>	5.40 ^b	4.67 ^b	4.04 ^b	2.71 ^b	2.42 ^b	<i>Peanut</i>	9.32 ^b	8.85 ^a	7.23 ^{ab}	4.31 ^{ab}	5.09 ^a
<i>Soybean</i>	5.32 ^b	4.40 ^b	3.67 ^b	2.43 ^b	1.97 ^b	<i>Soybean</i>	7.36 ^b	6.92 ^a	5.41 ^b	3.19 ^b	3.41 ^a
Mean	9.73	9.60	7.89	4.90	8.55	Mean	12.71	13.05	10.10	5.97	10.27
C.V.(%)	7.66	12.86	6.94	8.19	21.12	C.V.(%)	19.80	26.67	20.29	18.26	34.82
F value	102.60**	49.70**	158.27***	97.56**	36.32**	F value	7.94	5.92	8.90	10.84*	7.81

1) : RV1=Band6/Band1, RV2=Band6/Band2, RV3=Band6/Band3, RV4=Band6/Band4, RV5=Band6/Band5,

2) : Duncan's multiple range test ($\alpha=0.05$)

* 0.05>p>0.01, ** 0.01>p>0.001, *** p<0.001

4) Estimation of vegetative growth stage in corn variety(rs510) as a function of spectral reflectance

Hanway(1971) described the phenological growth stage of a corn plant as stage index from seeding(0.0) to physiological maturity(10.0). In this study, vegetative stage from seeding(0.0) to flowering(5.0) was examined for estimating developmental stage of a corn(rs510 variety) by using OSMI equivalent bands and their ratio values. Four kinds of variable set, which were OSMI equivalent bands(B), logB, ratio values(RV) of OSMI equivalent bands, and logRV, were prepared for multi-regression and stepwise-regression model. As results, five significant models, 2 multi-regression model with RVs(1, 2, 3, 4, and 5) and logRVs(1, 2, 3, 4, and 5) and 3 stepwise-regression model with B(1, 2, and 3), logB(1, 2, and 3), and RV(4 and 5), were obtained from the SAS procedures REG and STEPWISE(Table 3).

Table 3. Best growth stage prediction models generated by the SAS procedures REG and STEPWISE with OSMI equivalent bands and their band ratios.

Model name	R ²	Prob>F	Complete model
MR ¹⁾ 1	0.97	0.0044	-1.461 -0.073*RV ³⁾ 1 - 0.155*RV2 + 0.482*RV3 + 1.420*RV4 -0.493*RV5
MR 2	0.97	0.0039	-0.98 - 5.68*log ⁴⁾ RV1 - 16.48*logRV2 + 32.84*logRV3 + 5.61*logRV4 -7.10*logRV5
SR ²⁾ 1	0.94	0.0005	6.43 + 1.66*B ⁵⁾ 1 + 3.27*B2 -4.59*B3
SR 2	0.94	0.0004	12.18 + 10.20*logB1 + 23.06*logB2 -41.65*logB3
SR 3	0.97	0.0001	-2.000 + 2.066*RV4 -0.617*RV5

¹⁾ MR : Multi-regression model, ²⁾ SR : Stepwise-regression model

³⁾ RV1=Band6/Band1, RV2=Band6/Band2, RV3=Band6/Band3, RV4=Band6/Band4, RV5=Band6/Band5,

⁴⁾ log=log₁₀

⁵⁾ B : OSMI equivalent band

Corn(rs510) was about 2.5 meter tall at the end of vegetative stage. It almost stops its vegetative growing after flowering. It means no difference in plant height after flowering in corn while canopies of peanut and soybean are still growing even after flowering. Therefore, it is worth while estimating developmental stage of a corn(rs510) with OSMI equivalent bands and their ratio values at vegetative stage.

4. Conclusions

Time-series spectral reflectance data of major upland crops, corns, peanuts, and a soybean, were collected by using a spectroradiometer and analyzed for spectral separability assessment and estimation of vegetative growth stage in a corn depending on crop phenology.

Spectral reflectance behavior of OSMI equivalent band 5 had a unique characteristic, which was that crossing point of band 4 and 5 was at 7 to 9 leaf stage far before flowering in corns, it was at flowering stage in soybean and at after flowering in peanuts. This phenomenon is concerned with canopy cover and development, because chlorophyll has absorption peak around at 670 nm. And also, time difference of crossing points between spectral reflectance curves could be a parameter to discriminate the crops knowledge-based on developmental stages. Unpublished results showed no critical differences between single bands and averaged(20 nm) multibands(no data were shown in this paper). Values of averaged multibands were just smoother than those of single bands.

Corn was clearly discriminated from two other legumes, peanut and soybean in late June, according to the SAS procedure GLM with OSMI equivalent bands and their ratio values(RVs). And also, growth stage of a corn could be estimated as a function of OSMI equivalent bands and their ratio values in vegetative stage.

No remarkable difference of spectral reflectance curves was shown between peanut and soybean until July although their seeding and flowering periods were different with each other. It needed more study and spectral reflectance data to screen another parameters such as vinyl mulching and cropping pattern to discriminate peanut from soybean and to find out the appropriate time for classification.

This kind of approach to spectral data interpretation may be theoretical because it considered only spectral reflectance factor excluding cultivation unit of our country and spatial resolution of OSMI sensor. But we do say crop calendar and characteristics of spectral reflectance can be used as a strong ancillary and reference data for crop identification and area estimation, even though interpretation direction may be changed.

Acknowledgement

This paper was accepted to publish by the editorial committee of National Institute of Agricultural Science and Technology(NIAST, 98-1-2-60). Growth stages for corns and peanuts were examined by Dr. Hyeon-Gui Moon, Dr. Chang-Hwan Park, and Mr. Tae-Wook Jung working at National Crop Experimental Station, whose contribution is gratefully acknowledged.

References

- 秋山侃, 福原道一, 藤元也, 深山一, 1996. 農業リモートセンシング, 農林水産省農業環境技術研究所, p. 13.
- Panigrahy, S. and M. Chakraborty, 1998. An integrated approach for potato crop intensification using temporal remote sensing data. *ISPRS J. Photogrammetry & Remote Sensing*, 53:54-60.
- Janssen, L. L. F. and H. Middelkoop, 1992. Knowledge-based crop classification of a Landsat thematic mapper image. *ISPRS J. Photogrammetry & Remote Sensing*, 13:2827-2837.
- Yeo, W. S., 1998. Trends in satellite image market. Workshop on new technology and its application for satellite image processing in GIS. *Cadland Inc.* 1998. 7. 15. pp. 22-23.
- Hanway, J. J., 1971. *How a corn plant develops*. Special Report No. 48 Science and Technology Cooperative Extension Service. Iowa State University