

The Inhibitory Effect of Metronidazole and Doxycycline - HCl on proMMP - 3 Production in Gingival Fibroblast

Hak - Joo Kim · Ki - Jung Lim · Sang - Mok Kim · Byung - Ock Kim · Kyung - Yoon Han

Department of Periodontology, College of Dentistry, Chosun University

I. Introduction

Matrix metalloproteinases(MMP) are a family of proteolytic enzymes that mediate the degradation of extracellular matrix macromolecules, including interstitial and basement membrane collagens, fibronectin, laminin, and proteoglycan core protein. The major cell types(fibroblasts, keratinocytes, endothelial cells, and macrophages) in periodontal tissue are capable of responding to growth factors and cytokines, as well as to products released from the microbial flora by induction of transcription of one or more MMP genes¹.

These MMPs share some common properties: 1) secretion from the cell in a latent form(proenzyme) with its subsequent activation in the extracellular space; 2) containing zinc cation at the active site; 3) inhibition by chelators of calcium(e.g., EDTA), and zinc(e.g., 1, 10 phenanthroline); 4) inhibition by tissue inhibitors of metalloproteinases(TIMPs); and 5) degradation at least one component of the extracel-

lular matrix(e.g., collagen)². One prominent member of these MMPs, MMP - 3(stromelysin - 1) is capable of degrading the numerous extracellular matrix macromolecules(ECM) including fibronectin, laminin, proteoglycan core protein, collagen IV, V, IX, X, and elastin³.

Increase of MMP - 3 activity associated with several chronic inflammatory disease appear to be the result of specific inductive mechanisms. One of the mediators in induction of MMP - 3 is interleukin - 1(IL - 1), cell product that has important regulatory functions mediating the body's response to microbial invasion, inflammation, and tissue injury^{4,5,6,7}. Especially, IL - 1 is thought to play a important role in the tissue destruction associated with inflammatory diseases such as rheumatoid arthritis and periodontal disease^{8,9}. As periodontitis is specifically associated with the destruction of periodontal connective tissues, it is closely related to both IL - 1 and MMPs.

Gingivitis can trigger the initial cascade of

periodontal destruction. The human gingival fibroblast is prominent cell type in the gingival connective tissue and products cytokines induced by microbial infection in periodontal disease. In the periodontal disease, the upregulation of MMP expression in response to locally released IL - 1 may provide one component of this pathologic process. IL - 1 has been identified in both the gingiva and gingival crevicular fluids of periodontitis patients^{10,11}). In Vitro, fibroblasts derived from gingival tissue have shown enhanced production of both MMP - 1 (collagenase) and MMP - 3 with IL - 1 stimulation^{7,12}).

Tetracyclines (TCs) and their chemically modified analogues (CMTs) remain useful as antibiotics in periodontal therapy. TCs and CMTs also have non - antimicrobial properties which appear to modulate host response. For example, clinical studies indicated that tetracyclines may be useful in the treatment of certain medical conditions like epidermolysis bullosa, rosacea, alpha - antitrypsin - deficiency panniculitis, pyoderma gangrenosum and other inflammatory and bullous skin diseases which do not have a microbial etiology¹³). It has been proposed that the anti - inflammatory mechanisms are due in part to tetracycline's ability to inhibit leukocyte proliferation and activity and to scavenge hypochlorous acid and superoxide radicals produced by phagocytes¹⁴). In that aspect, TCs and CMTs have been shown to inhibit the activity of MMPs and collagenase^{15,16,17,18}). Also, metronidazole is widely used in the treatment of trichomoniasis, tropical ulcer, giardiasis, balantidiasis, amebiasis, dracun -

culiasis and acute necrotizing ulcerative gingivitis (ANUG). The previous study¹⁹) suggested metronidazole has a marked anti - inflammatory action, such as early subsidence of pain and inflammatory edema and healing of an ulcer.

In the present study, we investigated the inhibitory effects of metronidazole and doxycycline - HCl, one of the tetracycline analogues, on the proMMP - 3 level in human gingival fibroblast cells induced by IL - 1 .

II. Material and Method

1. Cell preparation and culture

Gingival fibroblasts were obtained from gingival connective tissue of a healthy adult with clinically and radiographically normal periodontal tissues. The tissue explants were resected and immersed immediately in Hanke's buffered salt solution (HBSS) (GIBCO/BRL, USA) containing antibiotics (penicillin: 1000 U/ml, Streptomycin: 1000 µg/ml, and fungizon: 50 µg/ml) (GIBCO/BRL, USA). The tissue samples were rinsed several times with the same medium and minced to 1 × 1 × 1 mm. The minced samples were placed in 60 mm culture plates (NUNC, Netherland) containing culture medium composed of Dulbecco's Modified Eagle's Medium (DMEM) (GIBCO/BRL, USA) with penicillin: 1000 U/ml, Streptomycin: 1000 µg/ml, and fungi - zon: 50 µg/ml, and 20% heat - inactivated fetal bovine serum (FBS) (GIBCO/BRL, USA). The cells were incubated at 37 in a humidified atmosphere of 95% air and 5% CO₂ for 2 weeks. The medium was

renewed every 3 or 4 days until the cells were confluent. After 0.25% trypsin/EDTA(GIBCO/BRL, USA) incubation for detachment, the cells were transferred into 90 mm tissue culture plates. The cells were maintained in culture medium with 10% FBS and passaged upon confluence. The cells between the fifth and tenth passages showing a fibroblast-like morphology under light microscopy were used for the experiment.

2. Treatment with metronidazole and doxycycline - HCl

Dense cultures of fibroblasts were treated with 0.25% trypsin. The detached cells were washed once with DMEM supplemented with 10% FBS and diluted to 4×10^4 cells/ml in DMEM supplemented with 10% FBS. A cell suspension of cells was distributed into each well of 6 - well plate dish. When the cells reached confluence, the medium was replaced with DMEM supplemented with 0.1% FBS. The cells were then treated with culture medium containing 0.1% FBS alone (as a control) or with increasing concentrations (10, 25, 50, 100, 200 $\mu\text{g/ml}$) of metronidazole (Sigma, USA)

and doxycycline - HCl (Sigma, USA) for an hour prior to adding the recombinant human IL - 1 .

3. Treatment with the recombinant human IL - 1

After the incubation, the cell cultures were treated with the optimal concentration (25ng/ml) of recombinant human IL - 1 (R & D systems, Minneapolis, MN) and incubated for 24 hours prior to ELISA.

4. Enzyme - linked immunosorbent assay (ELISA)

After the incubation for 24 hours with antibiotics and IL - 1 , cell culture supernatant was obtained, and diluted with sample diluent according to recommended dilution rate (1/21). ELISA procedure was performed according to the manual of proMMP - 3 ELISA kit (The Binding site, San Diego, CA). Briefly, control or diluted 100 μl samples were distributed to the microwell plate and incubated for 60 minutes. After incubation, wells were washed 3 times with washing buffer. The biotinylated antibody 100 μl was added to each well and

Table 1. The optical density of proMMP - 3 according to the concentration of metronidazole

Concentration ($\mu\text{g/ml}$)	Number	Optical density (Mean \pm S.D)
0 (control)	6	1.80 \pm 0.54
10	6	0.42 \pm 0.08*
25	6	0.41 \pm 0.07*
50	6	0.74 \pm 0.20*
100	6	0.72 \pm 0.08*
200	6	1.10 \pm 0.18*

*p<0.05: significantly different from the control




Figure 1. The expression of proMMP - 3 according to the concentration of metronidazole
O.D.: Optical Density

incubated for 60 minutes and washed. The streptavidin peroxidase 100µl was added to each well and incubated for 30 minutes and washed. The substrate 100µl was added to each well and incubated for 10 minutes.

Finally, The stopping solution 100µl was added to each well, and the optical density was measured at 450nm under the microwell plate reader (Bio - Tek instrument, USA). Each assay was carried out in

Table 2. The optical density of proMMP - 3 according to the concentration of doxycycline - HCl

Concentration(µg/ml)	Number	Optical density (Mean ±S.D)
0(control)	6	1.26 ± 0.12
10	6	0.52 ± 0.03*
25	6	0.67 ± 0.06*
50	6	0.73 ± 0.09*
100	6	1.00 ± 0.05*
200	6	1.45 ± 0.10*

*p<0.05: significantly different from the control




Figure 2. The expression of proMMP - 3 according to the concentraion of doxycycline - HCl
O.D.: Optical Density

triplicate.

5. Statistical analysis

The difference of obtained data between groups was statistically analyzed by independent t - test, ANOVA and Duncan test. And the percentile change from the control was obtained in the each concentration of metronidazole and doxycycline - HCl.

III. Result

1. proMMP - 3 expression inhibited by varying the concentrations of metronidazole

When the cells were incubated with 10, 25, 50, 100, 200 μ g/ml of metronidazole and IL - 1 , proMMP - 3 expression was significantly decreased in all of the concentration, compared with control without metronidazole (Table 1). The proMMP - 3 expression exhibited initial reduction and gradual increase (Figure 1).

Table 3. The percentile change of proMMP - 3 by varying the concentration of metronidazole and doxycycline - HCl

Concentration(μ g/ml)	Metronidazole	Doxy - HCl
10	23.89%	41.78%
25	23.45%	53.06%
50	47.18%	58.10%
100	44.45%	80.04%
200	63.89%	115.90%




Figure 3. The comparison between metronidazole and doxycycline - HCl in the inhibitory effect on proMMP - 3 production

2. proMMP - 3 expression inhibited by varying the concentrations of doxycycline - HCl

When the cells were incubated with 10, 25, 50, 100, 200 $\mu\text{g}/\text{ml}$ of doxycycline - HCl and IL - 1 , proMMP - 3 expression was significantly decreased in the concentration below 100 $\mu\text{g}/\text{ml}$, compared with the control without doxycycline - HCl (Table 2). But proMMP - 3 expression was significantly increased in the cells incubated with 200 $\mu\text{g}/\text{ml}$ of doxycycline - HCl. The proMMP - 3 expression exhibited initial reduction and gradual increase (Figure 2).

3. Percentile change of proMMP - 3 expression from the control

When the percentile changes of proMMP - 3 expression between doxycycline - HCl and metronidazole were compared, it was noted that metronidazole was superior to doxycycline - HCl in inhibitory effect (Table 3, Figure 3).

IV. Discussion

IL - 1 produced by the macrophages/monocytes has multiple biological activities and is important in the immune and inflammatory responses. Especially, IL - 1 has been detected in gingival crevicular fluid and the gingival tissue affected by periodontitis. Several studies have shown that IL - 1 stimulates the production of collagenase, and the mitogenesis of gingival fibroblasts.

The biological activities of IL - 1 are

mediated by the binding of specific cell surface receptors. Two types of IL - 1 receptors (IL - 1R) have been identified: type I is expressed on T cells, epithelial cells, and fibroblasts, and type II is expressed on B cells, macrophages, and myeloid cells. IL - 1's actions may be determined by the type of receptors expressed on different types of cells.

Various reagents and cytokines have been found to influence IL - 1R expression in various cell types. PDGF, PGE₂, and cAMP - generating reagents up - regulate IL - 1R expression in fibroblasts. It has been found that IL - 1 itself rapidly up - regulates IL - 1R expression in fibroblast^(20,21). Chieko et al.⁽²²⁾ showed that IL - 1R expression on the inflamed gingival fibroblast cells stimulated by IL - 1 continuously increased. In the previous study, proMMP - 3 expression from gingival fibroblast stimulated by IL - 1 showed increasing trend corresponding to the IL - 1 concentration. Especially, in the concentration over 25ng/ml, proMMP - 3 was significantly increased⁽²³⁾.

Previous investigation⁽¹⁾ of IL - 1 regulation of MMP - 3 expression in periodontal ligament fibroblasts supported its role in the destruction of periodontal tissue associated with periodontitis. Because periodontitis is clinically distinct from gingivitis, there is necessarily some differences on a cellular level. Birkedal - Hansen⁽²⁴⁾ summarized the key understanding why gingival inflammation may or may not give rise to tissue destruction and attachment loss as follows: 1) different cell types express different complements of MMP; 2) different cytokines elicit different transcriptional

effects for MMP genes; and 3) different cell types do not necessarily respond in the same fashion to a given cytokine. Present results didn't provide a evidence for the direct effect of the IL - 1 in the periodontal ligament. But, as regarding the triggering role of gingival connective tissue inflammation in the periodontal disease, the present study provided another indirect evidence for IL - 1 regulation of MMP - 3 expression in periodontitis.

In the early to mid - 1980s, Golub and co-workers¹⁷⁾ reported that: 1) tetracyclines could inhibit the activity of mammalian collagenase; and 2) this inhibition was unrelated to the antimicrobial efficacy of these drugs. In fact, the initial observation of tetracyclines' effect on mammalian collagenase has been extended to other metalloproteinases including gelatinase, type IV/V collagenase, and macrophage elastase. Tetracyclines are known to inhibit collagenases from a variety of cells: neutrophils, macrophages, osteoblasts, chondrocytes, and a wide range of tissues: skin, gingiva, cornea, cartilage, and rheumatoid synovium.

Barry et al.²⁾ summarized that the ability of TCs may 1) indirectly inhibit the activity of extracellular collagenase and other MMPs such as gelatinase; 2) prevent the activation of its proenzyme by scavenging reactive oxygen species generated by other cell types (e.g., PMNs, osteoclasts); 3) inhibit the secretion of other collagenolytic enzymes (e.g., lysosomal cathepsins); and 4) directly affect other aspects of osteoclast structure and function. Several studies have also documented the therapeutic potential of TCs and CMTs in periodontal disease. In

our study, antibiotics were treated before the IL - 1 application, and then the inhibition of MMP - 3 was quantitatively measured in the pro - enzyme level. The results suggest that MMP - 3 expression can be inhibited by the antibiotics before the protein synthesis step.

Human fibroblast collagenase (MMP - 1) appears to be relatively resistant to tetracyclines treatment²⁵⁾. Since the neutrophils (PMN) likely provide the principal source of collagenase for tissue destruction during periodontal disease, fibroblast collagenase may be required for normal connective tissue remodeling. So, Barry et al.^{10,25)} suggested that the differential sensitivity of PMN and fibroblast collagenases to tetracycline treatment may have substantial therapeutic benefits. In this aspect, pharmacologic concentrations might inhibit the activity of PMN, but not the fibroblast enzyme. Such a selective inhibition might reduce collagenolytic activity and tissue destruction during inflammation, but not the normal collagen turnover required for the maintenance of tissue integrity. Because the present study was only performed for proMMP - 3 expression in the gingival fibroblast, we could not compare the differences between the inhibitory effect of tetracycline and CMTs in gingival fibroblast and PMNs. But it is distinct that MMP - 3 originated from gingival fibroblast can play a role in the inflammatory reaction induced by IL - 1 of inflammatory cells (PMN, monocyte, macrophage).

Metronidazole and doxycycline - HCl have been used because of not only the antimicrobial property, but also the anti - inflam -

matory property. Burns et al.²⁶⁾ compared the relative potencies of different collagenase inhibitors including tetracyclines, and reported that doxycycline (with an $IC_{50}=15\ \mu\text{M}$) was a more potent collagenase inhibitor than two other tetracyclines, minocycline ($IC_{50}=190\ \mu\text{M}$) and tetracycline ($IC_{50}=15\ \mu\text{M}$). In the safety aspect of a low-dose regimen of these CMTs, Golub et al.^{16,27)} described the effectiveness of a low-dose regimen of minocycline in reducing GCF collagenase activity in the periodontal pocket although no significant changes in the make up of the crevicular microflora were detected. In addition, their other study suggested that subgingival plaque microorganisms (*Fusobacterium nucleatum*, *Actinomyces* spp, *Bacteroides* spp) do not develop tetracycline resistance when patients are administered a 2 weeks regimen of low dose doxycycline capsules. In contrast, when patients were administered regular-dose doxycycline capsules (50mg. b.i.d), the subgingival plaque appeared to become resistant (minimum inhibitory concentration of doxycycline = 25 - 100 $\mu\text{g}/\text{ml}$ in these subjects) to expected tissue fluid levels (1 - 5 $\mu\text{g}/\text{ml}$) of this antibiotic²⁸⁾. Golub et al.^{16,29)} suggested that a regimen of low-dose CMTs capsules may provide a safe and effective adjunct to instrumentation therapy without inducing the side effects (e.g., antibiotic-resistant microorganisms, gastro-intestinal upset) in the management of pathologic collagenolysis in the periodontal patient. In the present study, when the reduction of MMP-3 expression by metronidazole and doxycycline-HCl was compared through this

uncontrolled study, the more potent MMP-3 inhibitor was the metronidazole. The metronidazole showed the inhibitory effect in the all of concentration range. Especially, in comparing with the percentile change of doxycycline-HCl, the inhibitory effect of metronidazole was reached to about 2-fold in each concentration. This result suggests that metronidazole may be suitable for the periodontal treatment without any induction of antibiotic-resistant organism and for a long term therapy.

Although the in vivo activities of MMP-3 are still uncertain, the results suggest that low concentration of metronidazole and doxycycline-HCl can inhibit the activity of MMP-3 without antibiotic-resistance in gingival connective tissue with gingivitis and periodontitis. Further studies, regarding the drug concentration in gingival crevice, drug delivery system, and recommended dosage for control of periodontal inflammation, are required.

V. Conclusion

The purpose of the present study was to investigate the inhibitory effect of metronidazole and doxycycline-HCl on proMMP-3 production in the activated gingival fibroblast by IL-1. The cultured human gingival fibroblasts (4×10^4 cell/ml) were treated with metronidazole and doxycycline-HCl at various concentrations (10 - 200 $\mu\text{g}/\text{ml}$) incubated for 1 hour, and treated with 25ng/ml of IL-1 to induce the MMP-3. The proMMP-3 level was assayed with proMMP-3 ELISA Kit. The following results were obtained through the in vitro

study

1. Metronidazole inhibited significantly the expression of proMMP - 3 in wide range of concentration from 10 to 200 $\mu\text{g/ml}$ ($p < 0.05$).
2. Doxycycline - HCl inhibited significantly the expression of proMMP - 3 at the concentration lower than 100 $\mu\text{g/ml}$, but increased significantly the expression of proMMP - 3 at the concentration of 200 $\mu\text{g/ml}$ ($p < 0.05$).
3. The percentile change of the proMMP - 3 expression was more reduced at metronidazole treated groups than doxycycline - HCl treated groups.

The results showed that the low concentration of doxycycline - HCl ($< 100 \mu\text{g/ml}$) and metronidazole ($< 200 \mu\text{g/ml}$) could inhibit effectively the activity of MMP - 3 induced by IL - 1 in gingival fibroblasts, and suggest that metronidazole and doxycycline - HCl may play important roles in anti - inflammatory effect as well as in antibiotic effect, and metronidazole may be superior to doxycycline - HCl in inhibitory effect.

VI. References

1. Nip, L.H., Uitto, V - J., Golub, L.M.: " Inhibition of epithelial cell matrix metalloproteinases by tetracyclines ", J. Periodont. Res., 28 : 379 - 385, 1993.
2. Rifkin, B.R., Vernillo, T.V., Golub, L.M.: " Blocking periodontal disease progression by inhibiting tissue destructive enzymes: a potential therapeutic role for tetracyclines and their chemically - modified analogs ", J. Periodontol., 64 : 819 - 827, 1993.
3. Preus, H., Tollefsen, T., M rland, B.: " Effect of tetracycline on human monocyte phagocytosis and lymphocyte proliferation ", Acta. Odontol. Scand., 45 : 297 - 302, 1987.
4. Conca, W., Kaplan, P.B., Krane, S.M.: " Increase in levels of procollagenase mRNA in cultured fibroblasts induced by IL - 1 or serum follow c - jun expression and are dependent on new protein synthesis ", J. Clin. Invest., 83 : 1753 - 1757, 1989.
5. Frisch, S.M., Ruley, H.E.: " Transcription of the stromelysin promoter is induced by IL - 1 and repressed by dexamethasone ", J. Biol. Chem., 262 : 16300 - 16304, 1987.
6. Ganu, V.S., Hu, S - I., Melton, R., et al.: " Biochemical and molecular characterization of stromelysin synthesized by human osteoarthritic chondrocytes stimulated with recombinant human interleukin - 1 ", Clin. Exp. Rheumatol., 12 : 489 - 496, 1994.
7. Richards, D., Rutherford, R.B.: " Interleukin - 1 regulation of procollagenase mRNA and protein in periodontal fibroblasts in vitro ", J. Periodont. Res., 25 : 222 - 229, 1990.
8. McCulloch, C.A.G., Birek, P., Aitken, S., Lee, W.: " Low - dose tetracyclines decrease elastase and B - glucuronidase activities in gingival crevicular fluid ", J. Dent. Res., 69(Spec, issue) : AADR abst(submitted), 1990.
9. Tatakis, D.S.: " Interleukin - 1 and bone metabolism: A review ", J. periodontol.,

- 64 : 416 - 431, 1993.
10. Sorsa, T., Uitto, V - J., Suomolainen, M., Vauhkonen, M., Lindy, S.: " Comparison of interstitial collage - nases from human gingiva, sulcular fluid and polymorphonuclear leukocytes "; J. Periodont. Res., 23 : 386 - 393, 1988.
11. Stashenko, P., Fujiyoshi, P.,

- Obernesser, M.S., Prostack, L., Haffajee, A.D., Socransky, S.S.: " Levels of interleukin - 1 in tissue from sites of active periodontitis disease "; *J. Clin. Periodontol.*, 18 : 548 - 554, 1991.
12. Tewari, D.S., Qian, Y., Tewari, M., et al.: " Mechanistic features associated with induction of metalloproteinases in human gingival fibroblasts by interleukin - 1 "; *Arch. Oral. Biol.*, 39 : 657 - 664, 1994.
 13. Okada, Y.N., Nagase, H., Harris, E.D. Jr.: " A metalloproteinase from human rheumatoid synovial fibroblasts that digests connective tissue matrix components "; *J. Biol. Chem.*, 261 : 14245 - 14255, 1986.
 14. Richards, D., Rutherford, R.B.: " Interleukin - 1 regulation of procollagenase mRNA and Protein in periodontal fibroblasts in vitro "; *J. Periodont. Res.*, 25 : 222 - 229, 1990.
 15. Golub, L.M., Mcnamara, T.F., D'Angelo G., Greenwald, R.A., Ramamurthy, S.: " A nonantibacterial chemically - modified tetracycline inhibits mammalian collagenase activity "; *J. Dent. Res.*, 66 : 1310 - 1314, 1987.
 16. Golub, L.M., Lee, H.M., Lehrer, G., et al.: " Minocycline reduces gingival collagenolytic activity during diabetes: preliminary observations and a proposed new mechanism of action "; *J. Periodontal. Res.*, 18 : 516 - 526, 1983.
 17. Golub, L.M., Ramamurthy, N.S., McNamara, T.F., et al.: " Tetracycline inhibits tissue collagenase activity: a new mechanism in the treatment of periodontal disease "; *J. Periodont. Res.*, 19 : 651 - 655, 1984.
 18. Golub, L.M., Ramamurthy, N.S., McNamara, T.F., Greenwald, R.A., Rifkin, B.R.: " Tetracyclines inhibit connective tissue breakdown: new therapeutic applications for an old family of drugs "; *Critical Reviews in Oral Biology and Medicine.*, 2 : 297 - 322, 1991.
 19. Tanga, M.R., Antani, J.A., Kabade, S.S.: " Clinical evaluation of metronidazole as an anti - inflammatory agent "; *Int. Surgery.*, 60 : 75 - 76, 1975.
 20. Bonin, P.D., Chiou, W.J., McGee, J.E., Singh, J.E.: " Two signal transduction pathways mediate interleukin - 1 receptor expression in balb/c3T3 fibroblasts "; *J. Biol. Chem.*, 265 : 18643 - 18649, 1990.
 21. Bonin, P.D., Singh, J.P.: " Modulation of interleukin 1 receptor expression and interleukin - 1 response in fibroblasts by platelet - derived growth factor "; *J. Biol. Chem.*, 263 : 11052 - 11055, 1988.
 22. Chieko, K - N., Yuichi, I., Takeshi, S.: " Increased expression of interleukin - 1 receptors on fibroblasts derived from inflamed gingiva "; *J. Periodontol.*, 67 : 1267 - 1273, 1996.
 23. , , : " MMP - 3 "; 29 (4) : 617 - 631, 1999.
 24. Birkedal - Hansen H.: " Role of matrix metalloproteinases in human periodontal disease "; *J. Periodontol.*, 64:474 - 484, 1993.
 25. Sorsa, T., Ingman, T., Suomalainen, K., et al.: " Cellular source and tetracycline -

inhibition of gingival crevicular fluid collagenase in patients with labile diabetes ”, J. Clin. Periodontol., 19 : 146 - 149, 1992.

26. Burns, F.R., Stack, M.S., Gray, R.D., Paterson, C.A.: “ Inhibition of purified collagenase from alkaline - burned rabbit corneas ”, Invest. Ophthalmol. Vis. Sci., 30 : 1569 - 1575, 1989.
27. Golub, L.M., Wolff, M., Vidal, A.M., Ramamurthy, N.S., McNamara, T.F.: “ Low - dose minocycline therapy: effects on crevicular fluid collagenase and subgingival microflora ”, IADR abstract. J. Dent. Res., 66(Spec. Issue) : 384, 1987.
28. Nakaya, H., Oates, T.W., Hoang, A.H., Kamoi, K., Cochran, D.L.: “ Effect of interleukin - 1 on matrix metalloproteinase - 3 levels in human periodontal ligament cells ”, J. Periodontol., 68 : 517 - 523, 1997.
29. H nig, J., Rordorf - Adam, C., Seigmund, C., Wiedemann, W., Erard, F.: “ Increased interleukin - 1 beta(IL - 1) concentration in gingival tissue from periodontitis patients ”, J. Periodontol. Res., 24 : 230 - 242, 1991.

3 proMMP -
metronidazole
doxycycline - HCl

matrix metalloproteinase - 3(MMP - 3)
interleukin - 1 (IL - 1)
MMP -
3 tetracycline tetracycline
metronidazole doxy -
cycline - HCl
IL - 1 MMP - 3
MMP - 3
2
metronidazole (10 - 200µg/ml)
doxycycline - HCl(10 - 200µg/ml)
1 proMMP - 3
25ng/ml IL - 1
24
proMMP - 3 ELISA kit
t - test
(ANOVA) Duncan test
1. Metronidazole 10 - 200µg/ml
proMMP - 3
가 (p<0.05).
2. Doxycycline - HCl 100µg/ml
proMMP - 3
가 (p<0.05), 200µg/ml
proMMP - 3 가
(p<0.05).
3. Metronidazole doxycycline - HCl

proMMP - 3

metronidazole doxycy -
cline - HCl .

metronidazole(10 - 200
 $\mu\text{g/ml}$) doxycycline - HCl(100 $\mu\text{g/ml}$)
IL - 1
MMP - 3
.