

Regeneration and selection of root resistant *Coleus forskohlii* A threatened medicinal plant

Manju M.George, R.B.Subramanian and Hiren A.Prajapati

Department of Biosciences, Sardar Patel University

Vallabh Vidyanagar, Gujarat, INDIA - 388 120

ABSTRACT

Coleus forskohlii Briq. of the family Lamiaceae yields a valuable secondary metabolite known as forskolin which is a labdane diterpenoid. *Coleus forskohlii* is the only known source of this compound. Forskolin is used in medicine for the treatment of glaucoma, congestive cardiomyopathy and asthma.

Morphogenic callus was induced from young leaves on MS medium augmented with NAA and BA. These calli, when subcultured on MS with KN alone gave rise to shoots. The regenerated shoot developed good root system on MS medium fortified with NAA. The fully grown plantlets were transferred to soil for acclimatization.

Coleus plant is mainly infected by a fungi *Lasiodiplodia theobromae* which causes root rot disease. The fungal culture filtrate (FCF) of *Lasiodiplodia theobromae*, has been used in regeneration media to find the MIC and further to select resistant plants to the pathogen. In the present study 40% FCF in the medium showed maximum inhibition and is therefore considered as the MIC level of *Coleus forskohlii*. This data could prove to be useful for the future for selecting a resistant *C.forskohlii* plant against the root disease caused by *L.theobromae*.

INTRODUCTION

The demand for the plant based herbal medicines is increasing every day due to their fewer side effects in comparison to synthetic chemical drugs. It is therefore very important to conserve and propagate medicinal plants of any country.

Throughout history, plants have been the most important source of medicines for human health. About 1,00,000 compounds are now known from the plants with about, 4,000 new ones being discovered every year (Verpoorte et al, 1999)

Medicinal and aromatic plants form an important

group among the economic plants. Due to the increasing population, especially in the developing countries, many natural growing sites and ecosystems of important medicinal and aromatic plants are being destroyed. It is therefore urgent to chalk out conservation measures for these invaluable natural resources.

Many modern medicines are either derived from the plants or artificially synthesized to copy plant chemical compounds. The World Health organization [WHO] estimates that up to 80% of the world's population rely on plants for their primary health care. India is one of the 12 mega biodiversity centers where almost all known medicinal plants can be cultivated in some part

Corresponding author : **Manju M.George** Department of Biosciences, Sardar Patel University Vallabh Vidyanagar, Gujarat, INDIA - 388 120

of the country or the other.

India has about 2,000 species of medicinal plants and vast geographical area with high production potential and varied agro climatic condition. Cultivation of medicinal plants offers considerable scope for rural employment and export for foreign exchange earnings.

India is already a major exporter of medicinal plants. The turn over of herbal medicines in India as over the counter products, ethical and classical formulations and home remedies of Ayurveda, Unani, and Siddha systems of medicine is about 1 million US \$ with an export of about 80 million US \$ (Khamboj, 2000)

Coleus forskohlii Briq. of the family Lamiaceae, is an aromatic herb 30-60 cm. high with dull orange tuberous roots. It grows wild in the sub-tropical Himalayas, distributed from the Kumaon hills to Nepal ascending up to 2000M, and in Bihar and Gujarat. (Anonymous, 1950). It is cultivated in Gujarat and Karnataka. The plant yields a valuable secondary metabolite known as forskolin which is a labdane diterpenoid. Though forskolin is found in all parts of the plant, roots are the main source of compounds. *Coleus forskohlii* is the only known source of the compound.

Forskolin is used in medicine for the treatment of Glaucoma, Congestive cardiomyopathy and Asthma (Valdes et al. 1987). It is characterized by hypertensive property, inhibitory action on thrombocyte aggregation and reduction of intra ocular pressure. In addition forskolin is used in purification of adenylate cyclase and in receptor binding assays (SukhDev, 1997:25)

Indiscriminate and large-scale collection of this plant from forest and the other natural sources and insufficient attempts to either allow its replenishment or its cultivation, have led to the rapid depletion of its resource. Gupta (1998) has listed this plant as one of the plant species in India vulnerable to extinction. Moreover this plant is affected by root rot disease caused by a fungus *Lasiodiplodia theobromae*. In severe case the entire plant dries up.

Lasiodiplodia theobromae belongs to the class-Deuteromycotina, Order-Spheropsidales, and Family-Sphereopsidaceae. The fungus has a wide host range around the world. The fungus is found throughout the tropics and sub-tropics causing numerous kinds of disease, especially rots of fruits and root crops during storage such as, stem and root rot in citrus, mango fruit rot in papaya (Gupta and Nema, 1979), crown rot in banana, storage rot in sweet potato and also attacks root crops of cassava. The fungus causes mycotic keratitis and oncomycosis in animals (Laverde et al, 1973 McGinnis 1985).

It forms pycnidia up to 5 mm in diameter. Conidia are initially hyaline and unicellular, sub ovoid to ellipsoidal with granular content. Mature conidia are two celled, cinnamon or light brown in color and often with longitudinal striations. Conidia are $18-30 \times 10-15$ micrometer with paraphyses up to 50 micrometer long. associated with a large number of rots (Sulton, 1980).

It is evident from the foregoing account that developing a method for the mass propagation of this economically useful, endangered plant species is highly essential. *In vitro* micropropagation technique offers a powerful tool for plant germplasm conservation as well as obtaining disease resistant plants.

In any selection scheme, it is desirable that the trait of interest be selectable at cellular level and also expressed in the regenerated plants. In the past *in vitro* techniques have been employed successfully in the selection of resistant plant at cellular level (Jones, 1990). The basis of such selection is that some agents of known importance in the disease reaction should be employed as the selection criterion. The selection agents that can be used for selection can be (i) pathogen itself or (ii) pathogen metabolites.

Use of pathogen metabolite for selection is the most commonly employed method and seems to be reliable in obtaining disease resistant plants.

Selection has been carried out with callus, or cell

suspension cultures and also protoplast. Culture filtrates represents an easy, common and cheap method of selection by incorporation in the culture media at appropriate concentrations. In several selection experiments culture filtrates, both purified and partially purified have been successfully used. For many diseases, well-characterized toxins are not known, and culture filtrates in these exhibit phytotoxic activities (Larkin and Scowcraft, 1983).

The pathogen causing root rot disease, *Lasiodiplodia theobromae* has been isolated and identified (MTCC 3086). Establishment of host-pathogen interaction *in vitro* permits screening and selection for disease resistant at cellular level (Vidyasekaran, 1993). A number of workers have used fungal culture filtrate (FCF) as the selection agent, rather than purified toxin to select resistant plants through tissue culture. This approach has been adapted in the present study also. The FCF of *Lasiodiplodia theobromae* has been used in callus induction and regeneration media to select resistant plants to the pathogen.

The scope of the project undertaken includes development of reliable, efficient *in vitro* propagation methods for *Coleus forskohlii* plant as well as the establishment of the Maximum Inhibitory Concentration (MIC) for the fungal toxin. This information will be highly useful for mass cultivation of this threatened, medicinally useful plant as well as for obtaining disease free/resistant plant.

MATERIALS AND METHODS

C. forskohlii plants were grown in Sardar Patel University botanical Garden, Vallabh Vidyanagar. Young leaves were taken from field grown mature plants.

The leaves were cut in to 1cm and were cultured in culture tube. The explants were sterilised using 0.1%

HgCl₂ for 3 minutes and thoroughly washed with sterile distilled water five times. For all the experiments, Murashige and Skoog (1962) medium supplemented with 3% sucrose (Hi Media, India) was used. The pH of the medium was adjusted to 5.8 followed by the addition of 0.8% agar (Hi Media, India) before autoclaving at 15 psi for 15-20 minutes (1230C). For callus induction and regeneration studies MS medium along with 2, 4-dichlorophenoxy acetic acid (2, 4-D), 1-naphthalene acetic acid (NAA), 3-indole butyric acid (IBA), 6-benzyl aminopurine (BA) and 6-furfuryl aminopurine (KN) were used as the growth hormones at varying concentrations. For shooting 150-200 mg green calli were subcultured on various concentration of cytokinins alone or in combination with auxins. Cultures were raised in 15 mm × 150 mm culture tubes under cool white fluorescent tubes emitting 50 mole m⁻² s⁻¹ for 14 hrs of photoperiod in a growth room maintained at 2520C. For hardening, the healthy plants were transferred to pro-trays containing sterile coco-peat kept in a small poly house to maintain humidity (80-85%RH) for 15 days and later transferred to soil for acclimatization.

Isolation of the fungus

The fungi *Lasiodiplodia theobromae* was isolated from decaying roots of *Coleus forskohlii* on potato dextrose Agar (PDA) medium and identified at IMTECH, Chandigarh (MTCC 3086). The pure cultures were transferred to PDA slants for subsequent maintenance of culture and bioassay. The fungus was grown on PD broth and the fungal culture filtrate was assayed for inhibitory/promotory effect by seed germination assay. The promotory effect of FCF from fresh subcultures was reconfirmed by wheat coleoptile elongation test.

The Maximum inhibitory concentration (MIC) of the FCF to *Coleus forskohlii* culture:-

The culture of *Lasiodiplodia theobromae* was

inoculated in 1L flask having 250ml PD broth (Himedia, Mumbai) and inoculated at 28°C for 10 days. The FCF was first filtered with Whatman's filter paper and further filtered through a 0.45 µm nylon membrane. The filtered crude toxin was added at different concentration of MS medium supplemented with 1 mg/l BA, (viz. 0%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%). The medium was then autoclaved at 121°C, 15 lbs. pressure for 15 minutes.

Calli weighing (~250 mg) were inoculated in these tubes and incubated in the growth room at 24°C with 14 hours of photoperiods using cool white fluorescent lamps of 3000 lux.

RESULTS

MS medium augmented with IBA+BA, IBA+KN, IBA+BA, NAA+KN at varying concentration gave poor response with the leaf explants (Table. 1) Out of the various combinations tried for callus induction with young leaf segments as explants. MS medium supplemented with NAA and BA showed better results. (Table.2)

The explants were placed in the medium with the abaxial and adaxial sides in contact with the medium. These explants placed with the abaxial surface in contact with the medium responded better. After 7 days of culture, the laminar portion as well as the petiolar portion showed callus initiation (Plate 1A).

MS medium augmented with 1-2 mg/l NAA +1mg/l BA gave better results than other combination of BA and NAA (Table.2)

Table 1. Media combinations tried for callus introduction from leaf disc explants of *C.forskohlii*

Media combinations	Response
MS+0.5mg/l IBA+1mg/l BA	+
MS+1mg/l IBA+1mg/l BA	+
MS+1mg/l IBA+0.5mg/l BA	+
MS+0.5mg/l IBA+1mg/l KN	-
MS+1mg/l IBA+1mg/l KN	-
MS+1mg/l IBA+0.5mg/l KN	-
MS+0.5mg/l IAA+1mg/l BA	+
MS+1mg/l IAA+1mg/l BA	+
MS+1mg/l IAA+0.5mg/l BA	+
MS+0.5mg/l IAA+1mg/l KN	-
MS+1mg/l IAA+1mg/l KN	-
MS+1mg/l IAA+0.5mg/l KN	-
MS+0.5mg/l NAA+1mg/l BA	+++
MS+1mg/l NAA+1mg/l BA	+++
MS+1mg/l NAA+0.5mg/l BA	+++
MS+0.5mg/l NAA+1mg/l KN	-
MS+1mg/l NAA+1mg/l KN	-
MS+1mg/l NAA+0.5mg/l KN	-

- no response

+ poor response

++ moderate

+++ good

Table 2. Media combinations tried for callus introduction from leaf explants of *C.forskohlii*

Media combinations	Response	%Reponse
MS+0.5mg/1 NAA+0.5mg/1 BAP	+	16
MS+1mg/1 NAA+0.5mg/1 BAP	+	20
MS+1.5mg/1 NAA+0.5mg/1 BAP	+	20
MS+2.0mg/1 NAA+0.5mg/1 BAP	+	25
MS+0.5mg/1 NAA+1mg/1 BAP	+++	85
MS+1mg/1 NAA+1mg/1 BAP	+++	83
MS+1.5mg/1 NAA+1mg/1 BAP	+++	66
MS+2.0mg/1 NAA+1mg/1 BAP	++++	100
MS+0.5mg/1 NAA+1.5mg/1 BAP	+	33
MS+1mg/1 NAA+1.5mg/1 BAP	+	33
MS+1.5mg/1 NAA+1.5mg/1 BAP	+	33
MS+2.0mg/1 NAA+1.5mg/1 BAP	+	50

Table 3. Combinations tried for shoot introduction from leaf derived calli of *C.forskohlii*

Media combinations	Observation	%Reponse
MS	shoot initiation	+
MS+0.5mg/1 BAP	shoot initiation	+
MS+1.0mg/1 BAP	shoots formed	+
MS+1.5mg/1 BAP	Greening of calli + shoots	+
MS+2.0mg/1 BAP	Greening of calli + shoots	+++
MS+0.5mg/1 KN	Callus remained green	+++
MS+1.0mg/1 KN	Callus remained green	+++
MS+1.5mg/1 KN	Callus remained green	++++
MS+2.0mg/1 KN	Callus remained green	+

- no response

+ poor

++ moderate

+++ good

The 50-60 days old greenish calli (Plate 2.B) were subcultured on either MS medium devoid of growth hormones or with KN or BA at varying concentration (Table.3). The medium augmented with KN alone (0.5 to 2mg/g) did not produce any shoot buds, though the calli remained green. But with BA (0.5 to 2 mg/l) shoot buds were initiated after 7-8 days of culture.


Out of these MS +1 mg/l BA produced maximum number of shoots (plate.1C).

Higher concentrations of BA have been found to be

inhibitory for the differentiation of shoots. More than 15 shoots were produced per culture (Fig.1)

Rooting of healthy shoots was tried on half strength as well as MS basal medium or Ms medium supplemented with growth hormones at varying concentrations (Table.4). MS medium augmented with IAA or IBA did not induce any roots or shoots, while with NAA at lower concentrations (0.1-0.4mg/l), the shoots produced roots (Plate 1.D, E). At higher concentrations however the bases of shoots produced

Fig 1.Effect of BA on shoot induction from leaf derived calli in *C.forskohlii*



calli. In half strength and full strength medium devoid of any hormones also the shoots produced roots. More number of lengthy roots were found in half strength MS basal medium.

Three weeks old rooted plants weaned in a small polyhouse for 10-12 days in high humid condition (70-80 %) were transferred to ambient conditions into cups filled with 50% soil and 50% coco peat.(Plate, 1 F) Before transfer, the *in vitro* raised plantlets were dipped in 20% glycerol and then transferred to the soil, to reduce the rate of transpiration.

Determination of maximum inhibitory concentration (MIC):

Standardization of the Maximum Inhibitory Concentration (MIC) of the Fungal culture Filtrate (FCF) to the callus tissue of *Coleus forskohlii* was carried out by incorporating various concentrations of the FCF in the MS medium. It was observed that the callus retained its morphogenetic potential at 5% and 10% concentrations. At concentrations above 15% the growth and differentiation of the callus were inhibited. At 45% and 50% concentration of FCF the callus turned

completely brown and eventually died.

DISCUSSION

Callus initiation from leaf disc explants has been reported from various Lamiaceae members, such as *Coleus blumei* (Dodds and Roberts, 1982; Hervey and Robbins, 1978), *C.purriiflorus* (Ashokan et al.1983), *Perilla frutescens* (tanimoto and Harada, 1980) *Mentha citrata* (Van Eck and Kitto, 1988;1992).

Higher concentrations of BA (1-2 mg/l) has been reported to induce callus formation from leaf explants of *Curculigo orchioides* (Augustine and D'souza. 1997) whereas in another member of Lamiaceae, *Mentha piperita*, 0.5 mg/leach BA and IBA was enough for callus induction from leaf discs (Faure et al. 1998). In the same species Jullien et al. (1998) induced calli in combination of 2, 4-D, NA, and BA.

The present observations with callus induction in *Coleus forskohlii* closely resemble the study on *Salvia officianalis*, another medicinal plant, where equimolar concentrations of auxin and cytokinin BA +NAA

Table 4. Media combinations tried for rhizogenesis of *C.forskohlii*

Media combinations	Reponse
MS half strength	+++
MS full strength	++
MS+0.1mg/l NAA	+
MS+0.2mg/l NAA	+
MS+0.4mg/l NAA	+
MS+0.5mg/l NAA	Callus induction from base of the green
MS+1.0mg/l NAA	Callus induction from base of the green
MS+0.2mg/l IAA	-
MS+0.5mg/l IAA	-
MS+1mg/l IAA	-
MS+0.2mg/l IBA	-
MS+0.5mg/l IBA	-
MS+1mg/l IBA	-

- no response + poor ++ moderate +++ good

stimulated callus formation from leaf explants. (Kintziouse et al. 1999). In Peppermint another Lamiaceae member, callus was obtained with 2, 4-D with NAA+BA at equal concentrations was found to be good for callus induction.

In *Mintostachys* (Chebel et al. 1998) MS +NAA and BA was found to induce higher number of shoots per nodal explants.

In earlier studies with *Pogostemon cablin* (Benth), another aromatic plant regenerating shoots were obtained from calli derived from leaf segments using MS +1 mg/l BA while Kinetin was found to be less effective for shooting (Misra, 1996). Similar results could be observed in the present study also. Sen and Sharma (1991) reported maximum number of shoots on MS medium augmented with 2 mg/l IBA within 20-25 days through shoot tip culture in

Coleus forskohlii while Sharma et al.(1991) reported shoot induction from nodal segment culture of *C.forskohlii* on MS+ 1 mg/l mg/l KN + 1mg/l IAA. In the present study our results are similar to those of Sen and Sharma, 1991.

Sharma et al. (1991) induced roots *in vitro* grown shoots of *Coleus forskohlii* on MS +1 mg/l NAA while Sen and Sharma (1991) obtained rooting on MS basal medium in a few plants. In the present study better and efficient rooting of shoots could be observed in half strength MS basal medium.

Acclimatization of *in vitro* plantlets is often difficult because they possess succulent stems and leaves due to high humidity with in the culture vessels and free water in the medium besides poorly developed cuticle, reduced differentiation of palisade cells large intracellular spaces and defective due to ambient conditions (Thomas, 1998)

The transition from *in vitro* to *ex vitro* is a critical phase, required to make the large heterotrophic plantlet completely autotrophic through photosynthesis and preparing it to withstand the unusual water loss by development of protective layer and regulation of stomatal functioning. The use of increased light intensity and gradual reduction in pH has been demonstrated to facilitate *ex vitro* establishment (Thomas, 1998; Mao et al, 2000).

Incorporation of an inhibitor or an anti metabolite at least two times the level of MIC, is one of the most common strategies used for the selection of plants resistant against diseases and various other environmental stresses (Suprasanna and Rao, 1998) A number of economically important plant species have been selected for disease resistance using the crude FCF of the pathogenic fungi as and inhibitor in the culture medium (Morpurgo et al, 1994; Venkatachalam et al, 1998) These include plant species such as Rice (Vidyasekaran et al, 1990), Tobacco (Thanutong et al, 1983), Potato (Behneke 1979) and Groundnut (Venkatachalam et al 1998).

Once the MIC is established the selection for disease resistance could be achieved using either a single step or a multi step selection method. In the single step selection procedure the inhibitor is added at least two to three times the level of MIC and cultures are maintained for several subculture regimes with the inhibitor (Gonzales, 1985) In the multi step selection procedure sub-lethal concentration (less than MIC) is added into the medium and subsequently a gradual increase in inhibitor level is maintained (McCoy, 1987; Miao et al, 1988).

In the present study 40% FCF in the medium showed maximum inhibition and is therefore considered as the MIC level of the FCF to *Coleus forskohlii*. This data could prove to be useful for the future for selecting a resistant *C.forskohlii* plant against the root disease caused by *L.theobromae*.

The protocol standardized for *in vitro* regeneration of *Coleus forskohlii* will be a useful tool for mass propagation of this important medicinal plant. In addition, the MIC value determined for the FCF of *L.theobromae* can be employed for the selection of disease resistant plants at cellular level.

LITERATURE CITED

- Anonymous (1950) The wealth of India Raw material Vol.2.Council of Scientific and Industrial Research (CSIR) India pp.308-309
- Ashokan MP, O`Hair SK, and Litz R.E. (1983) In vitro plant development from bulbil explants of two *Discorea* species. Hort. Science 18:702-703
- Augustine A.C and D`Souza L.(1997) Regeneration of anticarcinogenic herb, *Curculigo orchioides*. In vitro Cell.Dev.Biol-Plant. 33:111-113
- Behneke M (1979) Selection of potato callus for resistance to culture filtrates of *Phytophthora infestans* and regeneration of resistance plants. theor. Apple. Genet. 55:69-71
- Chebel A, Koroch AR, Juliani HR; and Trippi V.(1998) Micropropagation of *Mentha Stachys Mollia*(H.B.K) Grieseb and essential oil composition of clonally of clonally propagated plants. In vitro Cell. Dev. Biol. Plant 34:249-251
- Dodds JH, and Roberts LW (1982) Experiment in plant tissue culture, Cambridge University Press, Cambridge, London, New York.
- Faure O, Dimer F, Moja S and Jullien F(1998) Mannitol and Thidiazuron improve in vitro shoot regeneration from Spearmint and Peppermint leaf discs. Plant. Cell. Tiss. Org. Cult. 52:209-212
- Gozales GA and Widholm JM (1985) Selection of plant cells for desirable characteristics. Inhibitor resistance In:Plant Cell culture A practical Approach Ed.Dixon RA Irl Press Oxford UK. Pp. 67-78
- Gupta R (1998) India Journal of Plant Genetic Resources 1:98-102
- Gupta O and Nema KG(1979) Effect of different temperature and relative humidity on development of fruit rot of Papaya caused by *Botryodiplodia thebromae* and *colletotrichum papayae*. Indian Photopathology 32:106-107
- Hervey A, and Robbins WJ (1978) Development of plants from leaf Cell line selection. Edix. PJ, VCH Weinheim pp113-150

- Jullien F, Diemer F, Colson M Faure O (1998) An optimizing protocol for protoplast regeneration of three peppermint cultivators (*mentha xpiperita*) Plant Cell. Tiss. Org. Cult. 54:153-159
- Kintzious S, Nikolaou A, Shoula M (1999) Somatic embryogenesis and invitro rosmarinic acid accumulation in *Salvia officuinalis* & *S. Fruticosa* leaf callus culture. Plant Cell Reports 18:462-466
- Khamboj VP (2000) Herbal Medicine. Curr.Sci. 76(1):35-38
- Larkin PJ and Scowcraft WR (1983) Somaclonal variation and eye spot toxin tolerance in sugercane. Plant Cell Tiss. Org. Cult. 2:112-122
- Laverde S. Philip A and Frank L (1973) Mycotic Keratitis 5 cases caused by unusual fungi. Sabouraudia 11:119
- Mao AA, Wetten A, Fay and Caligari PDS(2000) Invitro prpagation of *Litsea cubeba* (lours) Pers. A multi purpose tree. Plant Cell Reports 19:263-267
- McCoy TJ (1987) Cherecterization of Alfalfa (*Medicago sativa* L.) plants regeneratnts from selected NaCl tolerant cell lines Plant Cell Reports 6:417-422
- Mc-Ginnis MR (1985) Mycotic diseases of Proposed nomenclature Int.Jour.Derm. 24:9
- Miao S, Duncan DR and Widholm JM(1988) Selection of reneerable maiza callus cultures resistant to 5-methy tryptophan 5-2-aminiethy L-cysteine and high levels of L-lysine plus L-trionine. Plant Cell, Tissue and Org. Cult 14:3-14
- Misra M. (1996) Regeneration of Patchouli(*Pogostemon cablin* Benth.) plants from leaf and node callus and evaluation after growth in the field Plant cell Reports 15:991-994
- Morpurgo R, Lopato SV, Afza RA and Novak FJ (late) (1994) Selection Parameters for resistance to oxysporom F.sp Cubense rceland race 4 on diploid banana (*Musa accuminata* Colla) Euphytica.00:1-9
- Murashige T and Skoog F(1962) A revised medium for rapid growth & bioassay with tobacco tissue culture. Physiol. Plant 15:473-497
- Sen J and Sharma AK (1991) *Invitro* propagation of *Coleus forskohlii* Briq. For forskolin synthesis. Plant Cell Reports 9:696-698
- Sharma N, Chandel K.P.S and Srivastava V.K(1991). *In vitro* propagation of *C.forskohli* Briq. A threatened medicinal plant. Plant Cell Report 10:67-70
- Sukh Dev (1997) Ethnotherapeutics and modern drug development;The potential of Ayurveda Curr.Sci.73(11)909-928
- Suprasanna P and Rao PS (1998) Selection of mutants using plant cell and tissue culture techniques. Advances in Plany Physiology Vol.1(ed) A Hemantarajah Scientific publishers India.
- Sulton BC (1980) The Coelomycetes. C.M.I., Kew, Surrey, UK
- Tanimoto S, and Harada H (1980) Hormonal of Morphogenesis in leaf explants of *Perilla frutescens*. Britton Var. Crispa, DeCaisnef. Viridi-crispa Makino. Ann. Bot. 45:321-327
- Thanutog. P Fursawa I, Yamamota M (1983) Resistance tobacco plants from protoplast derived calluses selected for their resistance to *Psuedamonas* and *Alternaria toxins*. Theor. Appl.Genet 66:209-215
- Thomas P(1998) Humid incubation period and plantlet age influence acclimatization and establishment of micropropagated grapes. *In vitro* Cell. Dev. Bio. Plant. 34:52-56
- Valdes III LJ, Mislanker SG and Paul A(1987) Economic Botany 41:474-483
- Van ECK JM and Kitto SL(1992) Regeneration of peppermintand orange mint from leaf discs. Plant. Tiss. Cell. Org. Cult, 30:41-49
- Van ECK JM and Kitto SL (1988) Regeneration of Mint. Hort. Sci. 23:803
- VanKatachalm P, Sanker rao K, Kavikshore PB and Jayabalan N (1998) Regeneration of late leaf spot resistance of ground nut plant *Cercosporidium*

personatum culture filtrates treated callus. Current Science 74(1):61-65

Verpoorte R, Heijden R, Hoopen HJG and Memclink J (1999) Metabolic engineering of plant secondary metabolism. Biotech Letter 21:467-479

Vidyasekharan P (1993) In: Genetic Eng. Mol. Bio. and tissue culture (ed. Vidyasekharan P.) pp 295-309

Vidyasekharan P, Ling DH, Borromeo ES, Zapata FJ and Men TW (1990) Annals of Applied Biology 117:515-523

Received 2000.12.1

Accepted 2001. 2.1