

Porphyrin을 포함하는 Crown Ether를 가진 사각포피린 형광물질 합성에 관한 연구

Synthesis and It's Luminescent Properties of Porphyrin Square Compound with Crown Ether Containing Pyridine

Seung-Hyun Chang*, Kwang-Bo Chung**, Kwan-young Lee***

<요약>

THF/toluene용매내에서 pyridine이 포함된 porphyrin과 $\text{Re}(\text{CO})_5\text{Cl}$ 의 반응에 의해 새로운 porphyrin square(1)을 합성하였다. 2-(Methylene 15-crown-5)-nicotinoyl chloride ester(2)는 CH_2Cl_2 용매내에서 2-Hydroxymethyl-15-crown-5와 nicotinoyl chloride로부터 합성하였다. 형광 실험에서 게스트(2)가 호스트(1)에 더해짐으로써 luminescence가 감소되는 것을 알았다. 호스트-게스트계에서 (2)의 농도가 감소함에 따라 결합상수 k 는 $1.13 \times 10^7 \text{M}^{-1}$ 이었다. 이 계에 Na^+ 농도를 증가시킴으로써 luminescence는 현저히 증가하였다.

Key Words : Porphyrin Square, Crown Ether, luminescence

Introduction

Rigid macrocycles based on cis bridging ligand of transition metals represented an unusual class of compounds having tremendous promise in host-guest, inclusion, and molecular recognition chemistry^[1,2]. The 50 or so available squares have been constructed from platinum, palladium, nickel, tungsten, and rhenium precursors and have been assembled in both homometallic and heterometallic form^[3,4]. New materials featuring exceptional nanometer scale porosity, enormous internal surface areas and enormous molecular binding capacities could be readily fabricated from an extended family of neutral, luminescent, tetrametallic cyclophanes recently developed in our lab^[5].

Depending on cavity size and overall charge, selected molecular squares were capable of functioning as solution phase hosts for either

complex anionic guests or neutral aromatic guests where binding was achieved, respectively, via coulombic or hydrophobic interactions. While host-guest binding was typically monitored via NMR spectroscopy, detection based on host luminescence had also been demonstrated^[6]. Particularly versatile in the respect were molecular squares derived from $\text{Re}(\text{CO})_5\text{Cl}$. Here, we intended to report on the synthesis of the square and on sensing and molecular recognition applications involving higher order assemblies of the porphyrin molecular squares in Figure 1.

Experimental

All reaction were conducted under a dry argon atmosphere with oven-dried glassware. All solvents were purified by distillation and dried, if necessary, prior to use. All chemicals

* 대구대학교 화학과 / 工博 / shchang@taegu.ac.kr
** 한서대학교 화학공학과 / 工博 / chemorg@hanseo.ac.kr
*** 대구대학교 건설환경공학부 / 工博 / kylee@taegu.ac.kr

* Dept. of Chemistry, Taegu Univ.
** Dept. of Chemical Engineering, Hanseo Univ.
*** Dept. of Civil Engineering, Taegu Univ.




Fig. 1 Synthetic route and structure of porphyrin square (3) and crown ether (4).

were purchased in reagent grade and used without further purification. ^1H -NMR and ^{13}C -NMR spectra were recorded on a Varian-300 spectrometer and referenced to residual protic solvent with chemical shifts being reported as δ ppm from TMS. IR-spectra were recorded on a Shimadzu FT IR-470 spectrometer. Mass spectra were recorded on a high resolution mass spectrometer.

The absorption and fluorescence spectra were recorded by using a UV/vis absorption spectrophotometer (Varian, Cary3) and a spectrometer (Spex, FluoreMAX).

3-Acetyl-2-heptanone

A mixture of 1-bromobutane(753.5g, 5.5 mole), 2,4-pentanedione(550g, 5.5 mole), potassium carbonate(912g, 6.6 mole), and dry acetone(2.5L) were stirred and heated at reflux for 48 hours with a 5 round bottom flask equipped with a paddle stirring device, condenser, and drying tube. During this time acetone was added as necessary to maintain solution volume. The solid were then filtered off and washed with acetone. The liquid residue was then filtered to remove precipitated solid and

heated at reflux for 30 minutes with 50 mL water and 5 mL concentrated sulfuric acid. The acid was then neutralized with aqueous potassium carbonate and the aqueous layer removed. The organic layer was dried over magnesium sulfate, filtered and distilled under reduced pressure to give the product as a liquid(420g,49%), bp 54-60 °C/3mm Hg. ¹H-NMR δ 0.8 (t,3H, CH₂CH₂CH₂CH₃) ; 1.3 (m, 4H, CH₂CH₂CH₂CH₃) ; 1.8 (q, 2H, CH₂CH₂CH₂CH₃); 2.15 (s, 6H, CH₃COCH-n-butyl-COCH₃) ; 3.6 (t, 1H, CH₃COCH(n-butyl)COCH₃). MS gave only fragment peaks at m/e = 113 (48%), 100 (56%), 85 (63%), and 71 (100%); no parent peak observed.

4-Butyl-3,5-dimethyl-2-ethoxycarbonylpyrrole

Oxime : A solution of sodium nitrite (80g, 1.15 mole) in water (215 mL) was added dropwise to a stirred solution of ethyl acetoacetate (137.5g, 1.05 mole) in acetic acid (350 mL), keeping the temperature between 0 °C and 10 °C with the use of an ice bath. After the addition was complete, the solution was stirred for an additional two hours at room temperature.




Fig. 2 Porphyrin square with one methylene crown ether moiety.

Pyrrole : A solution of 3-acetyl-2-heptanone (159g, 1.02 mole) in acetic acid (850 mL) was heated to 65 °C in a three necked 3L round bottom flask equipped with a paddle stirrer, a thermometer, a dropping funnel, and a powder funnel.

The above oxime was added dropwise simultaneously with portions of a intimate mixture of zinc dust(80g) and sodium acetate (80g).

The temperature was kept at 65 °C by use of an ice bath. More zinc dust (40g) was added during the latter course of the reaction to maintain a finely divided suspension in the flask.

After the oxime addition was complete, the contents of the flask were stirred for an additional 1.5 hours at 65 °C, then poured into a bucket of crushed ice. The solid product was collected by filtration and dissolved in methylene chloride. The methylene chloride was then decanted to remove zinc particles and the solvent removed on the rotary evaporator. The resulting solid residue was then recrystallized from absolute ethanol (300 mL) to give the product as soft white crystal (88g, 39%), mp 108-111 °C.

¹H-NMR δ 0.9 (t,3H, CH₂CH₂CH₂CH₃); 1.2-1.4 (m,7H, CH₂CH₂CH₂CH₃ + CO₂CH₂CH₃); 2.1-2.3 (s + s + t, 8H, CH₃ + CH₃ + CH₂CH₂CH₂CH₃); 4.3 (q, 2H, CO₂CH₂CH₃); 9.0 (br. s, 1H, NH). MS m/e = 223 (85%), 180 (100%), 134 (90%). HRMS. Calcd for C₁₃H₂₁NO₂: 223.15273. Found: 223.15767.

5-Acetoxymethyl-4-butyl-2-ethoxycarbonyl-3-methylpyrrole

Pyrrole(86.5g, 0.38 mole) was dissolved in glacial acetic acid (700 mL) with stirring in a 1L beaker. Lead tetraacetate (180g, 0.41 mole) was then added all at once. The resulting solution was stirred for two more hours and then poured into rapidly stirred water (1L), making sure that the acetic acid solution was thoroughly mixed with the water. The resulting viscous suspension was filtered through a buchner funnel. The collected solid was compressed with a rubber dam and then suspended in hot petroleum ether (1500 mL). Chloroform (50 mL) was added, and the clear solution was decanted into a crystallizing dish and placed in the refrigerator for 3-4 hours. The resulting solid was filtered and dried to yield the product as fluffy white needles (94g, 86%), mp 101-103 °C. ¹H-NMR δ 0.9 (t, 3H, CH₂CH₂CH₂CH₃); 1.25-1.45 (m, 7H, CH₂CH₃ + CH₂CH₂CH₂CH₃), 2.1 (s, 3H, CH₃); 2.3 (s, 3H, CH₂CO₂CCH₃); 2.45 (t,2H, CH₂CH₂CH₂CH₃); 4.3 (q,2H, CO₂CH₂CH₃); 5.05 (s,2H, CH₂CO₂CCH₃); 9.1 (br.s, 1H, NH). MS m/e = 281 (14%), 238 (27%), 223(26.3%), 180(88%), 134(100%), HRMS. Calcd for C₁₅H₂₃NO₄: 281.16271. Found: 281.16221.

Diethyl 3,7-Dibutyl-2,8-dimethyl-dipyrrol methane-1,9-dicarboxylate

Acetoxymethyl pyrrole(88.3g, 0.31 mole) was heated at reflux in 10% aqueous ethanol (660

mL) containing concentrated HCl(25 mL) for two hours and then in the refrigerator for four hours. The solid precipitate was filtered, washed with ethanol, and dried to give pure as a white crystalline powder(60g, 89%), mp 112–113°C. ¹H-NMR δ 0.9 (t, 6H, CH₂CH₂CH₂CH₃); 1.2–1.5 (m, 14H, CO₂CH₂CH₃) + 2x CH₂CH₂CH₂CH₃); 2.3 (s, 6H, CH₃); 2.4 (t, 4H, CH₂CH₂CH₂CH₃); 3.9 (s, 2H, CH₂); 4.3 (q, 4H, CO₂CH₂CH₃); 9.15 (br.s, 1H, NH). MS m/e = 430(26%), 221(100%), 208(14.3%). HRMS. Calcd for C₂₅H₃₈N₂O₄: 430.28316. Found: 430.28326.

3,7-Dibutyl-2,8-dimethyl-1,9-dipyrylmethane

Ester (71.6g, 0.17mole) was suspended in ethylene glycol (500mL) in a 1L roundbottom flask. Sodium hydroxide (29g, 0.72 mole) and the solution heated at reflux for 1 h under nitrogen. The solution first turned red and then to a yellow brown color. The solution was then poured into ice-water and filtered. The resulting solid was dissolved in hot ethanol (200 mL), filtered through celite, and concentrated in vacuo to give crude 1.09 (46g, 0.16mole). Recrystallization from ethanol gave colorless prisms, mp 96–97 °C. ¹H-NMR (300MHz, CDCl₃) δ 0.8 (t, 6H, CH₂CH₂CH₂CH₃), 1.4 (m, 8H, CH₂CH₂CH₂CH₃), 2.05(s, 6H, CH₃), 2.45(t, 4H, 2xCH₂CH₂CH₂CH₃), 3.7(s, 2H, CH₂), 6.25 (br. s, 2H, pyrrole CH), 7.1 (br. s, 2H, pyrrole NH): ¹³C-NMR (300 MHz, CDCl₃) δ 10.4, 14.0, 22.7, 24.0, 33.7, 113.7, 118.1, 119.7, 124.9 MS m/z = 286 (M⁺, 62%), 229 (30%), 149 (100%). HRMS: Calcd for C₁₉H₃₀N₂: 286.24090. Found: 286.24102.

2,8,12,18,-Tetrabutyl-3,7,13,17,-tetramethyl-5,15,-bis(4-pyridyl)porphyrin

The dipyrylmethane(2g, 6.98mmol) and freshly distilled 4-pyridine carboxaldehyde (0.748g, 6.98mmol) were dissolved in methanol(700mL) and the solution was deaired by bubbling with argon for 10 min while stirring. Trifluoroacetic acid (0.55 mL 6.98 mmol) was added and the solution was stirred overnight at room temperature under argon. At this time DDQ (2.37g, 10.5 mmol) dissolved in THF (100mL) was added and the resulting solution was stirred for an additional 4 h. The solvent was removed in vacuo. This residue was redissolved in dichloromethane, washed with saturated sodium bicarbonate solution, water, dried over anhydrous sodium sulfate, and concentrated in vacuo. The resulting solid was eluted through a silica column (1% MeOH/CH₂Cl₂) and the second

band was collected, concentrated to 80 mL, and crystallized by addition of methanol (200 mL). The dark red-purple crystals were filtered and dried in vacuo to yield (1.08g, 1.45 mmol) 41.5%: mp > 300°C Uv-vis (CH₂Cl₂) λ_{max} 406 (ε 181 000), 506 (15 300) 538 (5 820), 574(6 380) 624(1 820) ¹H-NMR(CDCl₃) δ -2.42 (br s, 2H, NH) 1.10 (t, 12H, CH₂CH₂CH₂CH₃), 2.50(s, 12H, CH₃), 3.97 (t, 8H, CH₂C H₂CH₂CH₃), 8.04 (d, 4H, β-pyridine H), 9.01 (d, 4H, α-pyridine H), 10.26 (s, 2H, 10, 20 meso H). MS m/z (%) 744 (M⁺, 7), 668 (100), 624 (25). HRMS: calcd. for C₅₀H₆₀N₆ 744.4879, found 744.4851. Anal calcd. for C₅₀H₆₀N₆ · H₂O 78.70, H 8.19, N 11.01; found C 78.68, H 7.99, N 10.97. On the basis of ¹H-NMR and Mass spectral data the leading band was determined to be 2,8,12,18,-Tetrabutyl-3,7,13,17-tetramethyl-5-(4'-pyridyl)porphyrin mp, 202 ¹H NMR (300 MHz, CDCl₃) δ -2.25 (br s, 2H, NH), 0.89 (t, 12H CH₂CH₂CH₂CH₃), 1.08 (m, 8H, CH₂CH₂CH₂CH₃), 2.03 (m, 4H, CH₂CH₂CH₂CH₃) 3.72 (t, 4H, CH₂CH₂CH₂CH₃), 3.81(t, 4H, CH₂CH₂CH₂C H₃), 7.87 (d, 2H, β-pyridine H), 8.92 (d, 2H, α-pyridine H), 9.56 (s, 1 H, 15 meso H), 9.59 (s, 2H, 10, 20 meso H). MS m/z (relative intensity %) 667(M⁺, 100), 624 (13), 334 (10)

2-(Methylene15-crown-5)-nicotinoyl ester

2-(Hyroxymethyl)-15-crown-5 250mg(1 mmol) was refluxed in CH₂Cl₂ with nicotinoyl chloride 178mg(1 mmol) and triethylamine for 12h. The mixture was filtered after cooling, and the filter cake was washed with ethylacetate. Following removal of the solvent in vacuo, the residue was crystallized from ethylacetate and hexane (10 : 1) to yield 159mg (45%) of the compound **2** as white solid. ¹H-NMR(CDCl₃, δ): 3.4~3.6(m, 16H, OCH₂CH₂O), 3.6~3.7(m, 2H, OCHCH₂O), 3.8~3.9(m, 1H, OCHCH₂O), 4.1~4.4(m, 2H, OCHCH₂O) 7.2~7.3(m, 1H, pyridine H), 8.1~8.2(d, 1H, pyridine H), 8.6~8.7(d, 1H, pyridine H), 9.0~9.1(s, 1H, pyridine H), MS(ED): m/e 372 .

Compound **1** was obtained as a maroon microcrystalline in high yield by combining Compound porphyrin 169mg with 82mg(0.227 mmol) of Re(CO)₅Cl dissolved in 100 mL of freshly distilled 4:1 tetrahydrofuran-toluene as a solvent and heated to reflux for 48 hours. Following cooling, 100mL of hexanes was added to get product precipitation, The product was isolated at 86% yield by filtration, washed with hexanes, and dried in vacuo. NMR indicated complete coordination of available pyridyl fragments. The analogous tetra-zincmetallated square was obtained in nearly quantitative

yield by first dissolving **1** in a minimum volume of CH₂Cl₂, adding a 5 mL volume of saturated Zn^{II} acetate in methanol, and stirring for 4h. Compound **1** was precipitated by adding a large volume of methanol-water mixture. The product was isolated by filtration, washed with water and then ether, and dried in vacuo.

Results and Discussion

In Figure 1 a new class tetraporphyrin square has been synthesized which introduces octahedral transition metal centers of square frameworks.




Fig. 3 Porphyrin square fluorescence emission intensity vs. methylene crown ether concentration.

The fac-Re(CO)₃(u-imine)₂Cl corners were capable of forming luminescent excited states (MLCT excited states) following visible ligands were employed as edges, however, the luminescence observed was that of the porphyrin. The tetraporphyrin square were highly soluble in CHCl₃, tetrahydrofuran, and CH₂Cl₂ but insoluble in water. This square also was enormously chromophoric, especially in the porphyrin Soret region (400 nm~). The square formation induced a bathochromic shift of 6 nm in the Soret region, consistent with net removal of electron density from the porphyrin π system upon rhenium-pyridine bond formation. Near-UV fluorescence excitation studies (MLCT region) established that rhenium corners served only a structural, rather than a direct photophysical, role within the square frame work. Luminescence measurements show that square was unresponsive to millimolar additions of pyridine, however, lead to significant red shift in the fluorescence the porphyrine square in Figure 2. From the known fluorescence response of monomeric Zn^{II} porphyrin species and the lack of

response by porphyrine square was clearly associated with axial ligation of porphyrine bound zinc by pyridine.




Fig. 4 Fluorescence emission from the porphyrin square/crown ether mixture as a function of [NaSCN].

As indicated in Figure 3 emission from tetra-zincmetallated square was decreased by added the crown ether binding to the square host cavity. From additional measurements at lower host concentrations, the host-guest association constant was $1.13 \times 10^7 \text{ M}^{-1}$. Further insight into the porphyrin square binding was provided by electronic absorption measurements. At micromolar concentrations, 1:1 mixtures exhibit an essentially complete loss of host Q band features at 542 and 576 nm. The absorption studies were consistent, therefore, with a binding geometry which entails pyridine ligation of all available Zn sites. In Figure 4 emission was enhanced again by adding NaSCN because the metal ion was bound by crown ether cavity in host-guest system. From addition of Na ion, binding constant was dramatically increased and the binding constant at that time was $4.35 \times 10^7 \text{ M}^{-1}$.

Acknowledgement

This research was financially supported by Taegu University Research Grant in 2001.

Reference

- 1) Fujita, M., Aoyagi, M., Ibukumo, F., Ogura, K., and Yamaguchi, K.; Made-to-Order Assembling of [2]Catenanes from Palladium(II)-Linked Rectangular Molecular Boxes. *J. Am. Chem. Soc.*, **120**, . pp. 611-612. (1998)

- 2) Lagrost, C., Lacroix, J. Aeiyaeh, S. Jouini M. and Lacaze, P.: Host-guest Complexation: A new Strategy for Electrodeposition of Processable Polythiophene Composites from Aqueous Medium. *Chem. Commun.*, pp. 489-490 (1998) .
- 3) Reek, J., Schenning A., Bosman, A., Meijer, E., and Crossly, M.: Templated assembly of a Molecular Capsule. *Chem. Commun.*, pp. 1289-1290 (1998) .
- 4) Mizutani, T., Horiguchi T., Koyama, H., Uratani, I. and Ogoshi, H., : Molecular Recognition and Atropisomerization of [5,10,15,20]-Tetrakis(1-pentyl-3-pyridino)porphyrinato]zinc(II) in Water. *The Chemical Society of Japan* , **71**, pp. 413-418 (1998)
- 5) Chang, S. H., Chung K., Slone, R. and Hupp, J. T.: Crown ether Functionalization of a Porphyrin-base "Molecular Square": Induction of Fluorescence sensitivity to Alkali Metal Cations. *Synthetic Metals.*, **117**, pp.215-217. (2001)
- 6) Inouye M., Akamatsu, K. and Nakazumi, H.: New Crown Spirobenzopyrans as Light- and Ion-Responsive Dual-Mode Signal Transducers. *J. Am. Chem. Soc.*, **119**, pp. 9160-9165 (1997)

(2002년 5월 25일 접수, 2002년 8월 20일 채택)