

# Telomerase: Key to Mortal or Immortal Road

Eun-Young Yang, Young Hoon Sung and Han-Woong Lee

---

*Samsung Biomedical Research Institute, Molecular Therapy Research Center, Samsung Medical Center, Department of Molecular Cell Biology, Sungkyunkwan University School of Medicine, Suwon, Korea*

## ABSTRACT

Gradual attrition of telomere to a critical short length elicits successive cellular response of cellular senescence and crisis. Cancer cells evade this process by maintaining functional telomeres via one of two known mechanisms of telomere maintenance. The first and most frequent mechanism involves reactivation of enzyme activity of telomerase, a ribonucleoprotein complex mainly via transcriptional up-regulation of TERT, a catalytic subunit of telomerase complex. The second mechanism utilizes telomerase-independent way termed ALT (for Alternative Lengthening of Telomere), which possibly involves recombination pathways. Thus master key for cellular immortalization is supposed to possess adequate telomere reserves. Indeed, telomerase can alone induce the immortalization under culture on feeder cell layers without generally known inactivation mechanism of tumor suppressor genes. Including this phenomena, this review will focus on telomerase and telomere-associated proteins, thereby implication of these proteins for cellular immortalization processes. (**Immune Network 2002;2(4):183-188**)

**Key Words:** Telomere, telomerase, telomere-binding protein, immortalization, tumorigenesis

## Introduction

Normal human cells possess a limited replicative lifespan when cultured in vitro (1). In these cells, telomeres, the ends of linear chromosomes, shorten with each cell division (2,3). This observation has led to the suggestion that telomeres act as a molecular counting device that checks the number of cell divisions and limits further division, upon shortening below a threshold length (4). In contrast, the majority of cells derived from spontaneously arising human tumors proliferate indefinitely and maintain stable telomere lengths (5). This observation has led to the inference that maintenance of telomere length is a prerequisite for the acquisition of immortalized replicative capacity. In unicellular eukaryotes and in the germline of multicellular organisms, replication-associated loss of telomeric DNA is counteracted in a variety of ways. The best studied of these is the ribonucleoprotein enzyme complex, telomerase, that uses an RNA template to add repeats onto the G-rich strand,

thus extending the single-stranded 3' overhang. TERT, catalytic subunit of telomerase, is important to maintain the telomere length and TERC, RNA subunit of telomerase, is essential for the enzymatic function of telomerase. Normal human somatic cells have the telomerase activity of either low or undetectable levels and progressive telomere loss occurs with each cell division. According to the telomere hypothesis of senescence, telomere erosion eventually acts as the trigger for cells to senesce, meaning that cell immortalization requires a mechanism for prevention of telomere attrition (6).

Here we summarize the regulation mechanism of telomere length, especially by telomere-binding proteins, and recent progress showing the importance of telomerase activity for telomere maintenance and their role in immortalization.

## Important roles of telomere-binding proteins on protection and maintenance of chromosomal end

Telomeres are predominately double stranded. However, these end in a 30~200 nucleotide single-stranded overhang (7-9). This 3' overhang can invade and anneal with the double-stranded region of telomeric DNA to form displacement (D)-loop in the

---

Correspondence to: Han-Woong Lee, Department of Molecular Cell Biology, Sungkyunkwan University School of Medicine, 300 Chonchon-dong, Jangan-gu, Suwon 440-746, Korea. (Tel) +82-31-299-6135, (Fax) +82-31-299-6435, (E-mail) hwl@skku.ac.kr

same telomere (10). This overall feature is called as a telomere (T)-loop. Although T-loops can be formed on relatively short telomeres, presumably there is a minimum size below, which they cannot form.

Because only the overexpression of the catalytic subunit of telomerase (TERT) reconstitutes the telomerase activity and increases telomere length in telomerase negative cells, it seems that activation of TERT is the limiting step for the induction of telomerase activity in most cells. Therefore, much effort has been focused on the understanding of the transcriptional regulation of TERT. Besides the transcriptional regulation of TERT, there may be complex and dynamic processes regulating the actual activity of telomerase (11).

However, sufficient length of telomeric DNA does not prevent abnormalities in the integrity of 3' overhang or mutation of telomere-binding proteins from inducing telomere dysfunctions and thereby several deleterious cellular effects. Functional telomeres require a minimum length of telomere repeats, the integrity of 3' overhang and functional telomere-binding proteins (Table I). To achieve and maintain the correct telomeric structure and the integrity of 3' overhang, several telomere-binding proteins are needed. Most of them participate in some steps involved in the regulation of telomere length and/or structure. For example, the formation of T-loops is critically dependent on TRF2. It directly binds double stranded telomeric DNA and can remodel the telomeric DNA into T-loop (10). Elimination of TRF2 from telomeric DNA induces immediate deprotection of chromosomal ends and the cells show some features resembling the cellular responses to a double strand break (DSB) in the genome. In fact,

dependent protein kinase complex are involved in signaling pathway induced by abnormal telomeres and regulate telomere integrity. For example, in cells from Ku86-deficient mice, telomeric fusions occur with sufficient length of telomeres at the fusion point (12). However, this Ku86 deficiency rescues the end-to-end chromosomal fusion induced by critically short telomeres in telomerase-deficient mice (13). Also, DNA-PKcs deficiency induces effects similar to Ku86 deficiency (14). These lines of evidence suggest that some components of DNA-PK complex have important roles in maintenance of telomere integrity and protection of telomeres from DNA repair activity.

Similar to TRF2, TRF1 binds double stranded telomeric DNA. TRF1 and its binding partner, TIN2 may facilitate or stabilize the formation and/or maintenance of T-loops (10,15). Another TRF1 binding protein, Tankyrase 1 shows poly (adenosine diphosphate-ribose) polymerase (PARP) activity and, indeed, exerts its activity on TRF1 (16). Tankyrase 1-mediated ADP-rybosylation of TRF1 decreases the telomeric DNA binding activity of TRF1 and promotes telomere elongation in a telomerase-dependent manner (17). Tankyrase 2, the homologue of Tankyrase 1, can bind to TRF1 and seems to have similar roles on telomere (18).

Whereas TRF1 and TRF2 bind double stranded telomeric DNA, Pot1 binds 3' overhang of telomeric DNA and seems to have important roles in maintaining chromosomal stability by capping the single-stranded telomeric DNA (19).

As a result, both telomere length and telomere structure are important in protection and maintenance of chromosomal end. Also, besides telomerase itself, telomere-binding proteins, as well as their

**Table I.** Telomere binding proteins

Genes	Mediator for telomere association	Roles in telomeric DNA
POT1	Single stranded telomeric DNA	Telomere-length maintenance and telomere protection
TRF1	Double stranded telomeric DNA	T-loop formation
TRF2	Double stranded telomeric DNA	T-loop formation
Tankyrase 1	Via TRF1	Ribosylate TRF1
Tankyrase 2	Via TRF1	Ribosylate TRF1
TIN2	Via TRF1	Negatively regulate telomere length
RAP1	Via TRF2	Regulate telomere length
RAD50/NBS1/MRE11	Via TRF2	Maybe participate in T-loop formation
PINX1	Via TRF1	Bind TERT and inhibit its activity
Ku86	Telomere repeat?	Maybe restrict telomerase and DNA repair activity
DNA-PKcs	Ku proteins	Protect telomeres

some evidence indicates that components of DNA-

associated proteins, have pivotal roles on modulation

and/or maintenance of telomere length or structure.

### Telomerase and immortalization

The primary hallmark of tumor cells is loss of growth control caused by gain-of-function mutations in proto-oncogenes, and loss-of-function mutations in tumor suppressor genes. A second hallmark of tumor cells, common to almost all malignant cancers, is cellular immortality. In human tumors, cell immortalization almost always involves derepression of the gene for the catalytic subunit of telomerase, hTERT (20,21). However, it remains controversial whether the cell culture immortality of many cancer cells is a fundamental property, required for tumorigenesis, or a by-product of other genetic changes.

After finite *in vitro* cell division, most human somatic cells face the loss of their replicative proliferating capacity (1), and are termed cellular senescence. Among a variety of triggers for cellular senescence, the telomere shortening that occurs in each cell division as a result of the end-replication problem of DNA polymerase is recognized as "mitotic clock" to control the cellular life span, at least in human somatic cells. Given that human cells can divide only a limited number of times in culture, many researchers have wondered whether there is a "clock" that measures cell divisions. Human somatic cells were found to have shorter telomeres than sperm from the same individuals, which suggests that human somatic telomeres shorten during development (3,22). Furthermore, telomere length in somatic cells, both fibroblasts and leukocytes, from older individuals was decreased on average compared with telomere length in somatic cells from younger people (2,3). Also, telomeres were found to shorten during the passage of fibroblasts in culture (3) and in cancers compared with adjacent normal tissues (2). These results have supported the hypothesis which telomerase is shut off in human somatic cells, limiting their lifespan, whereas it is activated in cells that become immortalized in culture (6,23).


Although oncogenically transformed cells can extend the lifespan of human cells beyond the first growth arrest point, known as senescence, these cells eventually enter a phase known as crisis and then suffer chromosome aberration and massive cell death. Rare immortal cells are survived by telomerase activation (24-26) and this was established by studies of telomere length and telomerase activity in mortal and immortal human cultured cells. Analysis of telomere lengths in human embryonic kidney cells transformed with simian virus 40 T antigen shows a steady decline until crisis, and then telomere maintenance in the survivors. Concordantly, the telomerase enzyme is inactive in the precrisis mortal cells and is activated

in the immortalized cell clones (26,27). Whereas the human telomerase RNA subunit is expressed in both telomerase-positive and telomerase-negative tissues (28,29), the hTERT mRNA is expressed in immortal, telomerase-positive cell lines but not in mortal, telomerase-negative cells (7,30,31). Furthermore, over-expression of hTERT in previously telomerase-negative cells generates an active telomerase enzyme in these cells, confirming that hTERT expression is a key step in regulating telomerase activity (32-36). Even more dramatically, ectopic expression of the telomerase catalytic subunit gene in certain telomerase-negative cells is able to extend the lifespan of these cells, confirming the model that telomerase can overcome the limits of human cell mortality, at least in some cases.

In addition, hTERT expression leads to significant lifespan extension in strains of mortal fibroblasts and retinal epithelial cells, allowing them to bypass the senescence limitation to cell growth (34,36). For example, hTERT-immortalized BJ foreskin fibroblasts have been found to proliferate for an additional 200 population doublings compared with control cell clones that can double for approximately 80 generations (37). Telomerase expression is sufficient to immortalize some cell types, such as fibroblasts, directly, but several reports have claimed that keratinocytes and other types of epithelial cells exhibit an additional mechanism of replicative aging involving the p16/pRB pathway that is independent of telomere length, and that inactivation of both mechanisms is also required for cellular immortalization (38-42).

However, in most of these reports, cells were cultured in chemically defined media and proliferated for only 15~20 doublings. It was dramatically less than the ~50 doublings described previously for the growth of keratinocytes on feeder layers (43). Furthermore, 15~20 doublings appear grossly inadequate to explain the estimated number of keratinocyte doublings occurring *in vivo* (44). For these reasons, several groups have developed the different model systems to examine whether the proposed p16/pRB pathway actually represented a telomere-independent second mechanism of cellular senescence or was a secondary consequence of particular culture conditions. They found that telomerase activity alone could immortalize without the inactivation of p16/pRB pathway. In other words, human keratinocytes grown on appropriate feeder layers could directly immortalized by hTERT maintenance of telomeres without any intervening inactivation of a p16/pRB pathway (45,46).

Human mammary epithelial (HME) cells have been described to exhibit the mechanism of senescence



**Figure 1.** Schematic of telomerase hypothesis in immortalization and tumorigenesis. This diagram is based on Neumann AA et al. and modified according to recent studies.

(38,47). Generally, it has been known that HME cells appear to senesce in two stages. One is an initial growth arrest, designated mortality stage 0 (M0), occurring after about 20 doublings. This stage is apparently controlled by the Rb/p16 pathway, because M0 can either be prevented by E7 (48) or a few cells escape spontaneously with reduced p16 expression due to promoter methylation (47). The other is subsequent M1 growth arrest bypassing by E6 (49,50). In some cases, only a subpopulation of cells can emerge from mortality stage 2, which is called, as E6/E7-expressing cells will enter a period of crisis in culture. However, recently, the replicative capacity of HME cells can also be extended without inactivating p16 if these cells are cultured on feeder layers (45,47). Finally, culture of human fibroblasts under inadequate culture conditions (chemically defined medium in the presence of 0.25% serum) recapitulates the keratinocyte and HME scenarios. Such fibroblasts arrest after ~25 doublings regardless of telomere length/telomerase, but can be rescued and immortalized following transfer to adequate culture media (46). Although one cannot theoretically exclude the existence of non-telomere based mechanisms for replicative aging, these results establish that at present there is no reliable evidence for the presence of a second mechanism of replicative aging specifically in these cells.

Analysis of telomerase activity in immortal cells has led to the identification of two classes of cells. In one class, telomerase is activated. In the other class of cells, telomerase activity remains undetectable, and telomeres seem to be maintained by a telomerase-independent mechanism known as alternative lengthening of telomeres (ALT). Some of primary human tumors as well as immortalized cell lines have likewise been observed to include a population of telomerase-negative tumors, presumably surviving by ALT (51,52). Recently, it was reported that ALT could not substitute for telomerase expression and an additional function of hTERT that goes beyond its ability to elongate telomeres was required for cell transformation (53). These results clearly indicate that the ability of hTERT to elongate telomeres is not essential for facilitating function in tumorigenesis although the telomere lengthening is important to acquire the immortality.

## Conclusions

Telomerase is the critical enzyme in overcoming growth limitations due to telomere dysfunction. Many scientists directly have addressed the role of telomere maintenance in control of proliferative potential and the mechanisms leading to tumorigenesis. Also, it has found that acquirement of immortality requires both telomerase activity and inactivation of oncogenes or tumor suppressors (Fig. 1a). However, from recent

results, under adequate culture conditions, telomerase is sufficient to immortalize normal cells leading to tumorigenesis. Besides, another functions of telomerase are essential for facilitating the tumorigenesis, as well as its ability to elongate telomere (Fig. 1b). Therefore, telomerase may alone extend the lifespan of mortal cells in adequate culture and contribute to experimental tumorigenesis. Among the many cellular regulatory mechanisms for immortality, the use of telomerase-based cell immortalization may offer a variety of practical applications in medical research and potentially even in improving therapies. To do this, continuing challenges for telomerase research will be required, including the elucidating the composition of telomerase complexes, the role of TERT variants, the interactions between telomerase and other components of the telomere, control of TERT transcription. To understand telomerase biology, we are currently studying the different regulation mechanisms of telomerase through interacting proteins in normal cells and immortalized cells.

## References

- Hayflick L, Moorhead PS: The serial cultivation of human diploid cell strains. *Exp Cell Res* 25; 585-621, 1961
- Harley CB, Futcher AB, Greider CW: Telomeres shorten during ageing of human fibroblasts. *Nature* 345;458-460, 1990
- Hastie ND, Dempster M, Dunlop MG, Thompson AM, Green DK, Allshire RC: Telomere reduction in human colorectal carcinoma and with ageing. *Nature* 346;866-868, 1990
- Harley CB, Vaziri H, Counter CM, Allsopp RC: The telomere hypothesis of cellular aging. *Exp Gerontol* 27;375-382, 1992
- Harley CB, Kim NW, Prowse KR, Weinrich SL, Hirsch KS, West MD, Bacchetti S, Hirte HW, Counter CM, Greider CW: Cold Spring Harbor Symp. *Quant Biol* 59;307-315, 1994
- Harley CB: Telomere loss: mitotic clock or genetic time bomb? *Mutat Res* 256;271-282, 1991
- Wellinger RJ, Ethier K, Labrecque P, Zakian VA: Evidence for a new step in telomere maintenance. *Cell* 85;423-433, 1996
- Makarov VL, Hirose Y, Langmore JP: Long G tails at both ends of human chromosomes suggest a C strand degradation mechanism for telomere shortening. *Cell* 88;657-666, 1997
- Wright WE, Tesmer VM, Huffman KE, Levene SD, Shay JW: Normal human chromosomes have long G-rich telomeric overhangs at one end. *Genes Dev* 11;2801-2809, 1997
- Griffith JD, Comeau L, Rosenfield S, Stansel RM, Bianchi A, Moss H, de Lange T: Mammalian telomeres end in a large duplex loop. *Cell* 97;503-514, 1999
- Aisner DL, Wright WE, Shay JW: Telomerase regulation: not just flipping the switch. *Curr Opin Genet Dev* 12;80-85, 2002
- Samper E, Goytisolo FA, Slijepcevic P, van Buul PP, Blasco MA: Mammalian Ku86 protein prevents telomeric fusions independently of the length of TTAGGG repeats and the G-strand overhang. *EMBO Rep* 1;244-252, 2000
- Espejel S, Franco S, Rodriguez-Perales S, Bouffler SD, Cigudosa JC, Blasco MA: Mammalian Ku86 mediates chromosomal fusions and apoptosis caused by critically short telomeres. *EMBO J* 21;2207-2219, 2002
- Goytisolo FA, Samper E, Edmonson S, Taccioli GE, Blasco MA: The absence of the DNA-dependent protein kinase catalytic subunit in mice results in anaphase bridges and in increased telomeric fusions with normal telomere length and G-strand overhang. *Mol Cell Biol* 21;3642, 2001
- Kim SH, Kaminker P, Campisi J: TIN2, a new regulator of telomere length in human cells. *Nat Genet* 23;405-412, 1999
- Smith S, Giriati I, Schmitt A, de Lange T: Tankyrase, a poly (ADP-ribose) polymerase at human telomeres. *Science* 282; 1484-1487, 1998
- Cook BD, Dynek JN, Chang W, Shostak G, Smith S: Role for the related poly (ADP-Ribose) polymerases tankyrase 1 and 2 at human telomeres. *Mol Cell Biol* 22;332-342, 2002
- Kaminker PG, Kim SH, Taylor RD, Zebajadian Y, Funk WD, Morin GB, Yaswen P, Campisi J: TANK2, a new TRF1-associated poly (ADP-ribose) polymerase, causes rapid induction of cell death upon overexpression. *J Biol Chem* 276;35891-35899, 2001
- Baumann P, Cech TR: Pot1, the Putative Telomere End-Binding Protein in Fission Yeast and Humans. *Science* 292; 1171-1175, 2001
- Meyerson M, Counter CM, Eaton EN, Ellisen LW, Steiner P, Caddle SD, Ziaugra L, Beijersbergen RL, Davidoff MJ, Liu Q, Bacchetti S, Haber DA, Weinberg RA: hEST2, the putative human telomerase catalytic subunit gene, is up-regulated in tumor cells and during immortalization. *Cell* 90;785-795, 1997
- Nakamura TM, Morin GB, Chapman KB, Winrich SL, Andrews WH, Lingner J, Harley CB, Cech TR: Telomerase catalytic subunit homologs from fission yeast and human. *Science* 277;955-959, 1997
- de Lange T, Shinue L, Myers RM: Structure and variability of human chromosome ends. *Mol Cell Biol* 10;518-527, 1990
- Greider CW: Telomeres, telomerase and senescence. *Bioessays* 12;363-369, 1990
- Huschtscha LI, Holliday R: Limited and unlimited growth of SV40-transformed cells from human diploid MRC-5 fibroblasts. *J Cell Sci* 63;77-99, 1983
- Wright WE, Pereira-Smith OM, Shay JW: Reversible cellular senescence: Implications for immortalization of normal human diploid fibroblasts. *Mol Cell Biol* 9;3088-3092, 1989
- Counter CM, Avilion AA, Le Feuvre CE, Stewart NG, Greider CW, Harley CB, Bacchetti S: Telomere shortening associated with chromosome instability is arrested in immortal cells which express telomerase activity. *EMBO J* 11; 1921-1929, 1992
- Counter CM, Botelho FM, Wang P, Harley CB, Bacchetti S: Stabilization of short telomeres and telomerase activity accompany immortalization of Epstein-Barr virus-transformed human B lymphocytes. *J Virol* 68;3410-3414, 1994
- Avilion AA, Piatyszek MA, Gupta J, Shay JW, Bacchetti S, Greider CW: Human telomerase RNA and telomerase activity in immortal cell lines and tumor tissues. *Cancer Res* 56;645-650, 1996
- Bestilny LJ, Brown CB, Miura Y, Robertson LD, Riabowol KT: Selective inhibition of telomerase activity during terminal differentiation of immortal cell lines. *Cancer Res* 56;3796-3802, 1996
- Nakamura TM, Morin GB, Chapman KB, Weinrich SL, Andrews WH, Lingner J, Harley CB, Cech TR: Telomerase catalytic subunit homologs from fission yeast and human. *Science* 277;955-959, 1997
- Kilian A, Bowtell DD, Abud HE, Hime GR, Venter DJ, Keese PK, Duncan EL, Reddel RR, Jefferson RA: Isolation of a candidate human telomerase catalytic subunit gene,

- which reveals complex splicing patterns in different cell types. *Hum Mol Genet* 6;2011-2019, 1997
32. Nakayama J, Tahara H, Tahara E, Saito M, Ito K, Nakamura H, Nakanishi T, Tahara E, Ide T, Ishikawa F: Telomerase activation by hTERT in human normal fibroblasts and hepatocellular carcinomas. *Nat Genet* 18;65-68, 1998
  33. Weinrich SL, Pruzan R, Ma L, Ouellette M, Tesmer VM, Holt SE, Bodnar AG, Lichtsteiner S, Kim NW, Trager JB, Taylor RD, Carlos R, Andrews WH, Wright WE, Shay JW, Harley CB, Morin GB: Reconstitution of human telomerase with the template RNA component hTR and the catalytic protein subunit hTERT. *Nat Genet* 17;498-502, 1997
  34. Bodnar AG, Ouellette M, Frolkis M, Holt SE, Chiu CP, Morin GB, Harley CB, Shay JW, Lichtsteiner S, Wright WE: Extension of lifespan by introduction of telomerase into normal human cells. *Science* 279;349-352, 1998
  35. Counter CM, Meyerson M, Eaton EN, Ellisen LW, Caddle SD, Haber DA, Weinberg RA: Telomerase activity is restored in human cells by ectopic expression of hTERT (hEST2), the catalytic subunit of telomerase. *Oncogene* 16;1217-1222, 1998
  36. Vaziri H, Benchimol S: Reconstitution of telomerase activity in normal human cells leads to elongation of telomeres and extended replicative life span. *Curr Biol* 8;279-282, 1998
  37. Morales CP, Holt SE, Ouellette M, Kaur KJ, Yan Y, Wilson KS, White MA, Wright WE, Shay JW: Absence of cancer-associated changes in human fibroblasts immortalized with telomerase. *Nat Genet* 21;115-118, 1999
  38. Kiyono T, Foster SA, Koop JI, McDougall JK, Galloway DA, Klingelutz AJ: Both Rb/p16INK4a inactivation and telomerase activity are required to immortalize human epithelial cells. *Nature* 396;84-88, 1998
  39. Jarrard DF, Sarkar S, Shi Y, Yeager TR, Magrane G, Kinoshita H, Nassif N, Meisner L, Newton MA, Waldman FM, Reznikoff CA: p16/pRb pathway alterations are required for bypassing senescence in human prostate epithelial cells. *Cancer Res* 59;2957-2964, 1999
  40. Dickson MA, Hahn WC, Ino Y, Ronfard V, Wu JY, Weinberg RA, Louis DN, Li FP, Rheinwald JG: Human keratinocytes that express hTERT and also bypass a p16 (INK4a)-enforced mechanism that limits life span become immortal yet retain normal growth and differentiation characteristics. *Mol Cell Biol* 20;1436-1447, 2000
  41. Farwell DG, Shera KA, Koop JI, Bonnet GA, Matthews CP, Reuther GW, Coltrera, MD, McDougall JK, Klingelutz AJ: Genetic and epigenetic changes in human epithelial cells immortalized by telomerase. *Am J Pathol* 156;1537-1547, 2000
  42. Jones CJ, Kipling D, Morris M, Hepburn P, Skinner J, Bounacer A, Wyllie FS, Ivan M, Bartek J, Wynford-Thomas D, Bond JA: Evidence for a telomere-independent "clock" limiting RAS oncogene-driven proliferation of human thyroid epithelial cells. *Mol Cell Biol* 20;5690-5699, 2000
  43. Green H, Rheinwald JG, Sun TT: Properties of an epithelial cell type in culture: The epidermal keratinocyte and its dependence on products of the fibroblast. *Prog Clin Biol Res* 17;493-500, 1977
  44. Rubin H: Cell aging in vivo and in vitro. *Mech Aging Dev* 98;1-35, 1997
  45. Herbert BS, Wright WE, Shay JW: p16 (INK4a) inactivation is not required to immortalize human mammary epithelial cells. *Oncogene* 21;7897-7900, 2002
  46. Ramirez RD, Morales CP, Herbert BS, Rohde JM, Passons J, Shay JW, Wright WE: Putative telomere-independent mechanisms of replicative aging reflect inadequate growth conditions. *Genes Dev* 15;398-403, 2001
  47. Foster SA, Wong DJ, Barrett MT, Galloway DA: Inactivation of p16 in human mammary epithelial cells by CpG island methylation. *Mol Cell Biol* 18;1793-1801, 1998
  48. Foster SA, Galloway DA: Human papillomavirus type 16 E7 alleviates a proliferation block in early passage human mammary epithelial cells. *Oncogene* 12;1773-1779, 1996
  49. Shay JW, Wright WE, Brasiskyte D, van der Haegen BA: E6 of human papillomavirus type 16 can overcome the M1 stage of immortalization in human mammary epithelial cells but not in human fibroblasts. *Oncogene* 8;1407-1413, 1993
  50. Band V, Zajchowski D, Kulesa V, Sager R: Human papilloma virus DNAs immortalize normal human mammary epithelial cells and reduce their growth factor requirements. *PNAS* 87;463-467, 1990
  51. Bryan TM, Englezou A, Dalla-Pozza L, Dunham MA, Reddel RR: Evidence for an alternative mechanism for maintaining telomere length in human tumors and tumor-derived cell lines. *Nat Med* 3;1271-1274, 1997
  52. Murnane JP, Sabatier L, Marder BA, Morgan WF: Telomere dynamics in an immortal human cell line. *EMBO J* 13; 4953-4962, 1994
  53. Stewart SA, Hahn WC, O'Connor BF, Banner BN, Lundberg AS, Hana PM, M Brooks MW, Fleming M, Zimonjic DB, Popescu NC, Weinberg RA: Telomerase contributes to tumorigenesis by a telomere length-independent mechanism. *PNAS* 99;12606-12611, 2002