

Synthesis of 1,2,3,4-Tetrahydroisoquinoline-2-sulfonic Acids

Ji Sun Lee, Sun Hee Kim, Han Sik Yoon,[†] and Chai-Ho Lee*


Department of Chemistry, Wonkwang University, Jeonbuk 570-749, Korea

**School of Medical Radiation, Wonkwang Health Science College, Jeonbuk 570-750, Korea*


Received May 7, 2003

Key Words : Tetrahydroisoquinoline. Sulfamic acid. Iminium ion

The 1,2,3,4-tetrahydroisoquinoline (THIQ) alkaloids **1**, which are widely distributed in plant and animal kingdoms, have received much attention because of their important biological activities.¹ For example, 1-aryl- or 1-alkyl-1,2,3,4-tetrahydroisoquinolines present in mammalian brain play a major role in therapy of variety of neurological disorders.²



Two general, acid mediated procedures have been reported for the preparation of THIQ ring system. The first entails the Pictet-Spengler reaction of a 2-arylethylamine **2** with aldehyde.³ The second method requires the treatment of a *N*-substituted-2-arylethylamine **3** (*i.e.*, *N*-acyl⁴ and *N*-sulfonyl group⁵ on the nitrogen) with carbonyl compound (*i.e.*, aldehyde, and the corresponding acetal). In these processes, introduction of an electron withdrawing substituents on nitrogen is to increase the electrophilicity of iminium intermediate. Previously, we demonstrated the intramolecular- and intermolecular α -sulfamidoalkylation transformation proceeding through the intermediacy of an iminium ion provide an expeditious route for the preparation of cyclic sulfamides containing THIQ ring.⁶ The sulfamic acid also have a sulfonic acid group as an electron withdrawing group on the nitrogen atom and is readily hydrolyzed to the



Scheme 1. Synthesis of **8**. Reagents and conditions: (i) 1) $\text{ClSO}_3\text{H}\cdot\text{Et}_3\text{N}$, $-5\text{--}0^\circ\text{C}$, 2) $c\text{-HCl}$; (ii) HCO_2H .

corresponding amines.⁷

In this present study, we report on the reaction of *N*-(2-arylethyl)sulfonic acids **6** with aldehydes **7** (or the corresponding acetals **8**) in formic acid for generation of 1,2,3,4-tetrahydroisoquinoline-2-sulfonic acids **9**.

The starting sulfamic acid **6** were prepared according to established synthetic protocols.⁸ When amine **5** was reacted with $\text{ClSO}_3\text{H}\cdot\text{Et}_3\text{N}$ in chloroform, followed by treatment with hydrochloric acid, sulfamic acid **6** was formed in 60-70 % yield. Intramolecular cyclization of **6** with **7** or **8** in formic acid (96% in H_2O) gave the desired product **9** in good yield. Identification of all the isolated products **9** was accomplished with the aid of infrared, ^1H and ^{13}C NMR, and mass

Table 1. Reaction condition, mp, and yield of product **9**

Entry	No.	R ¹	R	Reaction conditions		mp (°C) ^a	Yield (%) ^b
				Temp. (°C)	Time (h)		
1	9a	H	H	rt	1	200-213	81
2	9b	H	OMe	rt	1	172-200	65
3	9c	CH ₂ Cl	H	rt	12	75-80	74
4	9d	CH ₂ Cl	OMe	rt	0.5	174-176	92
5	9e	CH ₂ CN	H	rt	24	205-215	65
6	9f	CH ₂ CN	OMe	rt	1	194-195	68
7	9g	COOEt	H	rt	48	104-108	58
8	9h	COOEt	OMe	rt	48	170-171	67
9	9i	Benzyl	H	rt	24	142-145	84
10	9j	Benzyl	OMe	rt	48	150-152	73
11	9k	Phenyl	H	50	24	169-171	65
12	9l	Phenyl	OMe	50	24	172-176	78
13	9m	3-methoxy-4-hydroxyphenyl	H	50	24	117-119	62
14	9n	3-methoxy-4-hydroxyphenyl	OMe	50	24	76-99	71
15	9o	3,4,5-trimethoxyphenyl	H	rt	48	159-160	66
16	9p	3,4,5-trimethoxyphenyl	OMe	50	24	169-176	68
17	9q	2-furyl	H	rt	48	180-188	70
18	9r	2-furyl	OMe	rt	48	134-140	74
19	9s	2-thiophenyl	H	rt	24	186-190	70
20	9t	2-thiophenyl	OMe	rt	24	230-238	76

^aIsolated yields. ^bMelting points are uncorrected.

spectroscopy. In the infrared spectrum, the compound **9** exhibited characteristic absorption bands at 1246-1277 and 1016-1119 cm^{-1} for the sulfonyl group.⁹ Diagnostic signals of compounds **9** were observed at 4.38-6.13 ppm in the ^1H NMR spectra and at 48.5-60.8 ppm in the ^{13}C NMR spectra for the methine (C-1) unit furnished by aldehydes **7** (or acetals **8**).

In conclusion, we have developed a general and versatile method for the synthesis of 1, 2, 3, 4-tetrahydroisoquinoline-2-sulfonic acids, by the reaction of *N*-(2-arylethyl)sulfamic acids with aldehydes (or acetals).

Experimental Section

Typical experimental procedure for synthesis of *N*-arylethylsulfamic acid **6.** A solution of arylethylamine **5** (10 mmol) and triethylamine (12 mmol) in 30 mL of CHCl_3 was stirred at $-5-0^\circ\text{C}$ and chlorosulfonic acid (10 mmol) was added dropwise so as to maintain the temperature below 0°C . The solution was acidified with 1 *N* HCl solution to pH 2. The solid that precipitated was filtered to give the desired products **6**.

2-(3-Methoxyphenyl)ethylsulfamic acid (6a): Beginning with 3-methoxyphenethylamine (1.51 g), compound **6a** was obtained in 55% yield (1.27 g); mp $144-146^\circ\text{C}$; IR (KBr) 3180 (NH), 1288, 1063 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 2.87 (t, $J = 8.3$ Hz, 2H), 3.21 (t, $J = 8.3$ Hz, 2H), 3.75 (s, 3H), 6.79-6.82 (m, 3H), 7.22-7.25 (m, 1H), 10.18 (br s, 1H) ppm; ^{13}C NMR (DMSO- d_6) δ 32.5, 45.6, 55.5, 112.9, 114.7, 121.3, 130.2, 139.4, 160.0 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 230.1, found 230.2.

2-(3,4-Dimethoxyphenyl)ethylsulfamic acid (6b): Beginning with 3,4-dimethoxyphenethylamine (1.81 g), compound **6b** was obtained in 67% yield (1.75 g); mp $164-170^\circ\text{C}$; IR (KBr) 3146 (NH), 1259, 1072 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 2.82 (t, $J = 8.3$ Hz, 2H), 3.19 (t, $J = 8.3$ Hz, 2H), 3.71 (s, 3H), 3.75 (s, 3H), 6.73 (dd, $J = 1.8$ Hz and $J = 7.8$ Hz, 1H), 6.82 (d, $J = 1.8$ Hz, 1H), 6.87 (d, $J = 7.8$ Hz, 1H), 10.18 (br s, 1H) ppm; ^{13}C NMR (DMSO- d_6) δ 32.1, 45.9, 56.0, 56.1, 112.6, 113.0, 121.0, 130.2, 184.2, 149.4 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 260.1, found 260.2.

Typical experimental procedure for synthesis of 1,2,3,4-tetrahydroisoquinoline-2-sulfonic acid **9.** A formic acid (96% in H_2O , 20 mL) solution of arylethylsulfamic acids **6** (2.0 mmol) and aldehydes **7** (or acetals **8**) (2.0 mmol) was stirred, and then the solution was quenched with 1 *N* HCl (50 mL). The solid that precipitated was filtered and then recrystallized from methanol-chloroform to give the desired products **9** (Table 1).

9a: IR (KBr) 1269, 1063 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.13 (t, $J = 6.4$ Hz, 2H), 3.59 (t, $J = 6.4$ Hz, 2H), 3.73 (s, 3H), 4.39 (s, 2H), 6.75 (d, $J = 2.3$ Hz, 1H), 6.78 (dd, $J = 2.3$ Hz and $J = 8.4$ Hz, 1H), 7.12 (d, $J = 8.4$ Hz, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 26.3, 45.9, 48.5, 54.9, 113.1, 113.4, 120.8, 128.1, 133.1, 159.3 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 242.1, found 242.7.

9b: IR (KBr) 1258, 1057 cm^{-1} ; ^1H NMR (acetone- d_6) δ

3.07 (t, $J = 6.4$ Hz, 2H), 3.59 (t, $J = 6.4$ Hz, 2H), 3.74 (s, 6H), 4.38 (s, 2H), 6.77 (s, 1H), 6.79 (s, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 25.6, 46.1, 48.5, 55.4, 55.5, 109.8, 111.5, 120.5, 123.8, 148.2, 148.7 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 272.1, found 272.7.

9c: IR (KBr) 1265, 1047 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.05 (ddd, $J = 5.5$ Hz, $J = 5.5$ Hz, and $J = 17.6$ Hz, 1H), 3.18 (ddd, $J = 5.5$ Hz, $J = 8.3$ Hz, and $J = 17.6$ Hz, 1H), 3.45 (ddd, $J = 5.5$ Hz, $J = 8.3$ Hz, and $J = 13.2$ Hz, 1H), 3.69 (ddd, $J = 5.5$ Hz, $J = 5.5$ Hz, and $J = 13.2$ Hz, 1H), 3.74 (s, 3H), 4.18 (dd, $J = 7.1$ Hz and $J = 12.8$ Hz, 1H), 4.29 (dd, $J = 3.4$ Hz and $J = 12.8$ Hz, 1H), 4.94 (dd, $J = 3.4$ Hz and $J = 7.1$ Hz, 1H), 6.79 (d, $J = 2.7$ Hz, 1H), 6.83 (dd, $J = 2.7$ Hz and $J = 8.7$ Hz, 1H), 7.30 (d, $J = 8.7$ Hz, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 25.3, 40.0, 44.6, 55.1, 55.7, 113.7, 113.8, 120.7, 127.9, 134.5, 159.5 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 290.0, found 290.2.

9d: IR (KBr) 1273, 1071 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.03 (ddd, $J = 5.5$ Hz, $J = 5.8$ Hz, and $J = 17.1$ Hz, 1H), 3.14 (ddd, $J = 5.5$ Hz, $J = 7.6$ Hz, and $J = 17.1$ Hz, 1H), 3.50 (ddd, $J = 5.8$ Hz, $J = 7.6$ Hz, and $J = 12.5$ Hz, 1H), 3.74 (ddd, $J = 5.5$ Hz, $J = 5.5$ Hz, and $J = 12.5$ Hz, 1H), 3.78 (s, 6H), 4.25 (dd, $J = 7.3$ Hz and $J = 12.9$ Hz, 1H), 4.38 (dd, $J = 3.6$ Hz and $J = 12.9$ Hz, 1H), 4.97 (dd, $J = 3.6$ Hz and $J = 7.3$ Hz, 1H), 6.83 (s, 1H), 7.01 (s, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 24.6, 40.1, 44.7, 55.3, 55.6, 55.8, 109.7, 112.0, 120.4, 125.3, 148.6, 149.5 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 320.0, found 320.2.

9e: IR (KBr) 1251, 1049 cm^{-1} ; ^1H NMR (methanol- d_3) δ 3.10 (ddd, $J = 6.4$ Hz, $J = 6.4$ Hz, and $J = 17.5$ Hz, 1H), 3.18 (ddd, $J = 6.4$ Hz, $J = 7.1$ Hz, and $J = 17.5$ Hz, 1H), 3.36 (dd, $J = 5.0$ Hz and $J = 17.6$ Hz, 1H), 3.40 (dd, $J = 6.1$ Hz and $J = 17.6$ Hz, 1H), 3.48 (ddd, $J = 6.4$ Hz, $J = 6.4$ Hz, and $J = 13.0$ Hz, 1H), 3.64 (ddd, $J = 5.5$ Hz, $J = 7.1$ Hz, and $J = 13.0$ Hz, 1H), 3.80 (s, 3H), 4.89 (dd, $J = 5.0$ Hz and $J = 6.1$ Hz, 1H), 6.84 (d, $J = 2.3$ Hz, 1H), 6.91 (dd, $J = 2.3$ Hz and $J = 8.7$ Hz, 1H), 7.31 (d, $J = 8.7$ Hz, 1H) ppm; ^{13}C NMR (methanol- d_3) δ 22.0, 24.9, 39.6, 51.1, 54.6, 113.5, 113.8, 115.8, 121.1, 127.6, 133.3, 160.2 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 281.1, found 281.7.

9f: IR (KBr) 1271, 1071 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.05-3.09 (m, 2H), 3.38 (dd, $J = 5.0$ Hz and $J = 17.9$ Hz, 1H), 3.54 (dd, $J = 5.5$ Hz and $J = 17.9$ Hz, 1H), 3.54 (ddd, $J = 5.0$ Hz, $J = 5.7$ Hz, and $J = 12.7$ Hz, 1H), 3.69 (ddd, $J = 5.5$ Hz, $J = 7.3$ Hz, and $J = 12.7$ Hz, 1H), 3.76 (s, 3H), 3.77 (s, 3H), 4.97 (dd, $J = 5.0$ Hz and $J = 5.5$ Hz, 1H), 6.82 (s, 1H), 6.96 (s, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 22.7, 24.3, 39.5, 50.6, 55.4, 55.6, 109.5, 111.9, 116.8, 120.9, 124.7, 148.4, 149.5 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 311.1, found 311.7.

9g: IR (KBr) 1248, 1016 cm^{-1} ; ^1H NMR (acetone- d_6) δ 1.29 (t, $J = 7.1$ Hz, 3H), 3.14 (ddd, $J = 5.5$ Hz, $J = 5.7$ Hz, and $J = 17.2$ Hz, 1H), 3.24 (ddd, $J = 5.5$ Hz, $J = 8.7$ Hz, and $J = 17.2$ Hz, 1H), 3.67 (ddd, $J = 5.5$ Hz, $J = 5.7$ Hz, and $J = 12.3$ Hz, 1H), 3.77-3.81 (m, 1H), 3.80 (s, 3H), 4.31 (q, $J = 7.1$ Hz, 2H), 5.45 (s, 1H), 6.85 (d, $J = 2.7$ Hz, 1H), 6.89 (dd, $J = 2.7$ Hz and $J = 8.7$ Hz, 1H), 7.46 (d, $J = 8.7$ Hz, 1H)

ppm: ^{13}C NMR (acetone- d_6) δ 13.4, 25.0, 39.9, 54.9, 55.7, 62.8, 113.4, 113.7, 117.9, 129.1, 134.1, 160.0, 167.6 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 315.1, found 315.7.

9h: IR (KBr) 1273, 1061 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.33 (t, $J = 7.3$ Hz, 3H) 2.96 (ddd, $J = 3.2$ Hz, $J = 3.2$ Hz, and $J = 16.5$ Hz, 1H), 3.14 (ddd, $J = 4.1$ Hz, $J = 12.4$ Hz, and $J = 16.5$ Hz, 1H), 3.45 (ddd, $J = 3.2$ Hz, $J = 12.4$ Hz, and $J = 12.4$ Hz, 1H), 4.21 (ddd, $J = 3.2$ Hz, $J = 3.2$ Hz, and $J = 12.4$ Hz, 1H), 3.86 (s, 3H), 3.87 (s, 3H), 4.35 (q, $J = 7.3$ Hz, 2H), 5.43 (s, 1H), 6.65 (s, 1H), 6.93 (s, 1H) ppm; ^{13}C NMR (CDCl_3) δ 14.1, 26.2, 45.9, 56.1, 56.2, 60.8, 64.3, 109.7, 111.2, 124.8, 148.8, 149.8, 149.7, 169.0 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 344.1, found 344.7.

9i: IR (KBr) 1246, 1059 cm^{-1} ; ^1H NMR (methanol- d_3) δ 3.07 (ddd, $J = 6.0$ Hz, $J = 6.9$ Hz, and $J = 16.3$ Hz, 1H), 3.12 (dd, $J = 8.2$ Hz and $J = 13.9$ Hz, 1H), 3.16 (ddd, $J = 6.0$ Hz, $J = 7.1$ Hz, and $J = 16.3$ Hz, 1H), 3.33 (ddd, $J = 6.0$ Hz, $J = 7.1$ Hz, and $J = 12.6$ Hz, 1H), 3.49 (dd, $J = 6.4$ Hz and $J = 13.8$ Hz, 1H), 3.53 (ddd, $J = 6.0$ Hz, $J = 6.9$ Hz, and $J = 12.6$ Hz, 1H), 3.78 (s, 3H), 4.77 (dd, $J = 6.4$ Hz and $J = 8.2$ Hz, 1H), 6.79 (d, $J = 2.3$ Hz, 1H), 6.81 (dd, $J = 2.3$ Hz and $J = 8.2$ Hz, 1H), 7.05 (d, $J = 8.2$ Hz, 1H), 7.31-7.35 (m, 3H), 7.37-7.41 (m, 2H) ppm; ^{13}C NMR (methanol- d_3) δ 25.2, 39.2, 39.8, 54.5, 56.4, 113.1, 113.2, 123.4, 127.5, 127.9, 128.9, 129.4, 132.8, 135.3, 129.7 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 332.1, found 332.7.

9j: IR (KBr) 1262, 1061 cm^{-1} ; ^1H NMR (methanol- d_3) δ 3.04 (ddd, $J = 3.9$ Hz, $J = 6.2$ Hz, and $J = 17.4$ Hz, 1H), 3.09 (ddd, $J = 6.0$ Hz, $J = 7.6$ Hz, and $J = 17.4$ Hz, 1H), 3.18 (ddd, $J = 3.9$ Hz, $J = 7.6$ Hz, and $J = 13.3$ Hz, 1H), 3.36 (ddd, $J = 6.0$ Hz, $J = 6.2$ Hz, and $J = 13.3$ Hz, 1H), 3.42 (dd, $J = 7.3$ Hz and $J = 13.7$ Hz, 1H), 3.55 (dd, $J = 7.4$ Hz and $J = 13.7$ Hz, 1H), 3.67 (s, 3H), 3.81 (s, 3H), 4.74 (dd, $J = 7.3$ Hz and $J = 7.3$ Hz, 1H), 6.42 (s, 1H), 6.81 (s, 1H), 7.31-7.35 (m, 3H), 7.38-7.41 (m, 2H) ppm; ^{13}C NMR (methanol- d_3) δ 24.5, 39.1, 40.0, 53.8, 55.1, 55.3, 110.3, 111.8, 123.2, 124.0, 127.4, 129.0, 130.0, 135.9, 147.6, 149.0 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 362.1, found 362.7.

9k: IR (KBr) 1269, 1067 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.15 (ddd, $J = 5.5$ Hz, $J = 5.5$ Hz, and $J = 17.4$ Hz, 1H), 3.30 (ddd, $J = 6.5$ Hz, $J = 8.7$ Hz, and $J = 17.4$ Hz, 1H), 3.47 (ddd, $J = 5.5$ Hz, $J = 8.7$ Hz, and $J = 12.9$ Hz, 1H), 3.53 (ddd, $J = 5.5$ Hz, $J = 6.5$ Hz, and $J = 12.9$ Hz, 1H), 3.80 (s, 3H), 5.66 (s, 1H), 6.72 (d, $J = 8.7$ Hz, 1H), 6.78 (dd, $J = 2.8$ Hz and $J = 8.7$ Hz, 1H), 6.88 (d, $J = 2.8$ Hz, 1H), 7.35-7.37 (m, 2H), 7.46-7.48 (m, 3H) ppm; ^{13}C NMR (acetone- d_6) δ 25.3, 40.0, 55.0, 59.2, 113.1, 113.6, 123.7, 129.1, 129.4, 129.6, 130.1, 134.2, 137.0, 159.3 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 318.1, found 318.7.

9l: IR (KBr) 1262, 1063 cm^{-1} ; ^1H NMR (methanol- d_3) δ 3.11 (ddd, $J = 5.5$ Hz, $J = 6.0$ Hz, and $J = 17.4$ Hz, 1H), 3.23 (ddd, $J = 6.0$ Hz, $J = 6.4$ Hz, and $J = 17.4$ Hz, 1H), 3.43 (ddd, $J = 5.5$ Hz, $J = 7.3$ Hz, and $J = 13.0$ Hz, 1H), 3.49 (ddd, $J = 6.0$ Hz, $J = 6.4$ Hz, and $J = 13.0$ Hz, 1H), 3.59 (s, 3H), 3.86 (s, 3H), 5.69 (s, 1H), 6.34 (s, 1H), 6.90 (s, 1H), 7.35-7.37 (m, 2H), 7.47-7.49 (m, 3H) ppm; ^{13}C NMR (methanol- d_3) δ 24.5, 39.5, 55.1, 55.3, 59.3, 110.7, 111.4,

122.9, 124.8, 129.1, 129.7, 129.8, 136.4, 148.3, 149.5 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 348.1, found 348.7.

9m: IR (KBr) 1263, 1034 cm^{-1} ; ^1H NMR (methanol- d_3) δ 3.12 (ddd, $J = 5.5$ Hz, $J = 5.5$ Hz, and $J = 17.6$ Hz, 1H), 3.30 (ddd, $J = 6.2$ Hz, $J = 8.2$ Hz, and $J = 17.6$ Hz, 1H), 3.46 (ddd, $J = 5.5$ Hz, $J = 8.2$ Hz, and $J = 12.7$ Hz, 1H), 3.53 (ddd, $J = 5.5$ Hz, $J = 6.2$ Hz, and $J = 12.7$ Hz, 1H), 3.83 (s, 3H), 3.79 (s, 3H), 5.60 (s, 1H), 6.75 (d, $J = 1.9$ Hz and $J = 8.3$ Hz, 1H), 6.79 (s, 2H), 6.84 (d, $J = 8.3$ Hz, 2H), 6.95 (d, $J = 1.9$ Hz, 1H) ppm; ^{13}C NMR (methanol- d_3) δ 25.2, 39.8, 54.5, 55.2, 59.7, 112.7, 112.8, 113.5, 115.3, 122.7, 123.6, 127.5, 129.3, 133.5, 147.9, 148.2, 159.7 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 364.1, found 364.7.

9n: IR (KBr) 1260, 1059 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.08 (ddd, $J = 6.0$ Hz, $J = 6.0$ Hz, and $J = 17.2$ Hz, 1H), 3.20 (ddd, $J = 6.4$ Hz, $J = 7.6$ Hz, and $J = 17.2$ Hz, 1H), 3.42 (ddd, $J = 6.0$ Hz, $J = 7.6$ Hz, and $J = 12.8$ Hz, 1H), 3.49 (ddd, $J = 6.0$ Hz, $J = 6.4$ Hz, and $J = 12.8$ Hz, 1H), 3.62 (s, 3H), 3.84 (s, 3H), 3.85 (s, 3H), 5.61 (s, 1H), 6.39 (s, 1H), 6.76 (dd, $J = 1.9$ Hz and $J = 8.0$ Hz, 1H), 6.85 (d, $J = 8.0$ Hz, 1H), 6.87 (s, 1H), 6.94 (d, $J = 1.9$ Hz, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 24.8, 40.4, 55.2, 55.4, 55.9, 59.5, 111.3, 111.6, 113.5, 114.9, 123.2, 123.6, 125.2, 128.0, 147.1, 147.9, 148.4, 149.6 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 394.1, found 394.7.

9o: IR (KBr) 1277, 1057 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.13 (ddd, $J = 5.0$ Hz, $J = 5.5$ Hz, and $J = 17.5$ Hz, 1H), 3.41 (ddd, $J = 6.0$ Hz, $J = 9.9$ Hz, and $J = 17.5$ Hz, 1H), 3.55 (ddd, $J = 5.5$ Hz, $J = 9.9$ Hz, and $J = 12.7$ Hz, 1H), 3.68 (ddd, $J = 5.0$ Hz, $J = 5.5$ Hz, and $J = 12.7$ Hz, 1H), 3.71 (s, 3H), 3.73 (s, 6H), 3.80 (s, 3H), 6.03 (s, 1H), 6.62 (s, 2H), 6.80 (d, $J = 2.8$ Hz and $J = 8.7$ Hz, 1H), 6.87 (d, $J = 2.8$ Hz, 2H), 6.96 (d, $J = 8.7$ Hz, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 25.2, 40.6, 54.9, 55.9, 59.8, 59.9, 107.4, 113.1, 113.5, 123.6, 129.4, 132.3, 134.1, 138.8, 153.7, 159.4 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 408.1, found 404.7.

9p: IR (KBr) 1261, 1051 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.10 (ddd, $J = 5.5$ Hz, $J = 5.7$ Hz, and $J = 17.4$ Hz, 1H), 3.30 (ddd, $J = 6.4$ Hz, $J = 8.0$ Hz, and $J = 17.4$ Hz, 1H), 3.52 (ddd, $J = 5.5$ Hz, $J = 8.0$ Hz, and $J = 13.8$ Hz, 1H), 3.62 (ddd, $J = 5.7$ Hz, $J = 6.4$ Hz, and $J = 13.8$ Hz, 1H), 3.06 (s, 3H), 3.72 (s, 3H), 3.75 (s, 6H), 3.81 (s, 3H), 5.75 (s, 1H), 6.43 (s, 1H), 6.74 (s, 2H), 6.87 (s, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 25.0, 41.0, 54.8, 56.0, 56.2, 56.4, 60.4, 60.9, 107.2, 110.6, 111.2, 122.5, 124.7, 131.5, 138.8, 148.2, 149.3, 153.6 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 438.1, found 438.7.

9q: IR (KBr) 1246, 1115 cm^{-1} ; ^1H NMR (methanol- d_3) δ 3.17 (t, $J = 6.4$ Hz, 2H), 3.42-3.52 (m, 2H), 5.80 (s, 1H), 6.42 (d, $J = 3.2$ Hz, 1H), 6.49 (dd, $J = 1.8$ Hz and $J = 3.2$ Hz, 1H), 6.83 (dd, $J = 2.3$ Hz and $J = 8.7$ Hz, 1H), 6.86 (d, $J = 2.3$ Hz, 1H), 6.95 (d, $J = 8.7$ Hz, 1H), 7.63 (d, $J = 1.8$ Hz, 1H) ppm; ^{13}C NMR (methanol- d_3) δ 24.9, 39.0, 52.2, 54.8, 110.8, 112.9, 113.2, 113.6, 120.7, 128.9, 133.3, 144.7, 148.9, 159.8 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 308.1, found 308.7.

9r: IR (KBr) 1265, 1119 cm^{-1} ; ^1H NMR (methanol- d_3) δ

3.12 (t, $J = 6.4$ Hz, 2H), 3.41-3.50 (m, 2H), 3.70 (s, 3H), 3.85 (s, 3H), 5.81 (s, 1H), 6.44 (d, $J = 3.2$ Hz, 1H), 6.50 (dd, $J = 1.8$ Hz and $J = 3.2$ Hz, 1H), 6.57 (s, 1H), 6.87 (s, 1H), 7.65 (d, $J = 1.8$ Hz, 1H) ppm; ^{13}C NMR (methanol- d_3) δ 24.3, 38.3, 52.1, 55.3, 110.4, 110.8, 111.6, 113.0, 120.4, 124.5, 144.7, 148.4, 148.9, 149.7 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 338.1, found 338.6.

9s: IR (KBr) 1269, 1055 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.17 (ddd, $J = 6.4$ Hz, $J = 7.1$ Hz, and $J = 17.5$ Hz, 1H), 3.23 (ddd, $J = 5.9$ Hz, $J = 6.4$ Hz, and $J = 17.5$ Hz, 1H), 3.48 (ddd, $J = 6.4$ Hz, $J = 6.4$ Hz, and $J = 12.7$ Hz, 1H), 3.54 (ddd, $J = 5.9$ Hz, $J = 7.1$ Hz, and $J = 12.7$ Hz, 1H), 3.80 (s, 3H), 6.06 (s, 1H), 6.82 (dd, $J = 2.3$ Hz and $J = 8.7$ Hz, 1H), 6.97 (d, $J = 8.7$ Hz, 1H), 7.12 (dd, $J = 3.2$ Hz and $J = 5.4$ Hz, 1H), 7.25 (d, $J = 3.2$ Hz, 1H), 7.58 (dd, $J = 0.9$ Hz and $J = 5.4$ Hz, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 25.1, 28.9, 53.5, 55.0, 113.1, 113.6, 123.5, 127.7, 128.5, 129.6, 130.6, 133.6, 139.6, 159.5 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 324.0, found 324.2.

9t: IR (KBr) 1263, 1117 cm^{-1} ; ^1H NMR (acetone- d_6) δ 3.16 (t, $J = 6.4$ Hz, 2H), 3.47-3.56 (m, 2H), 3.79 (s, 3H), 3.80 (s, 3H), 6.13 (s, 1H), 6.60 (s, 1H), 6.85 (s, 1H), 7.07 (dd, $J = 3.2$ Hz and $J = 5.0$ Hz, 1H), 7.35 (d, $J = 3.2$ Hz, 1H), 7.53 (dd, $J = 0.9$ Hz and $J = 5.0$ Hz, 1H) ppm; ^{13}C NMR (acetone- d_6) δ 24.3, 38.5, 53.2, 55.3, 55.4, 111.2, 111.5, 123.1, 124.7, 127.7, 128.5, 130.7, 139.7, 148.2, 149.7 ppm; LR FBA MS: calcd for $[\text{M}-1]^-$ 354.1, found 354.1.

Acknowledgements. This work was supported by grant No. R01-2001-000-00133-0 from the Korea Science & Engineering Foundation.

References

- (a) Padwa, A.; Beall, L. S.; Heidelbaugh, T. M.; Liu, B.; Sheehan, S. M. *J. Org. Chem.* **2000**, *65*, 2684. (b) Glunewald, G. L.; Dahanukar, V. H.; Teoh, B.; Criscione, K. R. *J. Med. Chem.* **1999**, *42*, 1982. (c) Minor, D. L.; Wyrick, S. D.; Charifson, P. S.; Watts, V. J.; Nichols, D. E.; Mailman, R. B. *J. Med. Chem.* **1994**, *37*, 4328.
- (a) Shinohara, T.; Toda, J.; Sano, T. *Chem. Pharm. Bull.* **1997**, *45*(5), 813. (b) Ornstein, P. L.; Arnold, M. B.; Allen, N. K.; Bleisch, T.; Borromeo, P. S.; Lugar, C. W.; Leander, J. D.; Lodge, D.; Schoepp, D. D. *J. Med. Chem.* **1996**, *39*, 2219. (c) Munchhof, M. J.; Meyers, A. I. *J. Org. Chem.* **1996**, *61*, 4607. (d) Carifson, P. S.; Bowen, J. P.; Wyrick, S. D.; Hoffman, A. J.; Cory, M.; McPhail, A. T.; Mailman, R. B. *J. Med. Chem.* **1989**, *32*, 2050. (f) Clark, M. T.; Adejare, A. G.; Feller, D. R.; Miller, D. D. *J. Med. Chem.* **1987**, *30*, 86. (g) Battersby, A. R.; Southgate, R.; Staunton, J.; Hurst, M. *J. Chem. Soc. (C)* **1966**, 1052.
- (a) Eric, D.; Cook, J. M. *Chem. Rev.* **1995**, *95*(6), 1797. (b) Whaley, W. M.; Govindachari, T. R. *Org. React.* **1951**, *6*, 151.
- (a) Speekamp, W. N.; Moolenaar, M. J. *Tetrahedron* **2000**, *56*, 3817. (b) Venkov, A. P.; Lukanov, L. K. *Synthesis* **1989**, 59. (c) Molloy, N. M.; Venkov, A. P. *Synthesis* **1978**, 62.
- (a) Barn, D. R.; Caulfield, W. L.; Cottney, J.; MaGurk, K.; Morphy, J. R.; Rankovic, Z.; Roberts, B. *Bioorg. & Med.* **2001**, *9*, 2609. (b) Kohno, H.; Sekine, Y. *Heterocycles* **1996**, *42*(1), 141. (c) Lukanov, L. K.; Venkov, A. P.; Molloy, M. *Synthesis* **1987**, 204.
- (a) Lee, J. S.; Yang, I. D.; Kim, S. H.; An, S. I.; Lee, C.-H. *Bull. Korean Chem. Soc.* **2003**, *24*(1), 129. (b) Lee, J. D.; Lee, C.-H.; Nakamura, H.; Ko, J.; Kang, S. O. *Tetrahedron Lett.* **2002**, *43*, 5483. (c) Lee, J. S.; Lee, C.-H. *Bull. Korean Chem. Soc.* **2002**, *23*(1), 167. (d) Lee, J. S.; Lee, C.-H. *J. Korean Chem. Soc.* **2001**, *45*, 92. (e) Kong, Y. J.; Kim, S. H.; Lee, C.-H. *J. Korean Chem. Soc.* **1999**, *43*, 131. (f) Lee, C.-H.; Kohn, H. *J. Heterocyclic Chem.* **1990**, *27*, 2107. (g) Lee, C.-H.; Kohn, H. *J. Org. Chem.* **1990**, *55*, 6098. (h) Lee, C.-H.; Kohn, H. *Heterocycle* **1988**, *27*, 2581.
- (a) Kim, B. M.; So, S. M. *Tetrahedron Lett.* **1998**, *39*, 5381. (b) Benson, G. A.; Spillane, W. J. *Chem. Rev.* **1980**, *80*, 151.
- (a) Shi, D.-F.; Bradshaw, T. D.; Chua, M.-S.; Westwell, A. D.; Stevens, M. F. *Bioorg. & Med. Chem. Lett.* **2001**, *11*, 1093. (b) Curran, W. V.; Ross, A. A.; Lee, V. J. *J. Antibiot.* **1988**, *ALI*(10), 1418.
- Nakanish, K.; Solomon, P. H. *Infrared Absorption Spectroscopy*, Holden-Day: San Francisco, 1977.