

X-Ray Structure of Keggin-type Peroxo Polyoxometalate $A-\alpha-[Si(NbO_2)_3W_9O_{37}]^{7-}$

Gyu-Shik Kim,^{*} Huadong Zeng,[†] and Craig L. Hill[†]

Department of Science Education, Kangwon National University, Chunchon 200-701, Korea

[†]Department of Chemistry, 1515 Pierce Drive, Emory University, Atlanta, GA 30322, USA

Received November 29, 2002

Key Words : Polyoxometalates, Keggin, Tungsten, Niobium, Peroxo

The title polyoxometalate (POM), $A-\alpha-[Si(NbO_2)_3W_9O_{37}]^{7-}$ ($A-\alpha-1$)¹ was shown to have potent *anti*-HIV-1 activity (median effective antiviral concentration against HIV-1 in human peripheral blood mononuclear cells (PBMC), EC₅₀, ~1 μ M) while being minimally cytotoxic (median inhibitory concentration, IC₅₀, >100 μ M also in PBMC) in 1994.^{2,3} Unfortunately this POM isn't stable in aqueous solution and decomposes into other POMs with the corresponding Keggin complex, $A-\alpha-[SiNb_3W_9O_{40}]^{7-}$, depending on conditions, generally being the dominant product. Very recently this POM, often generated *in situ* given its instability, has proven to be an effective starting material for the preparation of a range of new or larger supramolecular POMs containing the $SiNb_3W_9$ unit.^{4,6} The instability of peroxoniobium POMs, in which the $(Nb^V O_2)^{3+}$ unit has been substituted for a $(W^{VI} O)^{4+}$ type I "octahedral" unit has rendered difficult the purification and structural characterization of such POMs in the past.⁷ Given the growing importance of $A-\alpha-1$, this study sought the specific goal associated with this unstable POM, *i.e.*, to obtain a good quality single crystal X-ray diffraction study to determine the nature, and in particular, the orientation of the peroxo groups. This goal has been met as described in this note. Furthermore, the possibility of using this peroxoniobium POM anion as a synthon in both aqueous and organic solvents is of interest and could lead to other supramolecular POM species based on the $SiNb_3W_9$ unit.

Experimental Section

Materials and Methods. $Na_2WO_4 \cdot 2H_2O$ (AESAR) was obtained from a commercial source and used as received. All solvents were at least reagent grade and were used without further purification. Both $\alpha-Na_{10}SiW_9O_{34} \cdot 23H_2O$ ⁸ and $K_7HNb_6O_{19} \cdot 13H_2O$ ⁹ were prepared using literature methods. Infrared spectra were recorded on a Nicolet 510 M FTIR spectrometer using 3-5 wt.% of sample in KBr pellets. Tungsten-183 NMR (ref = 2.0 M Na_2WO_4 in D_2O) spectra were acquired at ambient temperature using an INOVA 400 FT NMR spectrometer. Elemental analyses were performed by Galbraith Laboratories Inc., Knoxville, TN, U. S. A.

Synthesis of $A-\alpha-Cs_6H[Si(NbO_2)_3W_9O_{37}] \cdot 9H_2O$ ($A-\alpha-Cs_6HI$). $K_7HNb_6O_{19} \cdot 13H_2O$ (1.91 g, 1.39 mmol) was dissolved in 250 mL of 0.5 M aqueous H_2O_2 . To this stirred

solution, 20 mL of 1.0 M aqueous HCl was added dropwise, to give a bright yellow solution, followed by addition of 7.82 g (2.72 mmol) of solid $\alpha-Na_{10}SiW_9O_{34} \cdot 23H_2O$. After the solid $Na_{10}SiW_9O_{34}$ had fully dissolved, 25.0 g (46.5 mmol) of solid CsCl was added to precipitate the Cs salt of $A-\alpha-[Si(NbO_2)_3W_9O_{37}]^{7-}$ and the resulting orange yellow mixture stirred for 15 min. The precipitate was filtered on a medium glass frit and washed with two 20-mL portions of diethyl ether. Drying in air gave 9.11 g of yellow powder (86.9% yield based on the hexaniobate precursor). Anal. Calcd. (found) for $Cs_6H_{19}SiNb_3W_9O_{52}$: H, 0.48 (0.53); Si, 0.78 (0.78); Cs, 22.2 (22.1); Nb, 7.76 (7.72); W, 46.1 (45.8). FTIR (KBr, 1100-400 cm^{-1}): 994 (w), 957 (m), 903 (vs), 868 (sh), 789 (vs), 673 (vw), 592 (w), 534 (w), 482 (vw) cm^{-1} . The TGA data suggest the presence of ~9 H_2O molecules of hydration per POM, so this many were used in the elemental analysis above. This is within experimental error of the number of H_2O molecules per POM in the X-ray structure.

X-ray Structure of $A-\alpha-[Si(NbO_2)_3W_9O_{37}]^{7-}$ ($A-\alpha-1$). X-ray quality crystals of the cesium salt were grown from aqueous solution by slow evaporation at ambient temperature. A colorless block crystal with dimensions of 0.20 \times 0.20 \times 0.17 mm was mounted on a glass fiber and X-ray diffraction data were collected at room temperature using a Siemens SMART[®] platform diffractometer equipped with a CCD area detector and a normal focus molybdenum-target X-ray tube ($\lambda = 0.71073 \text{ \AA}$).¹⁰ Data frames were collected using a narrow frame method with scan width 0.30 $^\circ$ in ω and an exposure time of 10-30 s/frame ($2\theta_{max} = 56.78^\circ$). The data were integrated using the Siemens SAINT program.¹¹ Crystal was face-indexed for absorption corrections. An empirical absorption correction based on the entire data set was applied using SADABS.¹² Redundant reflections were averaged. The corrected data were used in subsequent structure solutions and refinements. The structure was solved by direct methods using SHELXTL-PC v. 5.03 and refined by full-matrix-least-squares-on- F^2 techniques (with $F^2 > 2\sigma(F_o^2)$) using SHELXL-97, incorporated in SHELXTL-PC v. 5.03.¹³ Direct methods located the heavy atoms (W, Cs and Nb); other non-hydrogen atoms were located by successive Fourier difference synthesis. Anisotropic temperature factors were applied to all atoms, except O8 and Ow5, Ow6, Ow8 and Ow9. Disorder of Cs8 and some hydrated water (Ow5, Ow6, Ow8 and Ow9) were observed as indicated by high temperature factors, which is not uncommon in the crystal structures of POM compounds. The occupancy of

^{*}Deceased, May 2003.

Table 1. Summary of X-ray crystallographic data for 1

	$\text{Cs}_{8.66}\text{H}_{13.34}[\text{Si}(\text{NbO}_2)_3\text{W}_9\text{O}_{37}] \cdot 7\text{H}_2\text{O}$ (A- α -1)
empirical formula	$\text{Cs}_{8.66}\text{H}_{13.34}\text{O}_{50}\text{SiNb}_3\text{W}_9$
f.w.	3660.9
crystal color/shape	colorless/block
crystal size, mm	$0.20 \times 0.20 \times 0.17$
crystal system	monoclinic
space group	$C2/c$ (No. 15)
T, K	298
a , Å	18.4852 (8)
b , Å	21.1362 (9)
c , Å	26.1679 (11)
α , deg	90
β , deg	92.98 (0)
γ , deg	90
V , Å ³	10210.2 (8)
Z	8
ρ_{calc} , g cm ⁻³	4.76
F(000)	12452
μ , mm ⁻¹	25.53
θ range, deg	1.46-28.39
no. measd reflns	25781
no. indep reflns	11938
no. parameters	634
Goodness of fit	0.978
final R_1 [$I > 2\sigma(I)$]	0.0586
final wR_2 [$I > 2\sigma(I)$]	0.1319
Largest diff. peak and hole (e.Å ⁻³)	3.542 and -4.193

$$^a R_1 = \sum |F_o| - F_c| / \sum F_o, \quad ^b wR_2 = [\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)]^{1/2}$$

Cs8 was refined using a program-supplied free variable to be 32%, therefore the total amount of Cs⁺ was determined to be 6.66 in the asymmetric unit. To balance the total negative charge of seven in the POM anion 0.34 equivalent of proton is assumed in the empirical formula. Occupancy of 50% was given to the O atoms in the above-mentioned disordered water molecules without further refining their occupancies. The X-ray data collection, structure solution and refinement parameters are summarized in Table 1. Crystallographic data for the structure reported here have been deposited with FIZ (Deposition # 413046). These data can be obtained free of charge via FIZ, D-76344, Eggenstein-Leopoldshafen, Germany, e-mail: crysdata@fiz-karlsruhe.de.

Results and Discussion

X-ray crystallography and ¹⁸³W NMR (of the lithiated cesium salt, A- α -Cs₈H₁₃) provided definitive evidence that the same isomer exists in both the solid state and in aqueous solution, and that A- α isomerism is maintained in both circumstances. ORTEP and polyhedral illustrations of the polyanion portion, A- α -[Si(NbO₂)₃W₉O₃₇]⁷⁻ (A- α -1) are provided in Figure 1. Selected bond distances and angles are given in Table 2.

The X-ray structure analysis reveals that the title compound




Figure 1. Structure of the polyanion unit A- α -[Si(NbO₂)₃W₉O₃₇]⁷⁻ in A- α -1. A: ORTEP drawing with 30%-probability thermal ellipsoids; B: polyhedral drawing. The Nb^V(O₂)O₅ heptahedra, the internal SiO₄ tetrahedron, and the WO₆ pseudo-octahedra are depicted in dark, intermediate and light shading, respectively.

contains Cs⁺/H⁺ cations, A- α -[Si(NbO₂)₃W₉O₃₇]⁷⁻ (A- α -1) anions, and water molecules. Each of the three Nb atoms in the anion is coordinated by four doubly-bridging O atoms, one quadruply-bridging O atom, and one terminal η^2 -coordinated peroxo group to form a distorted pentagonal bipyramidal polyhedron. Such coordination polyhedra containing η^2 -coordinated peroxo units are seen in three other structurally determined high nuclearity¹⁴ polyoxoanions, β_3 -[(Co^{II}O₄)W₁₁O₃₁(O₂)₄]^{10-15,17} A- β -[P(NbO₂)₃W₉O₃₇]⁶⁻¹⁸ [P₂(NbO₂)₅W₁₂O₅₆]¹²⁻⁷. The average peroxide (Op-Op) bond length in the polyanion (1.46(1) Å) is between those reported for α -[P₂(NbO₂)₅W₁₂O₅₆]¹²⁻ (1.43 Å),⁷ non-coordinated O₂²⁻ (1.49 Å)¹⁹ and A- β -[P(NbO₂)₃W₉O₃₇]⁶⁻ (1.51 Å).¹⁸ The average Op-Nb-Op (44.1°) is similar to those in α -[P₂(NbO₂)₅W₁₂O₅₆]¹²⁻ (43.7°)⁷ and [Nb(O₂)₄]³⁻ (43.3°),^{20,21} but is smaller than those of A- β -[P(NbO₂)₃W₉O₃₇]⁶⁻ (45.2°).¹⁸ The peroxo groups largely eclipse the trans O-Nb-O units in the semi-equatorial plane (actually a C_{4v} pyramidal cone resulting from the overall structural distortion of the NbO₇ units). The nine W atoms in the POM anion exhibit the

Table 2. Selected bond distances [Å] and angles [°] for **1**

Distances						
Nb(1)-O(32)	1.839(12)	Nb(2)-O(31)	1.859(12)	Nb(3)-O(34)	1.852(13)	W-O _t 1.712(13)
Nb(1)-O(33)	1.945(15)	Nb(2)-O(35A)	1.930(16)	Nb(3)-O(36A)	1.957(14)	W-O _b 1.910(13)
Nb(1)-O(33A)	1.959(14)	Nb(2)-O(35)	1.948(13)	Nb(3)-O(36)	1.958(14)	W-O _c 2.381(11)
Nb(1)-O(19)	2.035(11)	Nb(2)-O(34)	2.034(12)	Nb(3)-O(30)	2.039(12)	O(33)-O(33A) 1.44(2)
Nb(1)-O(31)	2.042(13)	Nb(2)-O(24)	2.057(13)	Nb(3)-O(32)	2.045(12)	O(35)-O(35A) 1.43(2)
Nb(1)-O(14)	2.080(13)	Nb(2)-O(21)	2.062(13)	Nb(3)-O(28)	2.059(12)	O(36)-O(36A) 1.51(2)
Nb(1)-O(38)	2.298(12)	Nb(2)-O(39)	2.341(11)	Nb(3)-O(40)	2.314(12)	
Angles						
Nb(2)-O(31)-Nb(1)	148.4(6)	O(39)-Si(1)-O(38)	108.0(6)	O(39)-Si(1)-O(40)	110.1(6)	
Nb(1)-O(32)-Nb(3)	149.2(7)	O(39)-Si(1)-O(37)	109.4(6)	O(38)-Si(1)-O(40)	108.6(6)	
Nb(3)-O(34)-Nb(2)	148.7(7)	O(38)-Si(1)-O(37)	110.8(7)	O(37)-Si(1)-O(40)	110.0(6)	

O_t: terminal oxygen, O_b: doubly-bridged O atoms, O_c: quadruply-bridged O atoms.

conventional distorted octahedra (*ca.* C_{4v} local symmetry) typical of the "type I" octahedral geometry in Keggin polyanions.

The three peroxo groups of A- α -1 as a grouping or unit are not equivalent in the solid state. That is, they are not related by a C₃ axis of symmetry trisecting the three NbO₇ units. Given this point, there are 3 possible general descriptions of the orientation of these three terminal peroxo groups to one another. First, they could be related by an approximate (but non-crystallographically imposed) C₃ axis. This would lead to the highest symmetry arrangement, *ca.* C₃ (or C_{3v}). They could be related by a mirror plane (2 of the 3 peroxo groups have similar orientations, but the third peroxo group has a different orientation). This would lead to *ca.* C_s symmetry of the grouping. Finally, they could be best described as having no apparent symmetry relationship with respect to one another (C₁ symmetry). The structure of A- α -1 indicates that while the three peroxo groups appear to approach a C₃ orientation with respect to one another, they are rigorously C₁.

The two-line ¹⁸³W NMR (0.1 M lithiated A- α -Cs₈H₁ in D₂O; δ (intensity): -111.4 (6) and -138.4 (3)) spectrum shown in Figure 2 clearly indicates that A- α -Cs₈H₁ possesses apparent C_{3v} symmetry in solution. Yet, it is clear from the somewhat different orientations of the peroxo groups relative to one another in the crystal structure of A- α -Cs₈H₁ that the polyanion unit as a whole has lower symmetry than C_{3v}, and that the ¹⁸³W NMR spectrum should have more than two lines if the solid state symmetry is maintained in solution. There are two possible explanations for this. The first is that the peroxo groups are fluxional and equilibrating on the ¹⁸³W NMR time scale so that all are equivalent, and thus the proximal or "belt" tungsten atoms are all equivalent. This would lead to the observed two-line spectrum. The second is that the orientations of the three peroxo groups are, in fact, different as they are in the solid state, but that this difference represents such a small perturbation on the tungsten nuclei that they are effectively magnetically equivalent (*i.e.* the I = 1/2 ¹⁸³W nuclei "see" effectively the same magnetic environment). The first explanation is extremely unlikely because fluxionality of this type hasn't been seen in




Figure 2. ¹⁸³W NMR spectrum of lithiated A- α -Cs₈H₁[Si(NbO₂)₃-W₃O₃₇] (A- α -Cs₈H₁) (0.1 M) in D₂O with 3 equivalents of aqueous H₂O₂.

any other peroxo-containing polyoxometalate complexes nor the polyperoxometalates^{14,15} nor is a facile chemistry involving peroxo group equilibration by detachment or reattachment on such a rapid time scale reasonable. Furthermore, if such dynamic processes were operable, one would expect to see significantly larger thermal ellipsoids than those exhibited by A- α -1 (see Figure 1) or by those exhibited in many other X-ray structures of d⁰ polyperoxometalate species. Therefore, we are left to conclude that apparent C_{3v} symmetry of A- α -1 derives from the fact that the orientation of the peroxo groups has too little effect on the tungsten nuclei to significantly impact the chemical shifts of these atoms.

POM anions are linked together through Cs⁻ cations and crystalline water molecules. All the surface O atoms of the POM anion, including the terminal oxo and peroxo, the bridging μ_2 -O atoms between Nb and W atoms, and the bridging μ_2 -O atoms between W and W atoms except O4, O5, O7, O13 and O25, are involved in forming coordination with Cs⁻ cations of different asymmetric units. The Cs⁻ cations are surrounded by 9 to 12 oxygen atoms from POMs and crystalline water molecules, with Cs-O bond ranging from 2.92 to 3.79 Å. For example, Cs8 is surrounded by two POM anions with the (NbO₂)₃ unit head-on to each other and coordinated to three μ_2 -O-Nb₂ atoms (O31, O32, and O34) and three terminal η^2 -coordinated peroxo oxygens (O33A, O35A and O36A) that come from one POM anion to give a

total coordination number of 12.

Acknowledgment. We wish to thank the Korea Science and Engineering Foundation (KOSEF 2001-1-12200-004-1 to GK), the U.S. Army Research Office (DAAD19-01-1-0593 to CLH), and, in part, the Molecular Design Institute (Office of Naval Research, grant N00014-95-1-1116 to CLH) for the financial support of this research. In addition, sincere thanks go to Dr. Don VanDerveer for X-ray data collection, to Jeff Rhule for the W-183 NMR, and to Dr. Ira Weinstock for help in writing the manuscript.

References

1. For the first synthesis of tetrabutylammonium salt of this peroxyniobium anion, see Droege, M. W.; Finke, R. G. *J. Mol. Catal.* **1991**, *69*, 323.
2. Kim, G.-S.; Judd, D. A.; Hill, C. L.; Schinazi, R. F. *J. Med. Chem.* **1994**, *37*, 816.
3. Rhule, J. T.; Hill, C. L.; Judd, D. A.; Schinazi, R. F. *Chem. Rev.* **1998**, *98*, 327.
4. Kim, G.-S.; Zeng, H.; VanDerveer, D.; Hill, C. L. *Angew. Chem. Int. Ed.* **1999**, *30*, 3205.
5. Kim, G.-S.; Zeng, H.; Weinstock, I. A.; Rhule, J. T.; VanDerveer, D.; Hill, C. L. *Inorg. Chem.*, in submission.
6. See also Kim, G.-S.; Zeng, H.; Rhule, J. T.; Weinstock, I. A.; Hill, C. L. *J. Chem. Soc., Chem. Commun.* **1999**, *17*, 1651.
7. Judd, D. A.; Chen, Q.; Campana, C. F.; Hill, C. L. *J. Am. Chem. Soc.* **1997**, *119*, 5461.
8. Klemperer, W. G. In *Inorganic Syntheses*; John Wiley and Sons, Inc.: New York, 1990; Vol. 27, p 71.
9. Dabbabi, M.; Boyer, M. *J. Inorg. Nucl. Chem.* **1976**, *38*, 1011.
10. *SMART, version 4.05*; Siemens Analytical X-ray Systems, Inc.: Madison, WI, 1995.
11. *SAINTE, version 4.05*; Siemens Analytical X-ray Systems, Inc.: Madison, WI, 1995.
12. *SADABS: Siemens area detector absorption correction program*; Siemens Analytical X-ray Systems, Inc.: Madison, WI, 1995.
13. Sheldrick, G. M. *SHELX-97 program package for structure solution and refinement*, University Goettingen: Germany, 1997.
14. There are many structures of low nuclearity complexes with as or more η^2 -coordinated peroxy groups than oxo groups on the d^0 metal centers. These complexes, including "Venturello tetramers" (e.g. $\{\text{PO}_4[\text{WO}(\text{O}_2)_2]_4\}^{2-}$) and "Brégeault dimers" (e.g. $\{\text{HPO}_4[\text{WO}(\text{O}_2)_2]_2\}^{2-}$) are low nuclearity polyanions and are better regarded as class of complexes distinct from conventional POMs.
15. Dickman, M. H.; Pope, M. T. *Chem. Rev.* **1994**, *94*, 569.
16. Bas-Serra, J.; Todorut, I.; Jameson, G. B.; Acerete, R.; Baker, L. C. W. Abstract of Paper No. 77b Presented at the Intl. Chem. Congress of Pacific Basin Societies, Honolulu, 1989.
17. Server-Carrió, J.; Bas-Serra, J.; González-Núñez, M. E.; García-Gastaldí, A.; Jameson, G. B.; Baker, L. C. W.; Acerete, R. *J. Am. Chem. Soc.* **1999**, *121*, 911.
18. Harrup, M. K.; Kim, G.-S.; Zeng, H.; Johnson, R. P.; VanDerveer, D.; Hill, C. L. *Inorg. Chem.* **1998**, *37*, 5550.
19. Vaska, L. *Acc. Chem. Res.* **1976**, *9*, 175.
20. Mathem, G.; Weiss, R. *Acta Cryst.* **1971**, *B27*, 1598.
21. Roch, M.; Weber, J.; Williams, A. F. *Inorg. Chem.* **1984**, *23*, 4572.