

PREOPEN SETS IN SMOOTH BITOPOLOGICAL SPACES

EUN PYO LEE

ABSTRACT. In this paper, we introduce the notions of $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen sets and fuzzy pairwise (r, s) -precontinuous mappings in smooth bitopological spaces and then we investigate some of their characteristic properties.

1. Introduction

The concept of fuzzy sets was introduced by Zadeh [13] in his classical paper. Using the concept of fuzzy sets Chang [2] introduced fuzzy topological spaces and several other authors continued the investigation of such spaces. Chattopadhyay et al. [4] and Ramadan [9] introduced new definition of smooth topological spaces as a generalization of fuzzy topological spaces. Kandil [6] introduced and studied the notion of fuzzy bitopological spaces as a natural generalization of fuzzy topological spaces. Lee et al. [7] introduced the concept of smooth bitopological spaces as a generalization of smooth topological spaces and Kandil's fuzzy bitopological spaces.

In this paper, we introduce the concepts of $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen sets and fuzzy pairwise (r, s) -precontinuous mappings in smooth bitopological spaces and then we investigate some of their characteristic properties.

2. Preliminaries

In this paper, I will denote the unit interval $[0, 1]$ of the real line and

Received February 18, 2002.

2000 Mathematics Subject Classification: 54A40.

Key words and phrases: smooth bitopological spaces, preopen sets, pairwise precontinuous mappings.

$I_0 = (0, 1]$. For a set X , I^X denotes the collection of all mappings from X to I . A member μ of I^X is called a fuzzy set of X . By $\tilde{0}$ and $\tilde{1}$ we denote constant mappings on X with value 0 and 1, respectively. For any $\mu \in I^X$, μ^c denotes the complement $\tilde{1} - \mu$. All other notations are the standard notations of fuzzy set theory.

A *Chang's fuzzy topology* on X [2] is a family T of fuzzy sets in X which satisfies the following properties:

- (1) $\tilde{0}, \tilde{1} \in T$.
- (2) If $\mu_1, \mu_2 \in T$ then $\mu_1 \wedge \mu_2 \in T$.
- (3) If $\mu_k \in T$ for each k , then $\bigvee \mu_k \in T$.

The pair (X, T) is called a *Chang's fuzzy topological space*. Members of T are called T -fuzzy open sets of X and their complements T -fuzzy closed sets of X .

A system (X, T_1, T_2) consisting of a set X with two Chang's fuzzy topologies T_1 and T_2 on X is called a *Kandil's fuzzy bitopological space*.

A *smooth topology* on X [4, 9] is a mapping $\mathcal{T} : I^X \rightarrow I$ which satisfies the following properties:

- (1) $\mathcal{T}(\tilde{0}) = \mathcal{T}(\tilde{1}) = 1$.
- (2) $\mathcal{T}(\mu_1 \wedge \mu_2) \geq \mathcal{T}(\mu_1) \wedge \mathcal{T}(\mu_2)$.
- (3) $\mathcal{T}(\bigvee \mu_k) \geq \bigwedge \mathcal{T}(\mu_k)$.

The pair (X, \mathcal{T}) is called a *smooth topological space*. For $r \in I_0$, we call μ a \mathcal{T} -fuzzy r -open set of X if $\mathcal{T}(\mu) \geq r$ and μ a \mathcal{T} -fuzzy r -closed set of X if $\mathcal{T}(\mu^c) \geq r$.

A system $(X, \mathcal{T}_1, \mathcal{T}_2)$ consisting of a set X with two smooth topologies \mathcal{T}_1 and \mathcal{T}_2 on X is called a *smooth bitopological space*. Throughout this paper the indices i, j take values in $\{1, 2\}$ and $i \neq j$.

Let (X, \mathcal{T}) be a smooth topological space. For each $r \in I_0$, an r -cut

$$\mathcal{T}_r = \{\mu \in I^X \mid \mathcal{T}(\mu) \geq r\}$$

is a Chang's fuzzy topology on X .

Let (X, T) be a Chang's fuzzy topological space and $r \in I_0$. Then a smooth topology $T^r : I^X \rightarrow I$ is defined by

$$T^r(\mu) = \begin{cases} 1 & \text{if } \mu = \tilde{0}, \tilde{1}, \\ r & \text{if } \mu \in T - \{\tilde{0}, \tilde{1}\}, \\ 0 & \text{otherwise.} \end{cases}$$

Hence, we obtain that if $(X, \mathcal{T}_1, \mathcal{T}_2)$ is a smooth bitopological space and $r, s \in I_0$, then $(X, (\mathcal{T}_1)_r, (\mathcal{T}_2)_s)$ is a Kandil's fuzzy bitopological space. Also, if $(X, \mathcal{T}_1, \mathcal{T}_2)$ is a Kandil's fuzzy bitopological space and $r, s \in I_0$, then $(X, (\mathcal{T}_1)^r, (\mathcal{T}_2)^s)$ is a smooth bitopological space.

DEFINITION 2.1 ([8]). Let (X, \mathcal{T}) be a smooth topological space. For each $r \in I_0$ and for each $\mu \in I^X$, the *fuzzy r -closure* is defined by

$$\mathcal{T}\text{-Cl}(\mu, r) = \bigwedge \{ \rho \mid \mu \leq \rho, \mathcal{T}(\rho^c) \geq r \}$$

and the *fuzzy r -interior*

$$\mathcal{T}\text{-Int}(\mu, r) = \bigvee \{ \rho \mid \mu \geq \rho, \mathcal{T}(\rho) \geq r \}.$$

THEOREM 2.2 ([8]). For a fuzzy set μ of a smooth topological space (X, \mathcal{T}) and $r \in I_0$, we have:

- (1) $\mathcal{T}\text{-Int}(\mu, r)^c = \mathcal{T}\text{-Cl}(\mu^c, r)$.
- (2) $\mathcal{T}\text{-Cl}(\mu, r)^c = \mathcal{T}\text{-Int}(\mu^c, r)$.

DEFINITION 2.3 ([7]). Let μ be a fuzzy set of a smooth bitopological space $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $r, s \in I_0$. Then μ is said to be

- (1) a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -semiopen set if there is a \mathcal{T}_i -fuzzy r -open set ρ in X such that $\rho \leq \mu \leq \mathcal{T}_j\text{-Cl}(\rho, s)$,
- (2) a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -semiclosed set if there is a \mathcal{T}_i -fuzzy r -closed set ρ in X such that $\mathcal{T}_j\text{-Int}(\rho, s) \leq \mu \leq \rho$.

DEFINITION 2.4 ([7]). Let $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ be a mapping from a smooth bitopological space X to another smooth bitopological space Y and $r, s \in I_0$. Then f is called a fuzzy pairwise (r, s) -continuous $((r, s)$ -open and (r, s) -closed, respectively) mapping if the induced mapping $f : (X, \mathcal{T}_1) \rightarrow (Y, \mathcal{U}_1)$ is a fuzzy r -continuous (r -open and r -closed, respectively) mapping and the induced mapping $f : (X, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_2)$ is a fuzzy s -continuous (s -open and s -closed, respectively) mapping.

3. $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen sets

DEFINITION 3.1. Let μ be a fuzzy set of a smooth bitopological space $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $r, s \in I_0$. Then μ is said to be

- (1) a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen set if $\mu \leq \mathcal{T}_i\text{-Int}(\mathcal{T}_j\text{-Cl}(\mu, s), r)$,
- (2) a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preclosed set if $\mathcal{T}_i\text{-Cl}(\mathcal{T}_j\text{-Int}(\mu, s), r) \leq \mu$.

THEOREM 3.2. *Let μ be a fuzzy set of a smooth bitopological space $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $r, s \in I_0$. Then the following statements are equivalent:*

- (1) μ is a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen set.
- (2) μ^c is a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preclosed set.

PROOF. It follows from Theorem 2.2. □

THEOREM 3.3. (1) *Any union of $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen sets is a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen set.*

(2) *Any intersection of $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preclosed sets is a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preclosed set.*

PROOF. (1) Let $\{\mu_k\}$ be a collection of $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen sets. Then for each k , $\mu_k \leq \mathcal{T}_i\text{-Int}(\mathcal{T}_j\text{-Cl}(\mu_k, s), r)$. So

$$\bigvee \mu_k \leq \bigvee \mathcal{T}_i\text{-Int}(\mathcal{T}_j\text{-Cl}(\mu_k, s), r) \leq \mathcal{T}_i\text{-Int}(\mathcal{T}_j\text{-Cl}(\bigvee \mu_k, s), r).$$

Thus $\bigvee \mu_k$ is a $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen set.

(2) It follows from (1) using Theorem 3.2. □

THEOREM 3.4. *Let μ be a fuzzy set of a smooth bitopological space $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $r, s \in I_0$.*

- (1) *If μ is a \mathcal{T}_1 -fuzzy r -open set of (X, \mathcal{T}_1) , then μ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preopen set of $(X, \mathcal{T}_1, \mathcal{T}_2)$.*
- (2) *If μ is a \mathcal{T}_2 -fuzzy s -open set of (X, \mathcal{T}_2) , then μ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preopen set of $(X, \mathcal{T}_1, \mathcal{T}_2)$.*

PROOF. (1) Let μ be a \mathcal{T}_1 -fuzzy r -open set of (X, \mathcal{T}_1) . Then $\mu = \mathcal{T}_1\text{-Int}(\mu, r)$. Clearly, we have $\mu \leq \mathcal{T}_2\text{-Cl}(\mu, s)$ and hence

$$\mu = \mathcal{T}_1\text{-Int}(\mu, r) \leq \mathcal{T}_1\text{-Int}(\mathcal{T}_2\text{-Cl}(\mu, s), r).$$

Thus μ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preopen set of $(X, \mathcal{T}_1, \mathcal{T}_2)$.

(2) Similar to (1). □

But the converses in the above theorem need not be true which is shown by the following example.

EXAMPLE 3.5. Let $X = \{x, y\}$ and μ_1, μ_2, μ_3 and μ_4 be fuzzy sets of X defined as

$$\mu_1(x) = 0.2, \quad \mu_1(y) = 0.4;$$

$$\mu_2(x) = 0.6, \quad \mu_2(y) = 0.3;$$

$$\mu_3(x) = 0.2, \quad \mu_3(y) = 0.3;$$

and

$$\mu_4(x) = 0.5, \quad \mu_4(y) = 0.2.$$

Define $\mathcal{T}_1 : I^X \rightarrow I$ and $\mathcal{T}_2 : I^X \rightarrow I$ by

$$\mathcal{T}_1(\mu) = \begin{cases} 1 & \text{if } \mu = \tilde{0}, \tilde{1}, \\ \frac{1}{2} & \text{if } \mu = \mu_1, \\ 0 & \text{otherwise;} \end{cases}$$

and

$$\mathcal{T}_2(\mu) = \begin{cases} 1 & \text{if } \mu = \tilde{0}, \tilde{1}, \\ \frac{1}{3} & \text{if } \mu = \mu_2, \\ 0 & \text{otherwise.} \end{cases}$$

Then clearly $(\mathcal{T}_1, \mathcal{T}_2)$ is a smooth bitopology on X . Note that

$$\mathcal{T}_1\text{-Int}(\mathcal{T}_2\text{-Cl}(\mu_3, \frac{1}{3}), \frac{1}{2}) = \mathcal{T}_1\text{-Int}(\mu_2^c, \frac{1}{2}) = \mu_1 \geq \mu_3.$$

Thus μ_3 is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy $(\frac{1}{2}, \frac{1}{3})$ -preopen set which is not a \mathcal{T}_1 -fuzzy $\frac{1}{2}$ -open set. Also we have

$$\mathcal{T}_2\text{-Int}(\mathcal{T}_1\text{-Cl}(\mu_4, \frac{1}{2}), \frac{1}{3}) = \mathcal{T}_2\text{-Int}(\mu_1^c, \frac{1}{3}) = \mu_2 \geq \mu_4.$$

Hence μ_4 is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy $(\frac{1}{3}, \frac{1}{2})$ -preopen set which is not a \mathcal{T}_2 -fuzzy $\frac{1}{3}$ -open set.

REMARK 3.6. That $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -semiopen sets and $(\mathcal{T}_i, \mathcal{T}_j)$ -fuzzy (r, s) -preopen sets are independent notions is shown by the following example.

EXAMPLE 3.7. Let $X = \{x, y\}$ and $\mu_1, \mu_2, \mu_3, \mu_4, \mu_5$ and μ_6 be fuzzy sets of X defined as

$$\mu_1(x) = 0.1, \quad \mu_1(y) = 0.7;$$

$$\mu_2(x) = 0.8, \quad \mu_2(y) = 0.2;$$

$$\mu_3(x) = 0, \quad \mu_3(y) = 0.6;$$

$$\mu_4(x) = 0.1, \quad \mu_4(y) = 0.8;$$

$$\mu_5(x) = 0.5, \quad \mu_5(y) = 0.6;$$

and

$$\mu_6(x) = 0.9, \quad \mu_6(y) = 0.2.$$

Define $\mathcal{T}_1 : I^X \rightarrow I$ and $\mathcal{T}_2 : I^X \rightarrow I$ by

$$\mathcal{T}_1(\mu) = \begin{cases} 1 & \text{if } \mu = \tilde{0}, \tilde{1}, \\ \frac{1}{2} & \text{if } \mu = \mu_1, \\ 0 & \text{otherwise;} \end{cases}$$

and

$$\mathcal{T}_2(\mu) = \begin{cases} 1 & \text{if } \mu = \tilde{0}, \tilde{1}, \\ \frac{1}{3} & \text{if } \mu = \mu_2, \\ 0 & \text{otherwise.} \end{cases}$$

Then clearly $(\mathcal{T}_1, \mathcal{T}_2)$ is a smooth bitopology on X . The fuzzy set μ_3 is $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy $(\frac{1}{2}, \frac{1}{3})$ -preopen which is not $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy $(\frac{1}{2}, \frac{1}{3})$ -semiopen and μ_4 is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy $(\frac{1}{2}, \frac{1}{3})$ -semiopen set which is not a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy $(\frac{1}{2}, \frac{1}{3})$ -preopen set. Also μ_5 is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy $(\frac{1}{3}, \frac{1}{2})$ -preopen set which is not a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy $(\frac{1}{3}, \frac{1}{2})$ -semiopen set and μ_6 is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy $(\frac{1}{3}, \frac{1}{2})$ -semiopen set which is not a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy $(\frac{1}{3}, \frac{1}{2})$ -preopen set.

4. Fuzzy pairwise (r, s) -precontinuous mappings

DEFINITION 4.1. Let $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ be a mapping from a smooth bitopological space X to another smooth bitopological space Y and $r, s \in I_0$. Then f is called

- (1) a *fuzzy pairwise (r, s) -precontinuous* mapping if $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preopen set of X for each \mathcal{U}_1 -fuzzy r -open

- set μ of Y and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preopen set of X for each \mathcal{U}_2 -fuzzy s -open set ν of Y ,
- (2) a *fuzzy pairwise (r, s) -preopen mapping* if $f(\rho)$ is a $(\mathcal{U}_1, \mathcal{U}_2)$ -fuzzy (r, s) -preopen set of Y for each \mathcal{T}_1 -fuzzy r -open set ρ of X and $f(\lambda)$ is a $(\mathcal{U}_2, \mathcal{U}_1)$ -fuzzy (s, r) -preopen set of Y for each \mathcal{T}_2 -fuzzy s -open set λ of X ,
- (3) a *fuzzy pairwise (r, s) -preclosed mapping* if $f(\rho)$ is a $(\mathcal{U}_1, \mathcal{U}_2)$ -fuzzy (r, s) -preclosed set of Y for each \mathcal{T}_1 -fuzzy r -closed set ρ of X and $f(\lambda)$ is a $(\mathcal{U}_2, \mathcal{U}_1)$ -fuzzy (s, r) -preclosed set of Y for each \mathcal{T}_2 -fuzzy s -closed set λ of X .

REMARK 4.2. It is obvious that every fuzzy pairwise (r, s) -continuous mapping is also a fuzzy pairwise (r, s) -precontinuous mapping. But the converse need not be true which is shown by the following example.

EXAMPLE 4.3. (1) *A fuzzy pairwise (r, s) -precontinuous mapping need not be a fuzzy pairwise (r, s) -continuous mapping.*

Let $(X, \mathcal{T}_1, \mathcal{T}_2)$ be a smooth bitopological space as described in Example 3.5. Define $\mathcal{U}_1 : I^X \rightarrow I$ and $\mathcal{U}_2 : I^X \rightarrow I$ by

$$\mathcal{U}_1(\mu) = \begin{cases} 1 & \text{if } \mu = \tilde{0}, \tilde{1}, \\ \frac{1}{2} & \text{if } \mu = \mu_3, \\ 0 & \text{otherwise;} \end{cases}$$

and

$$\mathcal{U}_2(\mu) = \begin{cases} 1 & \text{if } \mu = \tilde{0}, \tilde{1}, \\ 0 & \text{otherwise.} \end{cases}$$

Then clearly $(\mathcal{U}_1, \mathcal{U}_2)$ is a smooth bitopology on X . Consider the mapping $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (X, \mathcal{U}_1, \mathcal{U}_2)$ defined by $f(x) = x$ and $f(y) = y$. Then f is a fuzzy pairwise $(\frac{1}{2}, \frac{1}{3})$ -precontinuous mapping which is not a fuzzy pairwise $(\frac{1}{2}, \frac{1}{3})$ -continuous mapping.

(2) *A fuzzy pairwise (r, s) -preopen mapping need not be a fuzzy pairwise (r, s) -open mapping.*

Let $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $(X, \mathcal{U}_1, \mathcal{U}_2)$ be smooth bitopological spaces as described in (1). Consider the mapping $f : (X, \mathcal{U}_1, \mathcal{U}_2) \rightarrow (X, \mathcal{T}_1, \mathcal{T}_2)$ defined by $f(x) = x$ and $f(y) = y$. Then f is a fuzzy pairwise $(\frac{1}{2}, \frac{1}{3})$ -preopen mapping which is not a fuzzy pairwise $(\frac{1}{2}, \frac{1}{3})$ -open mapping.

(3) *A fuzzy pairwise (r, s) -preclosed mapping need not be a fuzzy pairwise (r, s) -closed mapping.*

Let $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $(X, \mathcal{U}_1, \mathcal{U}_2)$ be smooth bitopological spaces as described in (1). Consider the mapping $f : (X, \mathcal{U}_1, \mathcal{U}_2) \rightarrow (X, \mathcal{T}_1, \mathcal{T}_2)$

defined by $f(x) = x$ and $f(y) = y$. Then f is a fuzzy pairwise $(\frac{1}{2}, \frac{1}{3})$ -preclosed mapping which is not a fuzzy pairwise $(\frac{1}{2}, \frac{1}{3})$ -closed mapping.

THEOREM 4.4. *Let $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ be a mapping and $r, s \in I_0$. Then the following statements are equivalent:*

- (1) f is a fuzzy pairwise (r, s) -precontinuous mapping.
- (2) $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preclosed set of X for each \mathcal{U}_1 -fuzzy r -closed set μ of Y and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preclosed set of X for each \mathcal{U}_2 -fuzzy s -closed set ν of Y .
- (3) For each fuzzy set μ of Y ,

$$\mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(f^{-1}(\mu), s), r) \leq f^{-1}(\mathcal{U}_1\text{-Cl}(\mu, r))$$

and

$$\mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(f^{-1}(\mu), r), s) \leq f^{-1}(\mathcal{U}_2\text{-Cl}(\mu, s)).$$

- (4) For each fuzzy set ρ of X ,

$$f(\mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(\rho, s), r) \leq \mathcal{U}_1\text{-Cl}(f(\rho), r)$$

and

$$f(\mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(\rho, r), s) \leq \mathcal{U}_2\text{-Cl}(f(\rho), s).$$

PROOF. (1) \Rightarrow (2) Let μ be any \mathcal{U}_1 -fuzzy r -closed set and ν any \mathcal{U}_2 -fuzzy s -closed set of Y . Then μ^c is a \mathcal{U}_1 -fuzzy r -open set and ν^c is a \mathcal{U}_2 -fuzzy s -open set of Y . Since f is a fuzzy pairwise (r, s) -precontinuous mapping, $f^{-1}(\mu^c)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preopen set and $f^{-1}(\nu^c)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preopen set of X . But $f^{-1}(\mu^c) = f^{-1}(\mu)^c$ and $f^{-1}(\nu^c) = f^{-1}(\nu)^c$. By Theorem 3.2, $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preclosed set and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preclosed set of X .

(2) \Rightarrow (1) Let μ be any \mathcal{U}_1 -fuzzy r -open set and ν any \mathcal{U}_2 -fuzzy s -open set of Y . Then μ^c is a \mathcal{U}_1 -fuzzy r -closed set and ν^c is a \mathcal{U}_2 -fuzzy s -closed set of Y . By (2), $f^{-1}(\mu^c)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preclosed set and $f^{-1}(\nu^c)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preclosed set of X . But $f^{-1}(\mu^c) = f^{-1}(\mu)^c$ and $f^{-1}(\nu^c) = f^{-1}(\nu)^c$. By Theorem 3.2, $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preopen set and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preopen set of X . Thus f is a fuzzy pairwise (r, s) -precontinuous mapping.

(2) \Rightarrow (3) Let μ be any fuzzy set of Y . Then $\mathcal{U}_1\text{-Cl}(\mu, r)$ is a \mathcal{U}_1 -fuzzy r -closed set and $\mathcal{U}_2\text{-Cl}(\mu, s)$ is a \mathcal{U}_2 -fuzzy s -closed set of Y . By (2),

$f^{-1}(\mathcal{U}_1\text{-Cl}(\mu, r))$ is $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preclosed and $f^{-1}(\mathcal{U}_2\text{-Cl}(\mu, s))$ is $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preclosed of X . Thus

$$\begin{aligned} f^{-1}(\mathcal{U}_1\text{-Cl}(\mu, r)) &\geq \mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(f^{-1}(\mathcal{U}_1\text{-Cl}(\mu, r)), s), r) \\ &\geq \mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(f^{-1}(\mu), s), r) \end{aligned}$$

and

$$\begin{aligned} f^{-1}(\mathcal{U}_2\text{-Cl}(\mu, s)) &\geq \mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(f^{-1}(\mathcal{U}_2\text{-Cl}(\mu, s)), r), s) \\ &\geq \mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(f^{-1}(\mu), r), s). \end{aligned}$$

(3) \Rightarrow (4) Let ρ be any fuzzy set of X . Then $f(\rho)$ is a fuzzy set of Y . By (3),

$$\begin{aligned} f^{-1}(\mathcal{U}_1\text{-Cl}(f(\rho), r)) &\geq \mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(f^{-1}f(\rho), s), r) \\ &\geq \mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(\rho, s), r) \end{aligned}$$

and

$$\begin{aligned} f^{-1}(\mathcal{U}_2\text{-Cl}(f(\rho), s)) &\geq \mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(f^{-1}f(\rho), r), s) \\ &\geq \mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(\rho, r), s). \end{aligned}$$

Hence

$$\mathcal{U}_1\text{-Cl}(f(\rho), r) \geq f f^{-1}(\mathcal{U}_1\text{-Cl}(f(\rho), r)) \geq f(\mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(\rho, s), r))$$

and

$$\mathcal{U}_2\text{-Cl}(f(\rho), s) \geq f f^{-1}(\mathcal{U}_2\text{-Cl}(f(\rho), s)) \geq f(\mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(\rho, r), s)).$$

(4) \Rightarrow (2) Let μ be any \mathcal{U}_1 -fuzzy r -closed set and ν any \mathcal{U}_2 -fuzzy s -closed set of Y . Then $f^{-1}(\mu)$ and $f^{-1}(\nu)$ are fuzzy sets of X . By (4),

$$\begin{aligned} f(\mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(f^{-1}(\mu), s), r)) &\leq \mathcal{U}_1\text{-Cl}(f f^{-1}(\mu), r) \\ &\leq \mathcal{U}_1\text{-Cl}(\mu, r) = \mu \end{aligned}$$

and

$$\begin{aligned} f(\mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(f^{-1}(\nu), r), s)) &\leq \mathcal{U}_2\text{-Cl}(f f^{-1}(\nu), s) \\ &\leq \mathcal{U}_2\text{-Cl}(\nu, s) = \nu. \end{aligned}$$

So

$$\begin{aligned} \mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(f^{-1}(\mu), s), r) &\leq f^{-1}f(\mathcal{T}_1\text{-Cl}(\mathcal{T}_2\text{-Int}(f^{-1}(\mu), s), r)) \\ &\leq f^{-1}(\mu) \end{aligned}$$

and

$$\begin{aligned} \mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(f^{-1}(\nu), r), s) &\leq f^{-1}f(\mathcal{T}_2\text{-Cl}(\mathcal{T}_1\text{-Int}(f^{-1}(\nu), r), s)) \\ &\leq f^{-1}(\nu). \end{aligned}$$

Therefore $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preclosed set and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preclosed set of X . \square

In general, it need not be true that if f and g are fuzzy pairwise (r, s) -precontinuous ((r, s) -preopen and (r, s) -preclosed, respectively) then so is $g \circ f$. But we have the following theorem.

THEOREM 4.5. *Let $(X, \mathcal{T}_1, \mathcal{T}_2)$, $(Y, \mathcal{U}_1, \mathcal{U}_2)$ and $(Z, \mathcal{V}_1, \mathcal{V}_2)$ be smooth bitopological spaces and let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be mappings and $r, s \in I_0$. Then the following statements are true.*

- (1) *If f is fuzzy pairwise (r, s) -precontinuous and g is fuzzy pairwise (r, s) -continuous then $g \circ f$ is fuzzy pairwise (r, s) -precontinuous.*
- (2) *If f is fuzzy pairwise (r, s) -open and g is fuzzy pairwise (r, s) -preopen then $g \circ f$ is fuzzy pairwise (r, s) -preopen.*
- (3) *If f is fuzzy pairwise (r, s) -closed and g is fuzzy pairwise (r, s) -preclosed then $g \circ f$ is fuzzy pairwise (r, s) -preclosed.*

PROOF. Straightforward. \square

The next two theorems show that a fuzzy pairwise precontinuous mapping is a special case of a fuzzy pairwise (r, s) -precontinuous mapping.

THEOREM 4.6. *Let $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $(Y, \mathcal{U}_1, \mathcal{U}_2)$ be smooth bitopological spaces and let $r, s \in I_0$. Then $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ is fuzzy pairwise (r, s) -precontinuous if and only if $f : (X, (\mathcal{T}_1)_r, (\mathcal{T}_2)_s) \rightarrow (Y, (\mathcal{U}_1)_r, (\mathcal{U}_2)_s)$ is fuzzy pairwise precontinuous.*

PROOF. Let $\mu \in (\mathcal{U}_1)_r$ and $\nu \in (\mathcal{U}_2)_s$. Then $\mathcal{U}_1(\mu) \geq r$ and $\mathcal{U}_2(\nu) \geq s$ and hence μ is a \mathcal{U}_1 -fuzzy r -open set and ν is a \mathcal{U}_2 -fuzzy s -open set of Y . Since $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ is a fuzzy pairwise (r, s) -precontinuous mapping, $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preopen set and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preopen set of $(X, \mathcal{T}_1, \mathcal{T}_2)$. So $f^{-1}(\mu)$ is $((\mathcal{T}_1)_r, (\mathcal{T}_2)_s)$ -fuzzy preopen and $f^{-1}(\nu)$ is $((\mathcal{T}_2)_s, (\mathcal{T}_1)_r)$ -fuzzy preopen of $(X, (\mathcal{T}_1)_r, (\mathcal{T}_2)_s)$. Thus $f : (X, (\mathcal{T}_1)_r, (\mathcal{T}_2)_s) \rightarrow (Y, (\mathcal{U}_1)_r, (\mathcal{U}_2)_s)$ is a fuzzy pairwise precontinuous mapping.

Conversely, let μ be any \mathcal{U}_1 -fuzzy r -open set and ν any \mathcal{U}_2 -fuzzy s -open set of $(Y, \mathcal{U}_1, \mathcal{U}_2)$. Then $\mathcal{U}_1(\mu) \geq r$ and $\mathcal{U}_2(\nu) \geq s$. So $\mu \in (\mathcal{U}_1)_r$ and $\nu \in (\mathcal{U}_2)_s$. Since $f : (X, (\mathcal{T}_1)_r, (\mathcal{T}_2)_s) \rightarrow (Y, (\mathcal{U}_1)_r, (\mathcal{U}_2)_s)$ is a fuzzy pairwise precontinuous mapping, $f^{-1}(\mu)$ is a $((\mathcal{T}_1)_r, (\mathcal{T}_2)_s)$ -fuzzy preopen set and $f^{-1}(\nu)$ is a $((\mathcal{T}_2)_s, (\mathcal{T}_1)_r)$ -fuzzy preopen set of $(X, (\mathcal{T}_1)_r, (\mathcal{T}_2)_s)$. So $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy (r, s) -preopen set and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy (s, r) -preopen set of $(X, \mathcal{T}_1, \mathcal{T}_2)$. Thus $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ is a fuzzy pairwise (r, s) -precontinuous mapping. \square

THEOREM 4.7. *Let $(X, \mathcal{T}_1, \mathcal{T}_2)$ and $(Y, \mathcal{U}_1, \mathcal{U}_2)$ be Kandil's fuzzy bitopological spaces and let $r, s \in I_0$. Then $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ is a fuzzy pairwise precontinuous mapping if and only if $f : (X, (\mathcal{T}_1)^r, (\mathcal{T}_2)^s) \rightarrow (Y, (\mathcal{U}_1)^r, (\mathcal{U}_2)^s)$ is a fuzzy pairwise (r, s) -precontinuous mapping.*

PROOF. Let μ be a $(\mathcal{U}_1)^r$ -fuzzy r -open set and ν a $(\mathcal{U}_2)^s$ -fuzzy s -open set of $(Y, (\mathcal{U}_1)^r, (\mathcal{U}_2)^s)$. Then $(\mathcal{U}_1)^r(\mu) \geq r$ and $(\mathcal{U}_2)^s(\nu) \geq s$ and hence $\mu \in \mathcal{U}_1$ and $\nu \in \mathcal{U}_2$. Since $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ is a fuzzy pairwise precontinuous mapping, $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy preopen set and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy preopen set of $(X, \mathcal{T}_1, \mathcal{T}_2)$. So $f^{-1}(\mu)$ is a $((\mathcal{T}_1)^r, (\mathcal{T}_2)^s)$ -fuzzy (r, s) -preopen set and $f^{-1}(\nu)$ is a $((\mathcal{T}_2)^s, (\mathcal{T}_1)^r)$ -fuzzy (s, r) -preopen set of $(X, (\mathcal{T}_1)^r, (\mathcal{T}_2)^s)$. Thus $f : (X, (\mathcal{T}_1)^r, (\mathcal{T}_2)^s) \rightarrow (Y, (\mathcal{U}_1)^r, (\mathcal{U}_2)^s)$ is a fuzzy pairwise (r, s) -precontinuous mapping.

Conversely, let $\mu \in \mathcal{U}_1$ and $\nu \in \mathcal{U}_2$. Then $(\mathcal{U}_1)^r(\mu) \geq r$ and $(\mathcal{U}_2)^s(\nu) \geq s$, and hence μ is a $(\mathcal{U}_1)^r$ -fuzzy r -open set and ν is a $(\mathcal{U}_2)^s$ -fuzzy s -open set of Y . Since $f : (X, (\mathcal{T}_1)^r, (\mathcal{T}_2)^s) \rightarrow (Y, (\mathcal{U}_1)^r, (\mathcal{U}_2)^s)$ is a fuzzy pairwise (r, s) -precontinuous mapping, $f^{-1}(\mu)$ is a $((\mathcal{T}_1)^r, (\mathcal{T}_2)^s)$ -fuzzy (r, s) -preopen set and $f^{-1}(\nu)$ is a $((\mathcal{T}_2)^s, (\mathcal{T}_1)^r)$ -fuzzy (s, r) -preopen set of $(X, (\mathcal{T}_1)^r, (\mathcal{T}_2)^s)$. So $f^{-1}(\mu)$ is a $(\mathcal{T}_1, \mathcal{T}_2)$ -fuzzy preopen set and $f^{-1}(\nu)$ is a $(\mathcal{T}_2, \mathcal{T}_1)$ -fuzzy preopen set of $(X, \mathcal{T}_1, \mathcal{T}_2)$. Thus $f : (X, \mathcal{T}_1, \mathcal{T}_2) \rightarrow (Y, \mathcal{U}_1, \mathcal{U}_2)$ is a fuzzy pairwise precontinuous mapping. \square

References

- [1] K. K. Azad, *On fuzzy semicontinuity, fuzzy almost continuity and fuzzy weakly continuity*, J. Math. Anal. Appl. **82** (1981), 14–32.
- [2] C. L. Chang, *Fuzzy topological spaces*, J. Math. Anal. Appl. **24** (1968), 182–190.
- [3] K. C. Chattopadhyay and S. K. Samanta, *Fuzzy topology : Fuzzy closure operator, fuzzy compactness and fuzzy connectedness*, Fuzzy Sets and Systems **54** (1993), 207–212.
- [4] K. C. Chattopadhyay, R. N. Hazra and S. K. Samanta, *Gradation of openness : fuzzy topology*, Fuzzy Sets and Systems **49** (1992), 237–242.
- [5] R. N. Hazra, S. K. Samanta and K. C. Chattopadhyay, *Fuzzy topology redefined*, Fuzzy Sets and Systems **45** (1992), 79–82.
- [6] A. Kandil, *Biproximities and fuzzy bitopological spaces*, Simon Stevin **63** (1989), 45–66.
- [7] E. P. Lee, Y. B. Im and H. Han, *Semiopen sets on smooth bitopological spaces*, Far East J. Math. Sci. **3** (2001), 493–521.
- [8] S. J. Lee and E. P. Lee, *Fuzzy r -preopen sets and fuzzy r -precontinuous maps*, Bull. Korean Math. Soc. **36** (1999), no. 1, 91–108.
- [9] A. A. Ramadan, *Smooth topological spaces*, Fuzzy Sets and Systems **48** (1992), 371–375.
- [10] S. Sampath Kumar, *Semi-open sets, semi-continuity and semi-open mappings in fuzzy bitopological spaces*, Fuzzy Sets and Systems **64** (1994), 421–426.
- [11] ———, *On fuzzy pairwise α -continuity and fuzzy pairwise pre-continuity*, Fuzzy Sets and Systems **62** (1994), 231–238.
- [12] T. H. Yalvac, *Semi-interior and semi-closure of a fuzzy set*, J. Math. Anal. Appl. **132** (1988), 356–364.
- [13] L. A. Zadeh, *Fuzzy sets*, Inform. and Control **8** (1965), 338–353.

Department of Mathematics
Seonam University
Namwon 590-711, Korea
E-mail: eplee@seonam.ac.kr