

## Fluorescence Quenching of Bis-msb by Carbon Tetrachloride in Different Solvents

J. Thipperudrappa, D. S. Biradar, M.T. Lagare, S. M. Hanagodimath\*,  
S. R. Inamdar<sup>a</sup> and J. S. Kadadevaramath<sup>a</sup>

Department of Physics, Gulbarga University, Gulbarga - 585 106, Karnataka

<sup>a</sup>Department of Physics, Karnatak University, Dharwad - 580 003, Karnataka

Fluorescence quenching of 1,4-bis [2-(2-methylphenyl) ethenyl]-benzene (Bis-MSB) by carbon tetrachloride in five different solvents namely hexane, cyclohexane, toluene, benzene and dioxane has been carried out at room temperature with a view to understand the quenching mechanisms. The Stern-Volmer plot has been found to be non-linear with a positive deviation for all the solvents studied. In order to interpret these results we have invoked the Ground state complex and Sphere of action static quenching models. Using these models various rate parameters have been determined. The magnitudes of these parameters imply that sphere of action static quenching model agrees well with the experimental results. Hence the positive deviation in the Stern-Volmer plots is attributed to the static and dynamic quenching. Further, with the use of Finite Sink approximation model, it was possible to check whether these bimolecular reactions are diffusion limited and to estimate independently distance parameter  $R'$  and mutual diffusion coefficient  $D$ . Finally an effort has been made to correlate the values of  $R'$  and  $D$  with the values of the encounter distance  $R$  and the mutual diffusion coefficient  $D$  determined using the Edward's empirical relation and Stokes-Einstein relation.

**key words:** Fluorescence quenching, BIS-MSB, solvent effect, Finite sink approximation, model, Sphere of action static quenching model

### INTRODUCTION

Fluorescence quenching of organic molecules in solution by various quenchers like aniline, bromobenzene, carbon tetrachloride, ethyltrithiocarbonate, halide ions, metal ions etc. has been studied by several investigators [1-11]. The study has been mainly to understand the nature of bimolecular reactions taking place both under steady state and transient conditions. This study has not only been of importance in physical sciences but also in chemical, biological and medical sciences [12-18]. The fluorescence yield in bimolecular liquid systems is hindered due to several mechanisms such as static and dynamic quenching, excimer and exciplex formation, charge transfer processes, etc. One of the well known experimental techniques used to study the role of fluorescence quenching is to determine the quenching rate parameter using Stern-Volmer (S-V) plots. If the quenching mechanism is mainly due to dynamic process, then it will be largely due to diffusion in which case diffusion rate parameter  $k_d$  equals the quenching rate parameter  $k_q (=K_{sv}/\tau)$  where  $K_{sv}$  is the slope of the linear S-V plot and  $\tau$  is the decay time of the solute in the absence of the quencher. On the other hand, if the experimental

results do not simulate with the simple linear S-V equation, then it may be due to one of the above processes other than or along with diffusion processes.

In the present study we have used steady state experimental setup to investigate the quenching of 1,4-bis [2-(2-methylphenyl) ethenyl]-benzene (Bis-MSB) by carbon tetrachloride ( $CCl_4$ ) in five different organic solvents with a view to understand the nature of quenching mechanism involved in these systems.

### THEORY

The dynamic process in which quenching mechanism is mainly due to collision is governed by the linear S-V equation

$$\frac{I_0}{I} = 1 + K_{sv}[Q] \quad (1)$$

where  $I_0$  is the fluorescence intensity of the solute in the absence of quencher,  $I$  is the fluorescence intensity in the presence of quencher,  $K_{sv}$  ( $= k_q\tau$ ,  $k_q$  is quenching rate parameter) is the S-V constant and  $\tau$  is the life time of the solute molecule in the absence of quencher. Equation (1) is applicable as long as the experimental results show linear variation. The departure from linearity in S-V plot, suggests that quenching mechanism is not purely collisional and this may be attributed either to the ground state complex formation or to the sphere of action static quenching model [7].

\*To whom correspondence should be addressed.

E-mail : smhmath@rediffmail.com

Received September 2, 2003; Accepted March 19, 2004

In order to see whether the ground state complex formation is partly playing a role, one can use extended S-V equation [19, 20] given by

$$[(I_o/I) - 1]/[Q] = (K_{sv} + k_g) + (K_{sv}k_g)[Q] \quad (2)$$

where  $K_{sv}$  and  $k_g$  are S-V and ground state association constants respectively. From equation (2), the values of  $K_{sv}$  and  $k_g$  can easily be determined by least square fit method.

The ground state complex formation takes place if quenching constant  $k_q (= K_{sv}/\tau)$  obtained from equation (2) agrees well with that obtained from the lower portions of the plot according to equation (1) i.e in the low concentration region in the plot of  $I_o/I$  against  $[Q]$ , which is linear. In that case static quenching is very low. Apart from this the ground state complex formation may be noticed if there is a change in the absorption and fluorescence spectra even at higher concentrations of quencher.

In order to see whether static quenching process is playing a role, one can use "Sphere of action static quenching model". According to this model the instantaneous or static quenching occurs if the quencher molecule is very near to, or in contact with the fluorescent molecule at the exact moment it happens to be excited. This was explained by the fact that only a certain fraction 'W' of the excited state is actually quenched by the collisional mechanism. Some molecules in the excited state, the fraction of which is (1-W), are de-activated almost instantaneously after being formed, because a quencher molecule happens to be randomly positioned in the proximity at the time the molecules are excited and interacts very strongly with them. Several models were employed (Smoluchowski model)[21, 22] to describe this static quenching process, all leading to the following modified form of the S-V equation.

$$(I_o/I) = \frac{1 + K_{sv}[Q]}{W} \quad (3)$$

where  $K_{sv}$  has its usual meaning as explained earlier and  $[Q]$  is the quencher concentration. The Smoluchowski's diffusion controlled equation containing transient term is given by

$$K_d = 4\pi N'RD + 4R^2N'(\pi D)^{1/2}t^{-1/2} \quad (4)$$

where  $N'$  is the Avogadro's number per millimole,  $R$  is the encounter distance i.e. the sum of the radii of the solute and quencher molecules and  $t$  is the time. The retention of the later term of Eq. (4) leads to an additional factor,  $W$  in Eq. (3). This additional factor  $W$  is given by

$$W = e^{-V[Q]} \quad (5)$$

$$\ln(1/W) = V[Q]$$

where  $V$  is the static quenching constant and it represents an active volume element surrounding the excited solute

molecule.

Instantaneous (static) quenching occurs in a randomly distributed system when a quencher happens to reside within a sphere of action with a volume ' $V/N'$ ' and radius ' $r$ ' i.e

$$V/N' = (4\pi r^3)/3 \quad (6)$$

surrounding a solute molecule at the time of excitation.

As  $W$  depends on the quencher concentration  $[Q]$  the S-V plots for a quencher with a high quenching ability generally deviate from linearity. Thus it is worth rewriting Eq. (3) as

$$[1 - (I/I_o)]/[Q] = K_{sv}(I/I_o) + (1 - W)[Q] \quad (7)$$

From equation (7) one can easily calculate  $K_{sv}$  and the values of  $W$  by least square fit method. The static quenching constant  $V$  and the values of ' $r$ ' of sphere of action can also be calculated using equations (5), (6) and (7). According to Andre et al. and Zeng et al.[20, 24], if the distance between the quencher molecule and the excited molecule lies between the encounter distance and the kinetic distance, the static effect takes place especially in the case of steady state experiments irrespective of ground state complex formation provided reactions are limited by diffusion. In order to find whether the reactions are diffusion limited one can invoke the Finite Sink Approximation Model.

#### Finite Sink Approximation Model

Keizer[25-27] has proposed a nonequilibrium statistical modification of the Smoluchowski-Collins-Kimball (SCK) expression to fit the  $I_o/I$  ratio in fluorescence quenching.

In the case of SCK model time dependent rate coefficient  $k(t)$  for diffusion-limited reactions is given by[28]

$$k(t) = a + b \exp(c^2 t) \operatorname{erfc}(ct^{1/2}) \quad (8)$$

where

$$a = k_a \left[ 1 + \frac{k_a}{4\pi N'RD} \right]^{-1} \quad (9)$$

$$b = k_a \left[ 1 + \frac{4\pi N'RD}{k_a} \right]^{-1} \quad (10)$$

$$c = \left[ 1 + \frac{k_a}{4\pi N'RD} \right] \frac{D^{1/2}}{R} \quad (11)$$

Integration of Eq. (8) between the limits  $[Q]$  (8) at  $r \rightarrow \infty$  and  $[Q]$   $[R]$  at  $r = R$  provides the well known expression

$$1/k_d = (1/k_q) + (1/k_a) \quad (12)$$

where  $k_d = 4\pi N'DR$  and  $k_a$  is the activation energy controlled rate constant describing the reaction of encountered pairs at a

reactive distance  $R$  and  $D$  is the sum of the diffusion coefficients of the solute and quencher molecules. Following this expression,  $k_q$  is independent of  $[Q]$ . But for efficient quenching process in liquids,  $k_q$  is often observed to increase with  $[Q]$ . This might be attributed as discussed above to static quenching of solute molecule, in the vicinity of  $[Q]$ , transient effects arising from an initial time dependence of the concentration gradient or combination of them[20].

But if one assumes that only the first encounter is of interest in the case of efficient fluorescence quenching an initial average separation distance  $r_0$  can be defined (sink radius) so that the diffusive region of interest for a first encounter is in the range  $R \leq r \leq r_0$  such that all subsequent encounters are eliminated. Integration of the flux equation between the limits  $[Q](r_0)$  at  $r_0$  and  $[Q](R)$  at  $R$  leads to modification of expression (12) as

$$\frac{1}{k_q} = \frac{1 - (R/r_0)}{k_d} + \frac{1}{k_a} \quad (13)$$

This equation reduces to the reaction limited form ( $k_q = k_a$ ) both for inefficient quenching ( $k_a \ll k_d$ ) and for quenching in pure quenching solvents where  $R = r_0$ . In the diffusion controlled limit ( $k_a \gg k_d$ ) Eq. (13) reduces to

$$k_q = \frac{k_d}{1 - R/r_0} \quad (14)$$

and  $k_q$  depends on the quenching concentration through  $r_0$ . Since the sink radius ( $r_0$ ) is identified to the most probable nearest neighbour initial separation, the appropriate distribution requires that[26]  $r_0 = (2\pi N' [Q])^{-1/3}$ . Replacing  $r_0$  in Eq. (13) and  $k_d$  by its value ( $4\pi N' DR$ ) and dividing everywhere by the fluorescence lifetime of solute in the absence of quencher ( $\tau$ ) one obtains the modified S-V relationship as[20].

$$K_{sv}^{-1} = (K_{sv}^o) - \frac{(2\pi N')^{1/3}}{4\pi N' DR \tau} [Q]^{1/3} \quad (15)$$

where A plot of  $K_{sv}^{-1} = \frac{4\pi N' DR \tau k_a}{4\pi N' DR + k_a}$  against  $[Q]^{1/3}$  becomes

linear with negative slope. Mutual diffusion coefficient  $D$  becomes directly accessible from the slope of the graph exemplified in Eq. (15) and  $K_{sv}$  is obtained at  $[Q] = 0$  regardless of the relative magnitudes of  $k_a$  and  $k_d (=4\pi N' DR)$ , irrespective of quenching is diffusion limited or not. From  $K_{sv}$  we only have access to  $R'$  which is a composite of  $R$  and  $k_a$  through Eq. (16).

$$K_{sv} = 4\pi N' DR' \tau \quad (16)$$

where  $R'$  is the distance parameter and has the same meaning as in the long-time SCK model[20, 28] and is given as

$$R' = R[1 + 4\pi RDN'/k_a]^{-1} \quad (17)$$

Then according to the theory discussed above if  $k_a$  is greater than  $k_d$  (i.e Eq.12) then the reactions are said to be diffusion limited[20] (i.e. for  $R' < R$ ). But for  $R' > R$ , the bimolecular reactions of fluorescence quenching are said to be diffusion limited[28] if the values of  $k_q$  determined from Eq.(7) are greater than  $4\pi N' R' D$ .

## MATERIALS AND METHODS

The solute 1,4-bis[2-(2-methylphenyl) ethenyl]-benzene (Bis-MSB) was obtained from Sigma –Aldrich, USA and is of scintillation grade. The quencher  $CCl_4$  and all solvents were of spectroscopic grade and were used without any further purification. The solutions were prepared keeping the concentration of Bis-MSB fixed ( $5 \times 10^{-6} M/L$ ) and varying the quencher concentration ( $0.02M$  to  $0.10M$ ) in all the solvents. Fluorescence spectrophotometer F-2000 was used for fluorescence intensity measurements, with perpendicular geometry. Description of the experimental setup is discussed elsewhere[11]. Fluorescence measurements were made by taking fresh solution each time in a rectangular quartz cell having an airtight stopper. The solute has maximum absorption at 350 nm. The solute was excited at 350 nm and the fluorescence spectrum was recorded. The maximum fluorescence corresponds to 420 nm. First the fluorescence intensity  $I_0$  was measured without the quencher and then the fluorescence intensity  $I$  was measured at different quencher concentrations and at a fixed solute concentration. The experimental values are reproducible within 5% of the experimental error.

## RESULTS AND DISCUSSION

The S-V plots obtained by using the experimentally determined values of  $I_0$  and  $I$  are found to be nonlinear in all the solvents showing positive deviation and are shown in Fig. 1. Similar experimental results were also observed by others[1,6]. Thus, positive deviation from linearity suggests that quenching is not purely collisional and this may be attributed either to the ground state complex formation or to the sphere of action static quenching model[7].

The values of  $K_{sv}$  from equation (2) were found to be imaginary in all the solvents. Therefore the role of ground state complex formation is ruled out in the present case. This is also confirmed by the fact that there is no shift in the peak position in the emission spectrum of Bis-MSB in cyclohexane with different quencher concentrations and is shown in Fig. 2. These facts show that Eq. (2) is not applicable for the analysis of the data corresponding to the observed positive deviation in the S-V plots. Thus the analysis of the data for positive deviation in the S-V plot was made using "Sphere of action static quenching model". Fig. 3 shows the plots of  $[1 - (I/I_0)]/[Q]$  against  $I/I_0$  for Bis-MSB with  $CCl_4$  as quencher for different solvents which are linear according to equation (7).

The dynamic quenching constant  $K_{sv}$  is determined in all the cases by least square fit method using Eq.7 and the values




Fig. 1. Stern-Volmer plots of  $I_0/I$  against  $[Q]$  in different solvents.




Fig. 2. Emission spectra of Bis-MSB in cyclohexane with different quencher concentrations of  $\text{CCl}_4$

are given in the Table 1. Fluorescence lifetime  $\tau$  of the solute studied is obtained from the catalogue of the Koch-Light laboratories limited, England and is given at the bottom of the Table 1. Bimolecular quenching rate parameter  $k_q$  was determined from the experimentally determined values of  $K_{sv}$  and literature value of  $\tau$  according to the relation  $k_q = K_{sv}/\tau$  and the values are given in the Table 1. Here high values of  $k_q$  indicate efficient quenching of fluorescence.

In order to support static and dynamic effects, we have determined the magnitudes of static quenching constant  $V$  and radii  $r$  of sphere of action (or kinetic distance) using the above mentioned theory for sphere of action model. With the use of Eq. (5), (6) and (7) the values of  $V$  and  $r$  are determined by least square fit method in all the solvents and are given in Table 1. Similar results were also obtained by others [7, 12, 29].

The radii of the solute ( $R_Y$ ) and the quencher ( $R_Q$ ) molecules




Fig. 3. Plots of  $[1-(I_0/I)]/[Q]$  against  $I_0/I$  in different solvents.

Table 1. The values of Stern-Volmer constant  $K_{sv}$ , Bimolecular quenching rate parameter  $k_q$ , Static quenching constant  $V$  and kinetic distance  $r$  for different solvents.

Solvent	$K_{sv}$ ( $\text{M}^{-1}$ )	$k_q \times 10^{-10}$ ( $\text{M}^{-1}\text{s}^{-1}$ )	$V$ ( $\text{mol}^{-1}\text{dm}^3$ )	$r$ ( $\text{\AA}$ )
Hexane	3.26	0.251	4.62	12.23
Cyclohexane	7.55	0.581	5.97	13.33
Toluene	27.07	2.083	4.00	11.66
Benzene	40.59	3.122	5.28	12.79
Dioxane	73.91	5.685	9.94	15.79

Molecule: 1,4-bis [2-(2-methylphenyl) ethenyl]-benzene Quencher: Carbon Tetra Chloride ( $\text{CCl}_4$ )

$R_Y = 4.24 \text{ \AA}$ ,  $R_Q = 2.79 \text{ \AA}$ ,  $\tau = 1.3 \text{ ns}$

were determined by adding the atomic volumes of all the atoms constituting the molecule as suggested by Edward [23] and are given at the bottom of the Table 1. From these values of  $R_Y$  and  $R_Q$  the sum of the molecular radii  $[R]$  of the solute Bis-MSB and the quencher  $\text{CCl}_4$  is determined. This sum of the molecular radii  $R$  is referred to as encounter distance or contact distance or reactive distance. This value of  $R$  is then compared with the values of ' $r$ ' to verify whether the reaction is due to sphere of action model. From Table 1 we see that the values of kinetic distance ' $r$ ' are greater than the encounter distance  $R$ . Therefore according to Andre et al. and Zeng et al. [20, 24] the static effect takes place irrespective of ground state complex formation provided reactions are limited by diffusion indicating the sphere of action model holds well in our case also. Further it may also be noted that a positive

deviation in S-V plot is expected when both static and dynamic quenching occurs simultaneously[19].

In order to find whether the reactions are diffusion limited we have invoked the Finite Sink Approximation Model. To apply this model according to Eq.(15) we need to determine the values of  $K_{sv}^{-1}$  and  $[Q]^{1/3}$ . Where  $K_{sv} = [(I_0/I)-1]/[Q]$  and  $[Q]$  the quencher concentration from 0.02M to 0.10M. For efficient quenching processes (concentration dependent) the value  $K_{sv}$  is often observed to increase with  $[Q]$ . Hence the values of  $K_{sv}$  were determined at each quencher concentration in all the solvents and the values of  $K_{sv}^{-1}$  are also determined. Fig. 4 shows the plots of  $K_{sv}^{-1}$  against  $[Q]^{1/3}$ . From these figures we see that all the plots in different solvents are almost linear and small deviation may be due to experimental uncertainties. Hence, the linear dependence of  $K_{sv}^{-1}$  on the one-third power of quencher concentration within the error limits is confirmed[20]. Then the least square fit value of  $K_{sv}^0$




Fig. 4. Plots of  $K_{sv}^{-1}$  versus  $[Q]^{1/3}$  in different solvents.

(Stern-Volmer constant at  $[Q] = 0$ ) was obtained from the intercept of the plot of  $K_{sv}^{-1}$  against  $[Q]^{1/3}$  according to Eq. 15. Similarly, mutual diffusion coefficients  $D$  were determined from the slope of the Eq.15 by least square fit method and the values of  $K_{sv}^0$  and  $D$  are given in Table 2. Using these values of  $K_{sv}^0$  and  $D$ , the distance parameter  $R'$  was determined according to Eq.16 and the values are given in the Table 2. According to Joshi et al.[28] the bimolecular reactions are said to be diffusion limited if the values of  $k_q$  (determined from Eq. (7)) are greater than  $4\pi N'R'D$ . Hence the values of  $4\pi N'R'D$  are calculated using the experimentally determined values of  $R'$  and  $D$  of Eq. (15) and are given in the Table 2. We see from the Table that in three solvents the values of  $k_q$  are greater than  $4\pi N'R'D$ , which is an expected result for diffusion limited reaction[27].

It is important to note that, when an attempt was made to correlate the experimentally determined values of mutual diffusion coefficient  $D$  and distance parameter  $R'$  using finite sink approximation model with the corresponding values of  $D$  and  $R$  calculated from Stokes-Einstein relation and Edward's empirical relations respectively in all the systems, the values of  $D$  and  $R'$  (or  $R$ ) determined from the two methods are quite comparable only in some solvents. From the Table 3 it is clear that correlation fails in some other solvents. Similar discrepancies have also been observed by others[20, 28] in some bimolecular quenching reactions. This may be due to the uncertainties in

Table 3. The values of Mutual diffusion coefficients  $D^a$  and  $D^b$ , Distance parameter  $R'$  and encounter distance  $R$ .

Solvent	$D^a \times 10^5$ ( $\text{cm}^2\text{s}^{-1}$ )	$D^b \times 10^5$ ( $\text{cm}^2\text{s}^{-1}$ )	$R'$ ( $\text{\AA}$ )
Hexane	6.97	0.70	7.46
Cyclohexane	2.122	1.00	8.53
Toluene	3.762	3.42	6.77
Benzene	3.446	3.99	8.10
Dioxane	1.745	4.54	10.65

Molecule: 1,4-bis [2-(2-methylphenyl) ethenyl]-benzene

Quencher: Carbon Tetra Chloride ( $\text{CCl}_4$ )  $R(=R_Y + R_Q) = 7.03 \text{\AA}$

<sup>a</sup>Diffusion coefficients determined from Stoke's Einstein relation.

<sup>b</sup>Diffusion coefficients determined from Finite Sink Approximation Model.

Table 2. The values of  $K_{sv}^0$  (steady state quenching constant at  $[Q] = 0$ ), Mutual diffusion coefficient  $D$ , Distance parameter  $R'$ ,  $4\pi N'R'D$  and Quenching rate parameter  $k_q$ .

Solvent	$K_{sv}^0$ ( $\text{dm}^3\text{mol}^{-1}$ )	$D \times 10^5$ ( $\text{cm}^2\text{s}^{-1}$ )	$R'$ ( $\text{\AA}$ )	$4\pi N'R'D \times 10^{-10}$ ( $\text{M}^{-1}\text{s}^{-1}$ )	$k_q \times 10^{-10}$ ( $\text{M}^{-1}\text{s}^{-1}$ )
Hexane	5.141	0.70	7.46	0.39	0.251
Cyclohexane	8.402	1.00	8.53	0.64	0.581
Toluene	22.81	3.42	6.77	1.75	2.083
Benzene	31.80	3.99	8.10	2.44	3.122
Dioxane	47.62	4.54	10.65	3.66	5.685

Molecule: 1,4-bis [2-(2-methylphenyl) ethenyl]-benzene Quencher: Carbon Tetra Chloride ( $\text{CCl}_4$ )

$R(=R_Y + R_Q) = 7.03 \text{\AA}$

the values of adjustable parameter 'a' in the Stokes's-Einstein relation and the approximation in the values of atomic volume in the Edward's relation. Hence we may conclude that finite sink approximation model is valid in recovering the parameters D and R' (or R).

## CONCLUSION

From the above discussion we observe that: (1) the S-V plots show positive deviation leading to high values of  $k_q$ , indicating efficient fluorescence quenching. (2) Static quenching constant 'V' and kinetic distance 'r' are in agreement with the corresponding literature values. (3) The value of  $k_q$  is greater than  $4\pi N'R'D$  in three solvents. (4) Further the values of R' and D determined from the finite sink approximation model are close to the values determined from Edward's and Stokes-Einstein empirical relations respectively. In view of the above facts we may conclude that quenching reaction is diffusion limited and both static and dynamic quenching processes are partly playing a role in these systems.

## ACKNOWLEDGEMENT

The authors are grateful to the Chairman, Department of Physics, Karnatak University, Dharwad for providing the facilities to carry out experiments. Authors also thank Dr. S. Kumar B. Hadimani for providing the necessary scintillation samples used in the present work.

## REFERENCES

- Roy, R. and Mukherjee, S. (1987) Fluorescence quenching of carbazole and indole by ethylenetrithiocarbonate. *Chem. Phys. Lett.* **140**, 210-214.
- Moriya, T (1984) Excited-state reactions of Coumarins in aqueous solutions II The fluorescence quenching of 7-ethoxycoumarins by halide ions. *Bull. Chem. Soc. Jpn.* **57**, 1723-1729.
- Umarjee, D. M and Chikkur, G. C. (1987) Effect of dilution and temperature on the energy transfer and quenching in organic scintillators. *Phys. Lett. A* **123**, 23-26.
- Hanagodimath, S. M., Gadaginmath, G. S. and Chikkur, G. C. (1990) Role of Brownian diffusion and interreaction distance on energy transfer and quenching in an organic liquid scintillators. *Appl. Radiat. Isot.* **41**, 817-821.
- Arathi Rani, S., Sobhanadri, J. and Prasad Rao, T. A. (1996) Solvent and concentration effects on the steady state fluorescence of fluorenone. *J. Photochem. Photobiol.* **33**, 749-752.
- Kadadevaramath, J. S., Giraddi, T. P. and Chikkur, G. C. (1998) Fluorescence quenching of BBOT by aniline in different organic solvents. *J. Photoscience*, **4** (3), 105-112.
- Giraddi T. P., Kadadevaramath, J. S. Malimath, G. H. and Chikkur, G. C. (1996) Effect of solvent on the fluorescence quenching of organic liquid scintillators by aniline and carbon tetrachloride, *Appl. Radiat. Isot.*, (U.K) **47**[4], 461-466.
- Hariharan. C, Vijayasree, V. and Mishra, A. K. (1997) Quenching of 2,5-diphenyl oxazole (PPO) by metal ions, *Radiation Measurements*, **75** [3] 205-211.
- Hariharan. C and Mishra, A. K. (2000) Quenching of liquid scintillators by chloroalkanes and chloroalkenes, *Radiation Measurements*, **32**[2] 113- 121.
- Suresh Kumar. H. M., Ravikanth, M. N., Kunabenchi, Hanagodimath, S. M., Biradar, J. S., Renuka, P., Math, N. N., Kadadevaramath, J. S. and Inamdar, S. R. (2002) Fluorescence quenching of 6-Chloro-2-Phenyl [benz(b) Pymole]-1-carbazaldehyde thiosimicarbozone by  $CCl_4$  in different solvents, *Ind. J. Pure and Appl. Phys.*, **40**, 466-470.
- Shailaja, M. K., Hanagodimath, S. M., Kadadevaramath, J. S. and Chikkur, G. C. (2000) Effect of solvent polarity on the fluorescence quenching of liquid scintillators by aniline, *J. Photoscience*, **6**[4], 159-163.
- Hanagodimath. S. M., Chikkur, G. C. and Gadaginmath, G. S. (1990) Environmental effects on the energy migration coefficient, the effective energy transfer and quenching distance in an organic liquid scintillators. *Chem. Phys.* **148** [2,3] 347-358.
- Kadadevaramath, J. S., Giraddi, T. P. and Chikkur, G. C. (1996) Electronic excitation energy quenching of an organic liquid scintillator by carbon tetra chloride in different solvents, *Radiation Measurements (USA)* **26**[1], 117-121.
- Giraddi T. P., Kadadevaramath, J. S. Malimath, G. H. and Chikkur, G. C. (1996) Quenching of 2-phenylindole by carbon tetrachloride and aniline in different solvents. *Ind. J. Pure & Appl. Phys.*, **34**, 244-248,
- Hanagodimath. S. M., Chikkur, G. C. and Gadaginmath, G. S. (1992) On the mechanism of excitation energy transfer involving long range and short range interaction in dilute organic liquid scintillators system. *Acta Physical Polonica A.*, **81**[3], 361-368.
- Hanagodimath. S. M., Chikkur, G. C. and Gadaginmath, G. S. (1991) An unique method of determining the excitation energy migration coefficient in an organic liquid scintillator. *Pramana J of Physics (India)* **37** [2] 153-161.
- Lissi, E. A., Encians, M. V., Birtolotti, S. G., Cosa J. J., and Previtali C. M. (1990) Fluorescence quenching of indolic compounds in reverse micells of AOT. *Photochem. Photobiol.*, **51**, 53-58.
- Nigam, S., Belletete, M., Sarpal R. S., and Durocher G. (1995) Solubilization of two cyano 3H-indole molecular probes in CDS and CTAB micelles; spectral and photophysical characterization. *J. Chem. Soc. Faraday Trans.*, **91**, 2133-2139.
- Behera, P. K. and Mishra A. K. (1993) Static and dynamic model for 1-naphthol fluorescence quenching by carbon tetrachloride in dioxine-acetonitrile mixtures. *J. Photochem. Photobiol.A: Chem.*, **71**, 115-118.
- Zeng, H. and Durocher, G. (1995) Analysis of fluorescence quenching in some antioxidants from nonlinear Stern-Volmer plots. *J. Lumin.*, **63**, 75-84
- Frank, J. M. and Wawilow, S. J. (1931) Sphere of action of the extinction phenomenon in fluorescent liquids. *Z. Phys.*,

- 69, 100-110.
22. Moon, A. Y., Poland, D. C., and Scheraga, H. A. (1965) Thermodynamic data from fluorescence spectra I. The system phenol-acetate. *J. Phys. Chem.*, **69**, 2960-2966.
23. Edward, J. T. (1956) Molecular Volumes and Parachor. *Chem.Ind., (London)* 774.
24. Andre, J. C., Niclause M., and Ware W. R. (1978) Kinetics of partly diffusion controlled reactions. I. Transient and apparent transient effect in fluorescence quenching. *Chem. Phys.* **28**, 371-376.
25. Keizer, J (1982) Nonequilibrium statistical thermodynamics and the effect of diffusion on chemical reaction rates. *J. Phys. Chem.*, **86**, 5052-5067.
26. Keizer, J (1983) Nonlinear fluorescence quenching and the origin of the positive curvature in Stern-Volmer plots. *J. Am. Chem.Soc.*, **105**, 1494-1499; Keizer, J. (1985) Additions and corrections. *J. Am. Chem. Soc.*, **107**, 5319.
27. Keizer, J (1987) Diffusion effects on rapid bimolecular chemical reactions. *Chem. Rev.*, **87**, 167-180.
28. Joshi, G. C., R. Bhatnagar, S. Doraiswamy and N. Periasamy (1990) Diffusion controlled reactions: Transient effects in the fluorescence quenching of indole and N-acetyltryptophanamide in water. *J. Phys. Chem.*, **94**, 2908-2914.
29. Birks, J. B., Salet, M. and Leite, S. C. P. (1970) Energy transfer in organic systems 5. Quenching of naphthalene fluorescence by biacetyl. *Proc. Phys Soc., (London)* **3**, 417.