

Identification of Three Fungi Newly Intercepted from Importing Plants in Korea

Ik-Hwa Hyun*, Noh-Yeoul Heo, Seo-Yeon Chang¹, Jong-Young Heo² and Vadim Mel'nik³

Central Post-Entry Quarantine Station, National Plant Quarantine Service, Suwon 443-400, Korea

¹Honam Regional Office, National Plant Quarantine Service, Gunsan 573-879, Korea

²Pest Survey & Control Division, National Plant Quarantine Service, Anyang 430-016, Korea

³Komarov Botanical Institute, Russian Academy of Sciences, Saint Petersburg 197376, Russia

(Received September 7, 2005)

Three fungi newly intercepted from importing plants were identified in 2004. They were *Ascochyta chrysanthemi* on *Lactuca sativa* from China, *A. spinaciicola* on *Spinacia oleracea* from Denmark, and *Leptosphaerulina australis* on *Brassica oleracea* var. *capitata* from China. The characters of these fungi were described and illustrated.

KEYWORDS: *Ascochyta*, Identification, *Leptosphaerulina*, Plant quarantine

In making plant quarantine decisions it is essential that the fungi present in subject plant materials should be identified accurately, not only to exclude non-indigenous, pathogenic fungi, but also to allow the entry of material infected by fungi that are already present in a country and therefore not of plant quarantine concern (Palm, 1999).

Importing plant and plant products to Korea are inspected at the port by National Plant Quarantine Service (NPQS) inspectors. When a fungus is found during the inspection, the inspectors responsible for identifying fungi make a determination. If the fungus cannot be identified at the port, the specimen is forwarded to the consulting team of NPQS for pest identification. In 2004, the authors identified three fungal specimens newly intercepted from importing plants as the consulting team members of NPQS.

These specimens were made from *Lactuca sativa* (lettuce) sample importing from China, *Spinacia oleracea* (spinach) from Denmark, and *Brassica oleracea* var. *capitata* (Chinese cabbage) from China, respectively (Table 1). Leaf samples were incubated in a moist chamber for three to five days. Seed sample was incubated for seven days by the blotter method (Neergaard, 1977). After sporulation, morphological characters were observed for the identification.

On the basis of the morphology, three fungi intercepted from lettuce, spinach and Chinese cabbage were identified as *Ascochyta chrysanthemi* F. Stevens, *A. spinaciicola* Melnik, and *Leptosphaerulina australis* McAlp., respectively. Identification was done by referring to Mel'nik (2000) and Punithalingam (1980) for the *Ascochyta* species, and to Graham and Luttrell (1961) for *Leptosphaerulina australis*. These three fungi were characterized

as follows:

***Ascochyta chrysanthemi* F. Stevens.** Conidiomata pycnidial, epiphyllous, scattered on lesions of leaves, immersed, punctiform with papillate ostiole. Conidia cylindrical or oblong-ellipsoidal, straight or sometimes slightly curved, both ends rounded, sometimes narrowed to one end, not or only slightly constricted at the septa, hyaline, $10\text{--}22.5 \times 3\text{--}6.3 \mu\text{m}$ (Fig. 1A).

This fungus was detected from the leaves of lettuce showing spot symptom. Punithalingam (1980) recorded lettuce as a host by inoculation. However, principal hosts of this fungus are *Chrysanthemum cinerariifolium* and *C. morifolium* (Punithalingam, 1980). A chrysanthemum is one of the most important plants in floriculture in Korea. Once introduced and established in an area, this fungus is both difficult and costly to eradicate, and control of the disease remains as additional production expense. This fungus was recorded in Africa, Australasia & Oceania, Europe, North America and Asia (Punithalingam, 1980), however this fungus is not reported in Korea according to the Korean Society of Plant Pathology (2004). Its teleomorph is *Didymella ligulicola* (syn. *Mycosphaerella ligulicola*), designated as a plant quarantine pathogen in Korea.


***Ascochyta spinaciicola* Melnik.** Conidiomata pycnidial, scattered, immersed, $110\text{--}170 \mu\text{m}$ in diam. Conidia cylindrical or oblong-ellipsoidal, both ends rounded, straight or sometimes slightly curved, not constricted or rarely slightly constricted at the septa, hyaline, $10\text{--}17.5 \times 3.8\text{--}5 \mu\text{m}$ (Fig. 1B).

Mel'nik (2000) recorded Asia (George) as a distribution area of the fungus. Interestingly this fungus was intercepted from the spinach from Denmark. Mel'nik (2000) observed this fungus on leaves of spinach. In this work, it

*Corresponding author <E-mail: ihhyun@npqs.go.kr>

Table 1. List of plant specimens and fungi identified in this study

Accession No.	Host	Plant part	Symptom	Country importing	Fungi identified
PQ041	<i>Lactuca sativa</i> L.	Leaf	Spot	China	<i>Ascochyta chrysanthemi</i>
PQ042	<i>Spinacia oleracea</i> L.	Seed	No symptom	Denmark	<i>Ascochyta spinaciicola</i>
PQ043	<i>Brassica oleracea</i> var. <i>capitata</i> L.	Leaf	Spot	China	<i>Leptosphaerulina australis</i>

**Fig. 1.** Conidia of *Ascochyta chrysanthemi* (A) and *A. spinaciicola* (B). Bar = 15 μ m.


was detected from seeds of spinach. So far, information on this fungus is not known well.

***Leptosphaerulina australis* McAlp.** Ascomata pseudothecial, uniloculate, immersed. Asci bitunicate, thick-walled, 8-spored, 75–95 \times 45–58 μ m. Ascospores ellipsoid to oblong with gelatinous sheath, 3–5 transverse septa, 1–3 longitudinal septa, hyaline to brown, 26–38 \times 9–13 μ m (Fig. 2).

According to Graham and Luttrell (1961), *Leptosphaerulina trifolii*, *L. briosiana* and *L. arachidicola* were pathogenic, but *L. australis*, *L. argentinensis* and *L. americana* were saprobic. These six species of *Leptosphaerulina* are recognized on forage legumes in the U.S.A. However, Irwin and Davis (1985) reported that *L. argentinensis* was pathogenic to *Stylosanthes guianensis*. In the present work, the fungus was detected from the spot lesions of Chinese cabbages. Pathogenicity of the fungus is also needed to study.

On the other hand, Booth and Pirozynski (1967), and Irwin and Davis (1985) regarded *L. australis* as a taxonomic synonym of *L. trifolii*. *L. trifolii* is plurivorous but especially on economic plants from the following families: Cruciferae, Euphorbiaceae, Gramineae, Leguminosae and Solanaceae. It causes leaf spot or pepper spot on account of leaves and petioles of diverse plants, particularly forage legumes (Booth and Pirozynski, 1967).

All three fungi intercepted are not reported in Korea according to the Korean Society of Plant Pathology (2004), and may have risks to domestic plants. It is necessary to inspect and identify thoroughly the presence of

**Fig. 2.** Ascospores and ascus of *Leptosphaerulina australis*. Bar = 30 μ m.

these fungi during plant quarantine procedures.

References

- Baker, K. F., Dimock, A. W. and Davis, L. H. 1961. Cause and prevention of the rapid spread of the *Ascochyta* disease of chrysanthemum. *Phytopathology* **51**: 101.
- Booth, C. and Pirozynski, K. A. 1967. *Leptosphaerulina trifolii*. Commonw. Mycol. Inst. Descriptions of Pathogenic Fungi and Bacteria No. 146.
- Farr, D. F., Bills, G. F., Chamuris, G. P. and Rossman, A. Y. 1989. Fungi on Plants and Plant Products in the United States. APS Press, St. Paul. 1255pp.
- Graham, J. H. and Luttrell, E. S. 1961. Species of *Leptosphaerulina* on forage plants. *Phytopathology* **51**: 680-693.
- Irwin, J. A. G. and Davis, R. D. 1985. Taxonomy of some *Leptosphaerulina* spp. on legumes in eastern Australia. *Aust. J. Bot.* **33**: 233-237.
- Mel'nik, V. A. 2000. Key to the Fungi of the Genus *Ascochyta* Lib. (Coelomycetes). BBA, Berlin. 192pp.
- Neergaard, P. 1977. Seed Pathology. The Macmillan Press Ltd., London and Basingstoke. 839pp.
- Palm, M. E. 1999. Mycology and world trade: a view from the front line. *Mycologia* **91**: 1-12.
- Punithalingam, E. 1980. *Didymella chrysanthemi*. Commonw. Mycol. Inst. Descriptions of Pathogenic Fungi and Bacteria No. 662.
- The Korean Society of Plant Pathology. 2004. List of Plant Diseases in Korea. 779pp.