

# Communications

## An Efficient Oxidation of 1,4-Dihydropyridines to Pyridines Using Silver Carbonate on Silica Gel and Celite

Ahmad Reza Momeni,<sup>\*</sup> Tayebeh Sameh, Hosein Golmohammadi, Hamid Javaherian Naghash, Hamid Aliyan, Ahmad Reza Massah, and Shirin Solati

Department of Chemistry, Islamic Azad University Shahreza Branch, 86145-311, Isfahan, Iran. \*E-mail: momeni@iaush.ac.ir  
Received September 9, 2005

**Key Words :** 1,4-Dihydropyridines, Silver carbonate, Oxidation, Celite, Silica gel


The oxidation of Hantzsch 1,4-dihydropyridines to the corresponding pyridines has been extensively studied in view of the pertinence of the reaction to the metabolism of Hantzsch esters and the calcium channel blocking drugs used in the treatment of various cardiovascular disorders.<sup>1</sup> The reaction has also been used to study the biologically significant NADH redox processes.<sup>2</sup> Consequently, this aromatization reaction continues to attract the attention of researchers for the discovery of milder and general protocols applicable to a wide range of 1,4-dihydropyridines. Several oxidizing agents and methods such as  $\text{KMnO}_4$ ,<sup>3</sup> solid supported pyridinium chlorochromate (PCC),<sup>4</sup> silica gel supported ferric nitrate,<sup>5</sup> nitric oxide,<sup>6</sup>  $\text{MnO}_2$ ,<sup>7</sup> ceric ammonium nitrate (CAN),<sup>8</sup> *tert*-butylhydroperoxide,<sup>9</sup> clay supported cupric nitrate,<sup>10</sup>  $\text{Bi}(\text{NO}_3)_3$ ,<sup>11</sup>  $\text{I}_2/\text{MeOH}$ ,<sup>12</sup>  $\text{Zr}(\text{NO}_3)_4$ ,<sup>13</sup> tetrakispyridine cobalt(II) dichromate (TPCD),<sup>14</sup> iodobenzene diacetate,<sup>15</sup>  $\text{NaNO}_2/\text{NaHSO}_4/\text{SiO}_2$ ,<sup>16</sup> 3-Carboxypyridinium chlorochromate (CPCC),<sup>17</sup>  $\text{BaMnO}_4$ ,<sup>18</sup>  $\text{Ti}(\text{NO}_3)_3 \cdot 3\text{H}_2\text{O}$ ,<sup>19</sup> or photochemical reactions<sup>20</sup> have been introduced in the literature for this purpose. The introduction of supported reagents for bringing about various chemical transformations has provided an attractive option for organic synthesis. These reagents not only modify the activity but also may impart product selectivity.

Because of our interest in oxidation processes, we decided to develop a practical and general approach for oxidative conversion of 1,4-dihydropyridines using a mild oxidant, silver carbonate. This reagent is a mild oxidizing agent. This operates under neutral and heterogeneous conditions. Silver carbonate on celite was employed for organic reactions such

as oxidation of alcohols,<sup>21</sup> 1,2-diols,<sup>22</sup> phenols,<sup>23</sup> amines and hydrazines.<sup>24</sup> Silver carbonate on silica gel and celite can be readily available. We wish to indicate that it can also serve as an excellent oxidant for a variety of Hantzsch 1,4-dihydropyridines.

A series of 1,4-DHP derivatives (**1-12**, Scheme 1) were synthesized<sup>20,25</sup> to investigate their conversion to the corresponding pyridines. Initially 4-phenyl-dihydropyridine (entry 3) has been used as a substrate to test the feasibility of silver carbonate used as an oxidant for DHP's. Investigation on the oxidation efficiency of silver carbonate alone at ambient temperature as well as higher temperature indicated that, oxidation of (**3**) into aromatized product did not proceed effectively. A better activity of silver carbonate was obtained when combined with silica gel or celite. Silver carbonate on celite as mentioned above is known, but silver carbonate on silica gel was used for the first time in this study. Subsequent experiments revealed that at reflux conditions in acetonitrile a ratio of DHP :  $\text{Ag}_2\text{CO}_3 = 1 : 6$  on silica gel and DHP :  $\text{Ag}_2\text{CO}_3 = 1 : 12$  on celite was most effective. Under similar conditions various substituted 1,4-dihydropyridines were efficiently converted to the pyridine derivatives<sup>26</sup> and the results are given in Table 1.

According to the experiment we observed that oxidation of 1,4-dihydropyridines with secondary alkyl group (entry 5) and benzyl groups (entry 6) at the 4-position were performed efficiently and only dealkylated pyridine derivative (**13**) was obtained. This is a general trend in the oxidation of 1,4-dihydropyridines. 1,4-Dihydropyridine (entry 4) with propyl group at the 4-position yielded two products, dealkylated



Scheme 1

**Table 1.** Oxidation of 1,4-Dihydropyridines with Silver Carbonate on Silica gel and Silver Carbonate on Celite

Entry	R	Products <sup>a</sup>	Silver carbonate/SiO <sub>2</sub>		Silver carbonate/Celite	
			Time (h)	Yields (%) <sup>b</sup>	Time (h)	Yields (%) <sup>b</sup>
1	H	<b>13</b>	0.9	99	0.5	95
2	CH <sub>3</sub>	<b>14</b>	3.5	79	5.5	98
3	C <sub>6</sub> H <sub>5</sub>	<b>15</b>	1.75	94	3.5	90
4	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	<b>13+16</b>	2	76–15	3	55–20
5	(CH <sub>3</sub> ) <sub>2</sub> CH	<b>13</b>	0.8	95	5	98
6	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH	<b>13</b>	0.75	94	2.2	95
7	3-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	<b>17</b>	5	87	9	90
8	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	<b>18</b>	5	82	6.5	80
9	2-Pyridyl	<b>19</b>	0.8	93	8	85
10	4-ClC <sub>6</sub> H <sub>4</sub>	<b>20</b>	3	96	4	95
11	4-BrC <sub>6</sub> H <sub>4</sub>	<b>21</b>	3	91	4	80
12	4-CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	<b>22</b>	1.5	98	5	98

<sup>a</sup>All products were characterized by <sup>1</sup>H-NMR, IR and their m.p.s. Compared with literature values. <sup>b</sup>Isolated yields

compound (**13**), which is rarely observed by primary alkyl groups, and pyridine derivative (**16**) with retention of 4-substituent. However, 4-methyl, 4-aryl and 4-heteroaryl-1,4-dihydropyridines (entries 2, 3 and 7-12) were oxidized with retention of substitution at 4-position to afford the corresponding pyridines (**14-22**) in good to excellent yields. In comparison with silver carbonate on celite, silver carbonate on silica gel is more reactive.

The effect of different solvents on the oxidation reaction was investigated. In comparison with acetonitrile, the reaction time was longer and the yields of products were considerably lower when dichloromethane, carbontetrachloride and chloroform were employed as a solvent.

In summary, we have found that silver carbonate on silica gel or celite is a valuable addition to the existing methods available for the oxidation of 1,4-dihydropyridines with additional advantages of ease of procedure, workup and high yields.

## References

- (a) Janis, R. A.; Triggler, D. J. *J. Med. Chem.* **1983**, *26*, 775. (b) Gordeev, M. F.; Patel, D. V.; Gordon, E. M. *J. Org. Chem.* **1996**, *61*, 924. (c) Bossert, F.; Vater, W. *Med. Res. Rev.* **1989**, *9*, 291.
- Stout, D. M.; Meyers, A. I. *Chem. Rev.* **1982**, *82*, 223.
- Vanden Eynde, J. J.; D'Orazio, R.; Van Haverbeke, Y. *Tetrahedron* **1994**, *50*, 2479.
- Vanden Eynde, J. J.; Mayence, A.; Maquestiau, A. *Tetrahedron* **1992**, *48*, 463.
- Khadikar, B.; Brokar, S. *Synth. Commun.* **1998**, *28*, 207.
- Itoh, T.; Nagata, K.; Matsuya, Y.; Miyazaki, M.; Ohsawa, A. *J. Org. Chem.* **1997**, *62*, 3582.
- Vanden Eynde, J. J.; Delfosse, F.; Mayence, A.; Van Haverbeke, Y. *Tetrahedron* **1995**, *51*, 6511.
- Pfister, J. R. *Synthesis* **1990**, 689.
- Chavan, S. P.; Dantal, S. W.; Kalkote, U. R.; Jyothirmal, V. S.; Kharul, R. K. *Synth. Commun.* **1998**, *28*, 2789.
- Maquestiau, A.; Mayence, A.; Vanden Eynde, J. J. *Tetrahedron Lett.* **1991**, *32*, 3839.
- Mashraqui, S. H.; Kamik, M. A. *Synthesis* **1998**, 713.
- Yadav, J. S.; Subba Reddy, B. V.; Sabita, G.; Kiran Kumar Reddy, G. S. *Synthesis* **2000**, 1532.
- Sabita, G.; Kiran Kumar Reddy, G. S.; Srinivas, Ch.; Fatima, N.; Yadav, J. S. *Synthesis* **2003**, 1267.
- Wang, B.; Hu, Y.; Hu, H. *Synth. Commun.* **1999**, *29*, 4193.
- Cheng, D. P.; Chen, Z. C. *Synth. Commun.* **2002**, *32*, 793.
- Zolfigol, M. A.; Kiany-Borazjani, M.; Sadeghi, M. M.; Memarian, H. R.; Mohammadpoor-Baltork, I. *Synth. Commun.* **2000**, *30*, 2945.
- Mohammadpoor-Baltork, I.; Sadeghi, M. M.; Memarian, H. R.; Pairov, R. *J. Chem. Research (s)* **2000**, 40.
- Memarian, H. R.; Sadeghi, M. M.; Momeni, A. R. *Synth. Commun.* **2001**, *31*, 2241.
- Momeni, A. R.; Massah, A. R.; Javaherian Naghash, H.; Aliyan, H.; Solati, S.; Sameh, T. *J. Chem. Research* **2005**, 227.
- (a) Memarian, H. R.; Sadeghi, M. M.; Momeni, A. R. *Indian J. Chem.* **1999**, *38B*, 800. (b) Memarian, H. R.; Sadeghi, M. M.; Momeni, A. R. *Monatsheshefte für Chemie.* **2002**, *133*, 661. (c) Memarian, H. R.; Sadeghi, M. M.; Momeni, A. R. *Indian J. Chem.* **2001**, *40B*, 508.
- (a) Schwab, J. M.; Chornig-Kei, H. *J. Chem. Soc. Chem. Commun.* **1989**, 872. (b) Gardner, D.; Glen, A. J.; Turner, W. B.; Dimitriadis, E.; Massy-Westropp, R. A. *Aust. J. Chem.* **1980**, *33*, 2729.
- (a) Corey, E. J.; Ueda, Y.; Rudew, R. A. *Tetrahedron Lett.* **1975**, 4347. (b) Daily, O. D. Jr.; Fuchs, P. L. *J. Org. Chem.* **1980**, *45*, 216.
- Balogh, V.; Fetizon, M.; Golfer, M. *J. Org. Chem.* **1971**, *36*, 1339.
- Hedayatallah, M.; Dechatre, J. P.; Denivelle, J. P. *Tetrahedron Lett.* **1975**, 2039.
- Love, B.; Goodman, M. M.; Snader, K. M.; Tedeshi, R.; Macko, E. *J. Med. Chem.* **1974**, *19*, 956.
- Silver carbonate is commercially available and prepared according to the following procedure.  
*Preparation of solid-supported silver carbonate:* Silver nitrate (30 g) is dissolved in 200 mL of distilled water, and silica gel (30 g) is added. To the stirred mixture, a solution of sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>, 10 H<sub>2</sub>O) (30 g) in distilled water (300 mL) is slowly added. The yellow-green precipitate is filtered and washed to neutrality with distilled water. It is then dried (4 h) by rotatory evaporation on a steam bath, preferably in the dark. When prepared according to this procedure, 0.6 g of this reagent contains approximately 1 mmol of Ag<sub>2</sub>CO<sub>3</sub>.  
*Oxidation of 1,4-dihydropyridines with silver carbonate on silica gel: general procedure:* A suspension of 1,4-dihydropyridines (0.2 mmol) and silica gel supported silver carbonate (0.72 g) in acetonitrile (10 mL) was stirred at reflux condition for the time indicated in Table 1. The progress of the reaction was monitored by TLC. The mixture was filtered and then washed with acetonitrile. The filtrate was evaporated, the product obtained was recrystallized with ethanol. Products of entry 4 were purified by TLC.  
*Oxidation of 1,4-dihydropyridines with silver carbonate on celite:* Same procedure as above method but with 1.44 g celite supported silver carbonate.