

# Lysophosphatidylcholine Attenuates Endothelium-dependent Relaxation Responses through Inhibition of ACh-induced Endothelial $[Ca^{2+}]_i$ Increase

Seong-Chun Kwon<sup>1</sup>, Yong-Ho Lee<sup>2</sup>, Taicksang Nam<sup>2</sup>, and Duck-Sun Ahn<sup>2</sup>

<sup>1</sup>Department of Physiology, College of Medicine, Kwandong University, Gangneung 210–701, <sup>2</sup>Department of Physiology, Yonsei University College of Medicine, Seoul 120–752, Korea

Lysophosphatidylcholine (LPC), which accumulates in atherosclerotic arteries, has been reported to inhibit endothelium-dependent relaxation (EDR) in many different species. However, the underlying mechanism of LPC-induced inhibition of EDR is still uncertain. In the present study, we measured simultaneously both isometric tension and cytosolic free  $Ca^{2+}$  ( $[Ca^{2+}]_i$ ) in rabbit carotid strips, and examined the effect of LPC on tension and  $[Ca^{2+}]_i$ . In carotid strips with intact-endothelium, high  $K^+$  (70 mM) increased both tension and  $[Ca^{2+}]_i$ , and cumulative addition of acetylcholine (ACh) from 0.1 to 10  $\mu$ M induced dose dependent increase of  $[Ca^{2+}]_i$  with concomitant relaxation. In the presence of L-NAME (0.1 mM), ACh increased  $[Ca^{2+}]_i$  without affecting the amplitude of high  $K^+$ -induced tension. These ACh-induced change of  $[Ca^{2+}]_i$  and tension was abolished by removal of endothelium or 10 nM 4-DAMP (muscarinic receptor antagonist) pretreatment. Pretreatment of LPC (10  $\mu$ M) inhibited ACh (10  $\mu$ M)-induced change of tension and  $[Ca^{2+}]_i$  in endothelium-intact carotid artery. On the other hand, LPC had no effect on ACh-induced change of tension and  $[Ca^{2+}]_i$  in endothelium denuded artery. In  $Ca^{2+}$ -free external solution, ACh transiently increased  $[Ca^{2+}]_i$ , and pretreatment of LPC significantly inhibited ACh-induced transient  $[Ca^{2+}]_i$  change. Based on the above results, it may be concluded that LPC inhibits the ACh-induced  $[Ca^{2+}]_i$  change through inhibition of  $Ca^{2+}$  mobilization in vascular endothelial cells, resulting in decreased production of NO and concomitant inhibition of endothelium-dependent vascular relaxation.

**Key Words:** Vascular endothelium, LPC, EDR, Endothelial  $[Ca^{2+}]_i$

## INTRODUCTION

Impairment of endothelium-dependent relaxation (EDR) is thought to play an important role in the development of abnormal vascular responses in atherosclerotic arteries (Flavahan, 1992), and oxidized low-density lipoprotein (LDL), which is accumulated in the vessel wall in atherosclerosis, plays an important role in impairment of EDR (Berliner et al, 1995). The inhibitory effect of oxidized LDL, which is not shared by native LDL, is mainly mediated by lysophosphatidylcholine (LPC) (Murohara et al, 1994). When LDL is oxidized, the production of LPC from the phosphatidylcholine is increased 3 to 5 folds (Parthasarathy et al, 1985), and exogenous application of LPC causes an endothelial dysfunction similar to that observed in atherosclerotic arteries (Miwa et al, 1994; Murohara et al, 1994; Huang et al, 2002). Although LPC-induced inhibition of EDR has been examined in many studies, details of its mechanism are not clear at present.

Vascular endothelium secretes a variety of bioactive

agents including nitric oxide (NO), prostacyclin, and endothelium-dependent hyperpolarizing factors. These bioactive agents play an important role in controlling vascular tone (Gryglewski et al, 1988). A rise in endothelial cytosolic  $Ca^{2+}$  concentration ( $[Ca^{2+}]_i$ ) in response to various chemical or physical stimuli is an essential step in the synthesis or release of these bioactive agents (Boulaner & Vanhoutte, 1997). There is a positive correlation between increase in endothelial  $[Ca^{2+}]_i$  and release of NO in rat aortic endothelium, suggesting that  $Ca^{2+}$  is prerequisite for activation of nitric oxide synthase (Busse & Mulsch, 1990). In endothelial cells, the agonist-induced elevation of  $[Ca^{2+}]_i$  was due to the release of  $Ca^{2+}$  from internal stores by 1,4,5-triphosphate ( $IP_3$ ) or the influx of  $Ca^{2+}$  from external space (Adams et al, 1989). In fact, LPC has been reported to inhibit the agonist-induced increase in  $[Ca^{2+}]_i$  in cultured endothelial cells (Inoue et al, 1992) or in endothelium-intact tissue preparations (Miwa et al, 1994; Huang et al, 2002). Thus, LPC-induced inhibition of EDR may be due to inhibition of  $IP_3$ -induced  $Ca^{2+}$  release or  $Ca^{2+}$  influx from external space. However, which one is responsible for the

Corresponding to: Duck-Sun Ahn, Department of Physiology, Yonsei University College of Medicine, 134 Sinchon-dong, Seodaemun-gu, Seoul 120-752, Korea. (Tel) 82-2-2228-1707, (Fax) 82-2-393-0203, (E-mail) dsahn@yumc.yonsei.ac.kr

**ABBREVIATIONS:** ACh, addition of acetylcholine; EDR, endothelium-dependent relaxation; LDL, low-density lipoprotein; LPC, lysophosphatidylcholine; NO, nitric oxide;  $[Ca^{2+}]_i$ , cytosolic  $Ca^{2+}$  concentration.

inhibition of endothelial  $[Ca^{2+}]_i$  in intact vascular strips is still uncertain.

In the present study, we investigated the mechanism of LPC-induced inhibition of EDR in the rabbit carotid arteries. For this, we used a method that allows simultaneous measurements of endothelial  $[Ca^{2+}]_i$  and vascular tone, and then measured effect of LPC on ACh-induced endothelial  $[Ca^{2+}]_i$  and tension response in rabbit carotid arteries. Our results suggest that LPC suppressed ACh-induced endothelial  $[Ca^{2+}]_i$  elevation via inhibition of  $Ca^{2+}$  release from internal stores in rabbit carotid strips.

## METHODS

### *Tissue preparations*

New Zealand male white rabbits (1.5–2 kg) were anesthetized with pentobarbital sodium (30 mg/kg body weight IV), and the right and left carotid arteries were isolated and cleaned of surrounding tissues. Carotid arteries were cut into helical strips of 1.5- to 2-mm width and 8 to 10-mm length and placed in physiological salt solution (PSS) of the following composition (in mM): NaCl 136.9, KCl 5.4,  $CaCl_2$  1.5,  $MgCl_2$  1.2,  $NaHCO_3$  23.8, EDTA 0.01, glucose 5.5. Ethylenediaminetetraacetic acid (EDTA) at 0.01 mM was also added to chelate the contaminated heavy metal ions. High  $K^+$  solution was prepared by replacing NaCl with equimolar KCl. These solutions were saturated with 95%  $O_2$  and 5%  $CO_2$  at 37°C to maintain pH at 7.4.

### *Simultaneous measurements of isometric tension and $[Ca^{2+}]_i$ in isolated carotid strips*

$[Ca^{2+}]_i$  was measured according to the method described by Sato et al. (1990) using the fluorescent  $Ca^{2+}$  indicator fura-2. Muscle strips were exposed to the acethoxymethyl ester of fura-2 (fura-2/AM, 10  $\mu$ M) in the presence of 0.02% cremophor EL for 5–6 hr at room temperature (22–24°C). The muscle strip loaded with fura-2 was held horizontally in a temperature-controlled organ bath filled with normal PSS, and the endothelial side of strip was faced on the aperture of UV light. The carotid strip was illuminated alternately at a cycle of 48 Hz with two excitation wavelengths of 340 nm and 380 nm, and the fluorescence emitted from strip was collected into a photomultiplier tube through a 500 nm filter. The amount of emitted fluorescence induced by 340 nm excitation wave ( $F_{340}$ ) and that induced by 380 nm excitation wave ( $F_{380}$ ) was measured successively, and the ratio of these two fluorescence values ( $F_{340}/F_{380}$ ) was calculated. The ratio was used as an indicator of intracellular  $Ca^{2+}$  concentration ( $[Ca^{2+}]_i$ ). To measure the  $[Ca^{2+}]_i$  change in endothelial cells, endothelial surface of the strip was faced to the bottom of the chamber, and emitted fluorescence was detected in the presence or absence of endothelium. Experiments were performed with a bio-fluorimeter (CAF-110, Japan Spectroscopic, Tokyo, Japan). The resting  $[Ca^{2+}]_i$  in normal PSS and the sustained  $[Ca^{2+}]_i$  in the presence of 70 mM KCl were taken as 0% and 100%, respectively.

### *Statistics*

The results of the experiments are expressed as mean  $\pm$  S.E.M. Paired Student's *t*-test was used for statistical

analysis of the results, and the number of preparations taken from separate animals was indicated by *n*. *P* values of less than 0.05 were considered significant.


## RESULTS

### *Effect of ACh on $[Ca^{2+}]_i$ and tension in high $K^+$ -stimulated carotid strips*


Typical recordings of the response to ACh in carotid arterial strips with endothelium are shown in Fig. 1. Application of high  $K^+$  (70 mM  $K^+$ ) evoked the contraction with simultaneous increase of  $[Ca^{2+}]_i$  in endothelium-intact carotid strips. ACh, administered after high  $K^+$ -stimulated responses, reached to steady state, and elicited a concentration-dependent additional increase in  $[Ca^{2+}]_i$ , which was accompanied by a simultaneous decrease in muscle tension (Fig. 1A). ACh (10  $\mu$ M) caused a decrease of muscle tension by  $53.0 \pm 10.0\%$  with concomitant increase in  $[Ca^{2+}]_i$  by  $78.4 \pm 10.1\%$  from the steady state values of high  $K^+$ -induced responses. In contrast to endothelium-intact strips, ACh had no significant effect on steady-state values of high  $K^+$ -induced tension and  $[Ca^{2+}]_i$  in endothelium-denuded carotid strips (Fig. 1B). To test whether the ACh-induced changes in  $[Ca^{2+}]_i$  represent the endothelial  $[Ca^{2+}]_i$  transient, we measured the relationship between ACh-induced  $[Ca^{2+}]_i$  transients and the magnitude of relaxations (Fig. 1C), and found a positive correlation between them (correlation coefficient=0.97,  $p < 0.01$ ). We also measured the effects of muscarinic receptor antagonist and nitric oxide synthase (NOS) inhibitor on ACh-induced tension and  $[Ca^{2+}]_i$  transient in endothelium-intact carotid strips (Fig. 2). Pretreatment with 10 nM 4-DAMP, a muscarinic receptor antagonist, had no effect on high  $K^+$ -induced  $[Ca^{2+}]_i$  increase and tension development. However, it completely abolished the ACh-induced increase in  $[Ca^{2+}]_i$  (% change of  $[Ca^{2+}]_i$  in control vs. 4-DAMP treated group;  $53.3 \pm 7.8\%$  vs. 0%,  $n=5$ ) and relaxation responses (% change of relaxation in control vs. 4-DAMP treated group;  $48.1 \pm 6.7\%$  vs. 0%,  $n=5$ ). Pretreatment with 0.1 mM L-NAME, a NOS inhibitor, significantly inhibited the ACh-induced relaxation ( $48.1 \pm 6.7\%$  vs.  $3.4 \pm 2.3\%$  in control vs. L-NAME treated group,  $n=5$ ), while it had no significant effect on the ACh-induced  $[Ca^{2+}]_i$  increase ( $53.3 \pm 7.8\%$  vs.  $48.4 \pm 6.3\%$  in control vs. L-NAME treated group,  $n=5$ ).

### *Effect of lysophosphatidylcholine on ACh-induced additional $[Ca^{2+}]_i$ increase and tension*


As shown in Fig. 3, LPC (10  $\mu$ M) had no significant effect on high  $K^+$ -induced  $[Ca^{2+}]_i$  and tension development in endothelium-intact carotid strips. However, ACh-induced elevations of  $[Ca^{2+}]_i$  and relaxation responses were significantly attenuated by pretreatment with LPC (10  $\mu$ M). In control experiment, ACh caused a relaxation ( $48.1 \pm 6.7\%$  of high  $K^+$ -stimulated level) and elevation of  $[Ca^{2+}]_i$  ( $53.3 \pm 7.8\%$  of high  $K^+$ -stimulated level,  $n=8$ ). In the presence of 10  $\mu$ M LPC, the magnitude of the ACh-induced relaxation was decreased to  $5.2 \pm 4.2\%$  of the high  $K^+$ -stimulated level, and the ACh-induced  $[Ca^{2+}]_i$  transient was also decreased to  $5.5 \pm 3.6\%$  of the high  $K^+$ -stimulated level. This inhibitory effect of LPC on ACh-induced tension and  $[Ca^{2+}]_i$  responses was reversible, because the ACh-induced  $[Ca^{2+}]_i$  elevations and relaxation responses gradually returned to




**Fig. 1.** ACh-induced changes of  $[Ca^{2+}]_i$  and tension in endothelium-intact or denuded rabbit carotid strips. Typical recordings of  $[Ca^{2+}]_i$  (indicated by  $F_{340}/F_{380}$ ; upper trace) and tension (lower trace) in response to ACh in endothelium-intact (A) or endothelium-denuded (B) carotid arterial strips. Muscle strips were pre-contracted with high  $K^+$  (70 mM) and relaxed by cumulative addition of ACh. Changes in  $[Ca^{2+}]_i$  were expressed by measuring the fura-2 fluorescence ratio ( $F_{340}/F_{380}$ ). The steady-state response to high  $K^+$ , obtained before the application of ACh, was taken as 100% and the resting  $[Ca^{2+}]_i$  level as 0%. The magnitude of ACh-induced relaxation was expressed as a percent of high  $K^+$ -induced contraction. (C) Relationship between ACh-induced increases in  $[Ca^{2+}]_i$  (abscissa) and relaxation (ordinate) in endothelium-intact carotid strips. Each point represents the mean of 6 to 8 experiments. S.E.M. is shown by vertical and horizontal bar. The regression line was drawn by least-square method (correlation coefficient=0.97,  $p < 0.01$ ).



**Fig. 2.** Effect of 4-DAMP and L-NAME on ACh-induced changes of  $[Ca^{2+}]_i$  and tension. Typical recordings of the ACh-induced changes of  $[Ca^{2+}]_i$  (upper trace) and tension (lower trace) in the presence of 10 nM 4-DAMP (A) or 0.1 mM L-NAME (B).



**Fig. 3.** Effect of LPC on ACh-induced changes of  $[Ca^{2+}]_i$  and tension in carotid arterial strips. (A) Typical recordings of simultaneous measurement of ACh-induced changes of  $[Ca^{2+}]_i$  (upper trace) and tension (lower trace) in the presence of  $10 \mu M$  LPC. (B) Mean changes of LPC-induced inhibition of ACh-induced relaxation (B) and increase in  $[Ca^{2+}]_i$  (C). Results are expressed as mean  $\pm$  S.E. ( $n=4$ ). \*Significantly different from ACh-induced response with  $p < 0.05$ .



**Fig. 4.** Effect of LPC on ACh-induced changes of  $[Ca^{2+}]_i$  in the absence of external  $Ca^{2+}$ . Typical recordings of ACh-induced changes of  $[Ca^{2+}]_i$  in the absence of external  $Ca^{2+}$ . Muscle strips were stimulated by  $70 mM K^+$  followed by a wash with  $Ca^{2+}$ -free PSS containing  $2 mM$  EGTA. One minute after washing with  $Ca^{2+}$ -free PSS, ACh ( $10 \mu M$ ) was added to the bath and measured the fluorescence change in the presence and absence of LPC ( $10 \mu M$ ). Effect of LPC on  $Ca^{2+}$  transient was evaluated by comparing the peak amplitude of  $Ca^{2+}$  transients. 100% represents the  $70 mM K^+$ -stimulated  $[Ca^{2+}]_i$  before the addition of  $Ca^{2+}$ -free PSS and the resting  $[Ca^{2+}]_i$  level as 0%.

control level after washout of LPC (Fig. 3A).

To elucidate the mechanism of LPC-induced inhibition of ACh-evoked rise in  $[Ca^{2+}]_i$ , we measured the effect of LPC on  $[Ca^{2+}]_i$  transient in the absence of external  $Ca^{2+}$ , in which agonist-induced  $[Ca^{2+}]_i$  transients was solely due to

$Ca^{2+}$  release from internal stores. As shown in Fig. 4, removal of external  $Ca^{2+}$  decreased the high  $K^+$ -induced tension and  $[Ca^{2+}]_i$  to the resting level. Application of ACh after removal of external  $Ca^{2+}$  transiently increased  $[Ca^{2+}]_i$  ( $68.7 \pm 10.3\%$  of high  $K^+$ -stimulated level,  $n=6$ ). Pretrea-

tment with LPC significantly inhibited these ACh-induced transient change in  $[Ca^{2+}]_i$  ( $43.3 \pm 7.3\%$  of high  $K^+$ -stimulated level,  $n=6$ ).

## DISCUSSION

In this study, we simultaneously measured both endothelial  $[Ca^{2+}]_i$  and vascular tension, in order to elucidate the mechanism of LPC-induced suppression of endothelium-dependent relaxation (EDR) at the tissue level. We first examined whether ACh-induced change in Fura-2 signal was due to changes in endothelial  $[Ca^{2+}]_i$  or not, since in vascular strips which contain both endothelium and smooth muscle cells, the Fura-2 signals measured might be due to the changes in  $[Ca^{2+}]_i$  in endothelium or smooth muscle cells. As seen in Fig. 1A & B, ACh-induced changes in Fura-2 signal were observed only in endothelium-intact strips, but not in endothelium-denuded strips. Moreover, there was a positive correlation between the ACh-induced increase in  $[Ca^{2+}]_i$  and the magnitude of relaxation in endothelium-intact strips (Fig. 1C). These findings suggest that, in our experimental conditions, ACh-induced change of Fura-2 signals was almost entirely originated from the vascular endothelium, and it represents endothelial  $[Ca^{2+}]_i$  change. It is possible that ACh-induced increase in endothelial  $[Ca^{2+}]_i$  might have been underestimated, because ACh decreases  $[Ca^{2+}]_i$  in smooth muscle cells by releasing EDRF from endothelium (Rapoport et al, 1983; Grygleski et al, 1988; Karaki et al, 1988). However, the contribution of ACh-induced reduction of  $[Ca^{2+}]_i$  in smooth muscle to the ACh-induced changes in  $[Ca^{2+}]_i$  could be negligible, because there was a close relationship between the ACh-induced increase in  $[Ca^{2+}]_i$  and the magnitude of relaxation ( $p < 0.01$ ,  $n=6$ ).

Agonist-induced changes in endothelial  $[Ca^{2+}]_i$  is an essential steps in the production of bioactive agents, such as nitric oxide (NO). Chelation of endothelial  $Ca^{2+}$  inhibits the production of NO and abolishes the EDR in vascular strips (huang et al, 2001). In the present study, we showed that blocking the ACh-induced  $[Ca^{2+}]_i$  elevation by 4-DAMP completely abolished the ACh-induced EDR responses. And pre-treatment of the strips with L-NAME completely eliminated the ACh-induced relaxation response with little effect on endothelial  $[Ca^{2+}]_i$  elevation (Fig. 2). These findings suggest that elevation of endothelial  $[Ca^{2+}]_i$  is an essential step for the production of endothelial bioactive agents, and NO is the major bioactive agents released from endothelium in our experimental conditions.

Impairment of EDR is a characteristic feature of atherosclerotic artery, and it is well documented that LPC is involved in the pathogenesis of atherosclerosis through inhibition of EDR (Flavahan, 1992; Murohara et al, 1994; Berliner et al, 1995). Similarly, we also observed that LPC inhibited ACh-induced EDR in carotid arteries. There is a possibility that LPC-induced reduction in EDR might be due to lysis of endothelial cells by its cytotoxic effect (Weltzien, 1979). However, the inhibitory effect of LPC was gradually abolished by washing the strip several times with normal PSS solution (Fig. 3A). Therefore, the LPC-induced reduction in EDR was not due to lysis of endothelial cells, but rather due to reversible alteration of endothelial function. However, it is still possible that LPC-induced overproduction of superoxide anion might perturb the efficacy of NO by formation of peroxynitrite, contributing to the LPC-

induced inhibition of EDR (Ohara et al, 1994).

In endothelial cells, the agonist-induced elevation of  $[Ca^{2+}]_i$  was due to the  $Ca^{2+}$  released from internal stores by 1,4,5-triphosphate ( $IP_3$ ) or the influx of  $Ca^{2+}$  from external space. The release of  $Ca^{2+}$  from internal stores represents the transient component in agonist-induced increase in  $[Ca^{2+}]_i$ , whereas the influx of  $Ca^{2+}$  from external stores through capacitive  $Ca^{2+}$  entry pathway represents the sustained component (Putney, 1986; Adams et al, 1989). In the present study, LPC ( $10 \mu M$ ) significantly blocked the ACh-induced elevation of endothelial  $[Ca^{2+}]_i$  in the absence of external  $Ca^{2+}$  (Fig. 4), suggesting the possibility that LPC inhibits the release of  $Ca^{2+}$  from internal stores via disturbing the G-protein coupled  $Ca^{2+}$  signaling in endothelial cells (Hirata et al, 1992; Flavahan, 1993).

Transmembrane  $Ca^{2+}$  influx from external space depends on the membrane potential of endothelial cells. Agonist-induced hyperpolarization through activation of  $Ca^{2+}$ -activated  $K^+$  channels augments the driving force for the  $Ca^{2+}$  influx through capacitive  $Ca^{2+}$  entry pathway (Mehrkke et al, 1991; Nilus, 1991). Therefore, depolarization with any causes can attenuate the amount of  $Ca^{2+}$  influx from external space. Terasawa et al. (2002) reported that LPC depolarized the coronary smooth muscle cells through activation of nonselective cation channels. Thus, LPC may depolarize membrane potential of endothelial cells, leading to the inhibition of transmembrane  $Ca^{2+}$  influx. In the present study, we measured the effect of LPC after  $70 \text{ mM } K^+$ -induced contraction reached steady-state. In this high external  $K^+$  solution, it is expected that the membrane potential of endothelial cells shows strong depolarization, and that the possibility of further depolarization by LPC will be negligible. However, we did not directly evaluate this possibility in the present study. Whether or not LPC inhibits  $Ca^{2+}$  influx through depolarization needs to be investigated further in isolated endothelial cells.

In summary, LPC inhibits the ACh-induced EDR through perturbation of ACh-induced  $[Ca^{2+}]_i$  regulation in rabbit carotid strips. LPC-induced inhibition of endothelial  $[Ca^{2+}]_i$  transient might be due to inhibition of  $Ca^{2+}$  release from internal stores in endothelial cells. Present results are expected to provide us one possible explanation for LPC-induced impairment of endothelium-dependent vascular relaxation.

## ACKNOWLEDGEMENT

This study was supported by a grant from the Yonsei University (Grant No. 2001-01-01).

## REFERENCES

- Adams DJ, Barakeh J, Laskey R, Van Breemen C. Ion channels and regulation of intracellular calcium in vascular endothelial cells. *FASEB J* 3: 2389–2400, 1989
- Berliner JA, Navab M, Fogelman AM. Atherosclerosis: basic mechanisms, oxidation, inflammation and genetics. *Circulation* 66: 2488–2496, 1995
- Boulanger CM, Vanhoutte PM. G proteins and endothelium-dependent relaxations. *J Vasc Res* 34: 175–185, 1997
- Busse R, Mulsch A. Calcium-dependent nitric oxide synthesis in endothelial cytosol is mediated by calmodulin. *FEBS Lett* 265: 133–136, 1990
- Flavahan NA. Atherosclerosis or lipoprotein-induced endothelial

- dysfunction: potential mechanisms underlying reduction in EDRF/nitric oxide activity. *Circulation* 85: 1927–1938, 1992
- Flavahan NA. Lysophosphatidylcholine modifies G-protein dependent signaling in porcine endothelial cells. *Am J Physiol* 264: H722–H727, 1993
- Gryglewski RJ, Botting RM, Vane JR. Mediators produced by the endothelial cell. *Hypertension* 12: 530–548, 1988
- Hirata K, Akita H, Yokoyama M. Oxidized low density lipoprotein inhibits bradykinin-induced phosphoinositide hydrolysis in cultured bovine aortic endothelial cells. *FEBS Lett* 287: 181–184, 1991
- Huang TY, Chen HI, Liu CY, Jen CJ. Endothelial  $[Ca^{2+}]_i$  is an integrating signal for the vascular tone in rat aortae. *BMC Physiol* 1: 5, 2001
- Huang TY, Chen HI, Liu CY, Jen CJ. Lysophosphatidylcholine alters vascular tone in rat aorta by suppressing endothelial  $[Ca^{2+}]_i$  signaling. *J Biomed Sci* 9: 327–333, 2002
- Inoue N, Hirata K, Yamada M, Hamamori Y, Matsuda Y, Akita H, Yokoyama M. Lysophosphatidylcholine inhibits bradykinin-induced phosphoinositide hydrolysis and calcium transients in cultured bovine aortic endothelial cells. *Circ Res* 71: 1410–1421, 1992
- Karaki H, Sato K, Ozaki H, Murakami K. Effect of sodium nitroprusside on cytosolic calcium level in vascular smooth muscle. *Eu J Pharmacol* 156: 259–266, 1988
- Mehrke G, Pohl U, Daut J. Effects of vasoactive agonists on the membrane potential of cultured bovine aortic and guinea-pig coronary endothelium. *J Physiol* 439: 277–299, 1991
- Miwa Y, Hirata KI, Kawashima S, Akita H, Yokoyama M. Lysophosphatidylcholine inhibits receptor-mediated  $Ca^{2+}$  mobilization in intact endothelial cells of rabbit aorta. *Arterioscler Thromb Vasc Biol* 17: 1561–1567, 1994
- Murohara T, Kugiyama K, Ohgushi M, Sugiyama S, Ohta Y, Yasue H. LPC in oxidized LDL elicits vasoconstriction and inhibits endothelium-dependent relaxation. *Am J Physiol* 267: H2441–H2449, 1994
- Nilus B. Regulation of transmembrane calcium fluxes in endothelium. *News Physiol Sci* 6: 110–114, 1991
- Ohara Y, Peterson TE, Zheng B, Kuo JF, Harrison DG. Lysophosphatidylcholine increases vascular superoxide anion production via protein kinase C activation. *Arterioscler Thromb* 14: 1007–1013, 1994
- Parthasarathy S, Steinbrecher UP, Barnett J, Witztum JL, Steinberg D. Essential role of phospholipase  $A_2$  activity in endothelial cell-induced modification of low density lipoprotein. *Proc Natl Acad Sci* 82: 3000–3004, 1985
- Putney JW. A model for receptor regulated calcium entry. *Cell Calcium* 7: 1–12, 1986
- Rapoport RM, Murad F. Agonist-induced endothelium-dependent relaxation in rat thoracic aorta may be mediated through cGMP. *Circ Res* 52: 352–357, 1983
- Sato K, Ozaki H, Karaki H. Differential effects of carbachol on cytosolic calcium levels in vascular endothelium and smooth muscle. *J Pharmacol Exp Ther* 255: 114–119, 1990
- Terasawa K, Nakajima T, Iida H, Iwasawa K, Oonuma H, Jo T, Morita T, Nakamura F, Fujimori Y, Toyoka T, Nagai R. Non-selective cation currents regulate membrane potential of rabbit coronary arterial cell: modulation by lysophosphatidylcholine. *Circulation* 106: 3111–3119, 2002
- Weltzien HU. Cytolytic and membrane-perturbing properties of lysophosphatidylcholine. *Biochim Biophys Acta* 559: 259–287, 1979