

Src Kinase Regulates Nitric Oxide-induced Dedifferentiation and Cyclooxygenase-2 Expression in Articular Chondrocytes via p38 Kinase-dependent Pathway

Seon-Mi Yu¹, Won-Kil Lee¹, Eun-Kyung Yoon¹, Ji-Hye Lee¹, Sun-Ryung Lee² and Song-Ja Kim¹

¹Department of Biological Sciences, Kongju National University College of Natural Sciences, Gongju, ²Department of Biology, College of Natural Sciences, Cheju National University, Jeju, Korea

ABSTRACT

Background: Nitric oxide (NO) in articular chondrocytes regulates dedifferentiation and inflammatory responses by modulating MAP kinases. In this study, we investigated whether the Src kinase in chondrocytes regulates NO-induced dedifferentiation and cyclooxygenase-2 (COX-2) expression. **Methods:** Primary chondrocytes were treated with various concentrations of SNP for 24 h. The COX-2 and type II collagen expression levels were determined by immunoblot analysis, and prostaglandin E₂ (PGE₂) was determined by using a PGE₂ assay kit. Expression and distribution of p-Caveolin and COX-2 in rabbit articular chondrocytes and cartilage explants were determined by immunohistochemical staining and immunocytochemical staining, respectively. **Results:** SNP treatment stimulated Src kinase activation in a dose-dependent manner in articular chondrocytes. The Src kinase inhibitors PP2 [4-amino-5-(4-chlorophenyl)-7-(t-butyl)pyrazolo(3,4-d)pyrimidine], a significantly blocked SNP-induced p38 kinase and caveolin-1 activation in a dose-dependent manner. Therefore, to determine whether Src kinase activation is associated with dedifferentiation and/or COX-2 expression and PGE₂ production. As expected, PP2 potentiated SNP-stimulated dedifferentiation, but completely blocked both COX-2 expression and PGE₂ production. And also, levels of p-Caveolin and COX-2 protein expression were increased in SNP-treated primary chondrocytes and osteoarthritic and rheumatoid arthritic cartilage, suggesting that p-Caveolin may play a role in the inflammatory responses of arthritic cartilage. **Conclusion:** Our previous studies indicated that NO caused dedifferentiation and COX-2 expression is regulated by p38 kinase through caveolin-1 (1). Therefore, our results collectively suggest that Src kinase regulates NO-induced dedifferentiation and COX-2 expression in chondrocytes via p38 kinase in association with caveolin-1. (*Immune Network* 2006;6(4):204-210)

Key Words: Src kinase, Caveolin-1, Dedifferentiation, COX-2, p38 kinase

Introduction

Chondrocytes in cartilage are differentiated from mesenchymal cells during embryonic development. Differentiated chondrocytes, which are the only cell type found in normal mature cartilage, synthesize

sufficient amounts of cartilage-specific extracellular matrix (ECM) to maintain matrix integrity (1). This homeostasis is destroyed in degenerative diseases, such as osteoarthritis and rheumatoid arthritis. Arthritis is characterized by structural and biochemical changes in chondrocytes and cartilage, including degradation of cartilage matrix, insufficient synthesis of ECM because of loss of chondrocyte phenotype. However, the differentiated phenotype is unstable both *in vivo* and *in vitro* and thus lost by a process designated "dedifferentiation" upon exposure of cells to interleukin-1 β (1,2), nitric oxide (NO) (3), or retinoic acid (4,5) and during serial monolayer culture (6,7). NO mediates the regulation of chondrocyte

Correspondence to: Song-Ja Kim, Department of Biological Sciences, Kongju National University, Gongju 314-701, Korea (Tel) 82-41-850-8507, (Fax) 82-41-850-8479, (E-mail) ksj85@kongju.ac.kr

This work was supported by Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2005-202-100654), Rheumatoid Research Center Program (KOSEF Grant R11-2002-098-04005-0).

phenotype and survival by inducing dedifferentiation and apoptosis (3,8,9). We showed previously that direct production of NO by treatment of primary culture articular chondrocytes with a NO donor, sodium nitroprusside (SNP), led to apoptosis, dedifferentiation, and cyclooxygenase (COX)-2 expression via a complex protein kinase signaling cascade involving mitogen-activated protein (MAP) kinase and protein kinase C (PKC) (10-13). For example, NO-induced activation of extracellular signal-regulated protein kinase (ERK) promotes dedifferentiation, COX-2 expression, and inhibition of apoptosis, whereas NO-induced p38 kinase activation triggers apoptosis, COX-2 expression, and maintains differentiated phenotypes (10,11). And also, we recently observed that p38 kinase regulates dedifferentiation in association with caveolin-1 phosphorylation (14).

Caveolin, a family of 21 ~ 24 kDa integral membrane proteins are a principal component of caveolae membranes (15). Although expression of caveolae structure and expression of caveolin family, caveolin-1, -2 and -3, was known in chondrocytes, the exact functional role of caveolae and caveolins in chondrocytes remain unclear (16).

Src family kinases (SFKs) are one of the most studied groups of tyrosine kinases and can mediate a variety of signaling pathways. c-Src signaling played a major role in transducing MSU-induced NO production and MMP-3 expression via p38 activation. Thus, c-Src as novel sites for potential therapeutic intervention in cartilage degradation (17). In addition to, the specific activity of pp60c-Src kinase regulate matrix mineralization in hypertrophic chondrocyte culture (18). However, little is known about the expression of SFKs in articular chondrocytes.

In the studies described herein we examined the role of Src kinase using a PP2, a potent and selective inhibitor of the Src family tyrosine kinases (13) on SNP-induced dedifferentiation and inflammatory responses. We found that dedifferentiation, COX-2 expression, and PGE₂ production by NO was regulated by Src kinase signaling pathway via activation of p38 kinase and p-Caveolin.

Materials and Methods

Isolation and monolayer culture of rabbit articular chondrocytes and experimental culture condition. Articular chondrocytes were isolated from cartilage slices of 2-week-old New Zealand white rabbits by enzymatic digestion as described previously (19). Cartilage slices were dissociated enzymatically for 6 h in 0.2% collagenase type II (381 U/mg solid, Sigma) in Dulbecco's modified Eagle's medium (DMEM, Gibco-BRL, Gaithersburg, MD). Individual cells were suspended in Dulbecco's modified Eagle's medium supplemented with 10%

(v/v) fetal bovine-calf serum, 50 µg/ml streptomycin, and 50 units/ml penicillin, after which and they were then plated on culture dishes at a density of 5×10^4 cells/cm². The medium was changed every 2 days after seeding, and cells reached confluence in approximately 5 days. Differentiation status of articular chondrocytes was determined by examining the accumulation of sulfated glycosaminoglycan with Alcian Blue staining or expression of type II collagen expression was detected using antibodies purchased from Chemicon (Temecula, CA) by immunoblot analysis as described previously (19).

Immunoblot analysis. Whole cell lysates were prepared by extracting proteins using a buffer containing 50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Nonidet P-40, and 0.1% sodium dodecylsulfate, supplemented with protease inhibitors [10 µg/ml leupeptin, 10 µg/ml pepstatin A, 10 µg/ml aprotinin and 1 mM of 4-(2-aminoethyl) benzenesulfonyl fluoride] and phosphatase inhibitors (1 mM NaF and 1 mM Na₃VO₄). The proteins were size-fractionated by SDS-polyacrylamide gel electrophoresis and transferred to a nitrocellulose membrane. The nitrocellulose sheet was then blocked with 3% non-fat dry milk in Tris-buffered saline. COX-2 was detected using antibody purchased from Cayman Chemical (Ann Arbor, MI), and p-Caveolin and pSrc were detected using antibodies purchased from Santa Cruz Biotech. (Santa Cruz, CA). pp38 was purchased from Chemicon (Temecula, CA). Blotss were developed using a peroxidase-conjugated secondary antibody and visualized with an ECL system.

Immunohistochemistry and immunofluorescence microscopy. Rabbit joint cartilage explants or arthritic cartilage were in 4% paraformaldehyde in PBS for 24 h at 4°C, washed with PBS, dehydrated with graded ethanol, embedded in paraffin, and sectioned at 4 µm thickness. The sections were stained by standard procedures using Alcian blue or antibody against p-Caveolin or COX-2 and visualized by developing with a kit purchased from DAKO (Carpinteria, CA), following the procedure recommended by the manufacturer. Expression and distribution of type II collagen and COX-2 in rabbit articular chondrocytes were determined by indirect immunofluorescence microscopy, as described previously (19). Briefly, chondrocytes were fixed with 3.5% paraformaldehyde in PBS for 10 min at room temperature. The cells were permeabilized and blocked with 0.1% Triton X-100 and 5% fetal calf serum in PBS for 30 min. The fixed cells were washed and incubated for 1 h with antibody (10 µg/ml) against type II collagen or COX-2. The cells were washed, incubated with rhodamine-conjugated secondary antibodies for 30 min, and observed under a fluorescence microscope.

PGE₂ Assay. PGE₂ production was determined by measuring the levels of cellular and secreted PGE₂ using an assay kit (Amersham Pharmacia Biotech, NJ, UK). Briefly, chondrocytes were seeded in standard 96-well microtiter plates at 2×10^4 cells/well. Following addition of the indicated pharmacological reagents, total cell lysate was used to quantify the amount of PGE₂, according to the manufacturer's protocol. PGE₂ levels were calculated against a standard curve of PGE₂ and normalized against the amount of genomic DNA.

Data analyses and statistics. The results are expressed as the means \pm S.E. values calculated from the specified

number of determinations. A student's test was used to compare individual treatments with their respective control values. A probability of $p < 0.05$ was taken as denoting a significant difference.

Results

SNP treatment induces dedifferentiation of articular chondrocytes. To examine the effects of NO generator SNP on articular cartilage chondrocytes, cartilage explant cultures and primary culture chondrocytes were treated with 1 mM SNP for 24 h, and expression of cartilage-




Figure 1. NO causes dedifferentiation of chondrocytes. Cartilage explants (left panel) and primary culture chondrocytes (right panel) were untreated or treated with 1 mM SNP for 24 h. Proteoglycan and type II collagen were detected by Alcian blue staining and immunocytochemical staining and, respectively (A). Chondrocytes were treated with 1 mM SNP for various time periods (upper panel) or with the indicated concentrations of SNP for 24 h (lower panel). The expression of type II collagen was determined immunoblot analysis (B). Accumulation of sulfated glycosaminoglycan in cells treated with the indicated concentrations of SNP for 24 h was quantified by Alcian Blue staining (C). The data represent the results of a typical experiment conducted at least three times with similar results.




Figure 2. NO causes COX-2 expression and PGE₂ production in articular chondrocytes. Primary culture chondrocytes were untreated or treated with 1 mM SNP for 24 h. COX-2 expression was detected by immunocytochemical staining (A). Chondrocytes were treated with 1 mM SNP for various time periods (upper panel) or with the indicated concentrations of SNP for 24 h (lower panel). The expression of COX-2 was determined immunoblot analysis (B). Levels of cellular and secreted PGE₂ were determined by using a PGE₂ assay kit (C). The data in A represent the results of a typical experiment conducted four times, and B signifies the average values with standard deviation ($n=4$).

specific matrix molecules were determined. NO caused a dramatic loss of type II collagen and sulfated proteoglycan as determined by immunostaining and Alcian blue staining, respectively (Fig. 1A). NO also inhibited type II collagen expression in articular chondrocytes in a time- and dose-dependent manner (Fig. 1B). Similarly, SNP treatment of articular chondrocytes inhibited the accumulation of sulfated proteoglycan in a dose-dependent manner (Fig. 1C). These results indicate that NO induces dedifferentiation in both cartilage explants and primary culture cells. *SNP treatment induces COX-2 expression and PGE₂ production in articular chondrocytes.* The effect of NO on COX-2 expression and PGE₂ production was investigated in articular chondrocytes. In primary culture articular chondrocytes, NO stimulated COX-2 expression as determined by immunocytochemical staining (Fig. 2A). And also, NO induce COX-2 expression in a time- and dose-dependent manner as determined by immunoblot analysis (Fig. 2B). Consistent with the induction of COX-2 expression, NO stimulated PGE₂ production (Fig. 2C). These data indicate that NO stimulates COX-2 expression and

PGE₂ production in articular chondrocytes.

Src kinase stimulates dedifferentiation and COX-2 expression. To investigate the signaling pathway stimulating dedifferentiation and COX-2 expression of SNP-treated chondrocytes, we examined changes in the activity of Src and p38 kinase, and activation of caveolin-1. As shown in Fig. 3A, Src kinase activity was increased in a time-dependent manner as determined by phosphorylation status of the protein. Consistent with our previous observation, levels of p38 kinase phosphorylation began to increase at 3 h, reached peak levels at 12 h, and decreased thereafter. Similar to the pattern of Src activation, caveolin-1 activation was also increased in a time-dependent manner (Fig. 3A). To determine the association between Src kinase and p38 kinase, chondrocytes were treated with SNP in the presence and absence of PP2, and expression of p38 activation as determined by immunoblot analysis. The addition of PP2 to SNP-treated chondrocytes blocked the increased phosphorylation of p38 kinase in a dose-dependent manner (Fig. 3B, middle panel). And, caveolin-1 tyrosine phosphorylation is also blocked by inhibition of Src kinase (Fig. 3B, lower




Figure 3. Activation of Src kinase, p38 kinase and caveolin-1 in SNP-treated chondrocytes. Chondrocytes were treated with 1 mM SNP for various time periods (A). Chondrocyte cells were treated with 1 mM SNP for 24 h in the presence of indicated concentrations of PP2 (B). pSrc, pp38 and p-Caveolin-1 (p-Cav) were detected by immunoblot analyses. The data represent the results of a typical experiment conducted at least three times with similar results.




Figure 4. Src kinase induces dedifferentiation and COX-2 expression. Articular chondrocyte cells were treated with 1 mM SNP for 24 h in the presence of indicated concentrations of PP2. Type II collagen, COX-2 and pSrc were detected by immunoblot analyses (A). Chondrocytes were treated with 1 mM SNP for 24 h in the absence or presence of indicated concentrations of 20 μM PP2 or 20 μM SB203580. Accumulation of sulfated glycosaminoglycan was quantified by Alcian blue staining (B). Levels of cellular and secreted PGE₂ were determined by using a PGE₂ assay kit (C). The data represent the average values with standard deviation (n=4).




Figure 5. Increased levels of p-Caveolin and COX-2 in arthritic cartilage. Cartilage were untreated or treated with 1 mM SNP for 24 h. Expression levels of p-Caveolin and COX-2 was detected by immunohistochemical staining (A). Expression of p-Caveolin and COX-2 proteins was detected from normal (Control) and osteoarthritis-affected (OA) and rheumatoid-affected human joint cartilage (RA) (B). Tissue sections were counter stained with hematoxylin. The data represent the results of a typical experiment conducted at least three times.

panel).

To determine whether Src kinase stimulated dedifferentiation and COX-2 expression, chondrocytes were treated with 1 mM SNP in the presence and absence of PP2. As shown in Fig. 4A, type II collagen expression, a differentiation marker of articular chondrocytes (17), was completely blocked (upper panel), and COX-2 expression was blocked in a dose-dependent manner (middle panel). To elucidate the above data, we next examined sulfated proteoglycan accumulation and PGE₂ production as determined by Alcian blue staining and using a kit assay, respectively. Similarly, SNP treatment of primary culture cells blocked the accumulation of sulfated proteoglycan (Fig. 4B) and PGE₂ production by PP2 (Fig. 4C). The inhibition of sulfated proteoglycan accumulation and PGE₂ production by inhibition of p38 kinase with SB203580 is consistent with our previous data (14,20). Taken together, these results indicate that activity of Src kinase is required for COX-2 expression, and subsequent PGE₂ production induced by NO.

Increased levels of caveolin-1 and COX-2 in cartilage explants culture. To examine the in vivo significance of p-Caveolin and COX-2 expression in cartilage, explants of rabbit joint cartilage were cultured in the absence or presence of SNP. SNP treatment of cartilage explants caused increased p-Caveolin protein levels with a concomitant increase in COX-2 expression (Fig. 5A). We also examined the expression levels of p-Caveolin and COX-2 in human osteoarthritic and rheumatoid arthritic cartilage obtained from patients undergoing total knee arthroplasty. Relatively normal parts of the joint (*i.e.*, undamaged parts of the osteoarthritic joint) showed undetectable expression of p-Caveolin and COX-2 (Fig. 5B). However, ex-

pression of p-Caveolin and COX-2 was significantly increased in osteoarthritis-affected part of cartilage and rheumatoid arthritic cartilage. Taken together, these results suggest that the increased expression levels of p-Caveolin and COX-2 may contribute to the cartilage destruction during arthritic disease.

Discussion

Our previous studies in primary articular chondrocytes indicated that NO-induced apoptosis and dedifferentiation of articular chondrocytes were regulated by opposite functions of mitogen-activated protein (MAP) kinase subtypes, extracellular signal-regulated protein kinase (ERK), and p38 kinase (10). NO-induced activation of ERK-1/-2 induces dedifferentiation with the inhibitory effects on apoptosis, whereas activation of p38 kinase induces apoptosis and is responsible for the maintenance of differentiated phenotype. In addition to MAP kinase signaling, NO production stimulates expression of inflammatory molecules such as COX-2 and PGE₂ (10,11). We also previously demonstrated that activation of caveolin-1 is necessary for NO-induced dedifferentiation, COX-2 expression, and PGE₂ production through p38 kinase activation (14). Volonte et al. (21), demonstrated that tyrosine phosphorylation of caveolin-1 on tyrosine 14 was mediated by the activation of p38 kinase and Src tyrosine kinase.

We characterized signaling pathways involved in NO-induced dedifferentiation and COX-2 expression in articular chondrocytes, focusing on the functional relationship between Src kinase and p38 kinase signaling. Here, we show that pSrc and p38 kinase activated by NO in a similar kinetic pattern. The addition of PP2 to SNP-treated chondrocytes inhibited the increase phosphorylation of p38 kinase and

p-Caveolin and slowed the inhibition of both type II collagen expression and proteoglycan synthesis. Similar to the pattern of type II collagen, COX-2 expression and PGE₂ production were also blocked in SNP-treated cells in the presence of PP2.

Several inflammatory mediators, including nitric oxide and PG, have been implicated in the disease process of osteo- and rheumatoid arthritis. PGs are produced by cytokine-stimulated chondrocytes in arthritis-affected cartilage via induction of COX-2 expression (8,22-25). Our current observation of increased expression of COX-2 in arthritis-affected chondrocytes is consistent with the reports by Amin et al. (8) and Jacques et al. (26), showing increased expression and activity of COX-2 in osteoarthritic cartilage. And also, we first demonstrated increased expression levels of p-Caveolin in arthritis-affected chondrocytes. Based on the observation that high levels of p-Caveolin in arthritic cartilage might be associated with increased COX-2 expression.

These data suggest that Src kinase activation is necessary to NO-induced dedifferentiation and COX-2 expression in articular chondrocytes through p38 kinase-dependent pathway.

References

1. Goldring MB, Birkhead JR, Suen LF, Yamin R, Mizuno S, Glowacki J, Arbisser JL, Apperley JF: Interleukin-1 beta-modulated gene expression in immortalized human chondrocytes. *J Clin Invest* 94;2307-2316, 1994
2. Demoor-Fossard M, Redini F, Boittin M, Pujol JP: Expression of decorin and biglycan by rabbit articular chondrocytes. Effects of cytokines and phenotypic modulation. *Biochim Biophys Acta* 1398;179-191, 1998
3. Amin AR, Abramson SB: The role of nitric oxide in articular cartilage breakdown in osteoarthritis. *Curr Opin Rheumatol* 10;263-268, 1998
4. Cash DE, Bock CB, Schughart K, Linney E, Underhill TM: Retinoic acid receptor alpha function in vertebrate limb skeletogenesis: a modulator of chondrogenesis. *J Cell Biol* 136;445-457, 1997
5. Weston AD, Rosen V, Chandraratna RA, Underhill TM: Regulation of skeletal progenitor differentiation by the BMP and retinoid signaling pathways. *J Cell Biol* 148;679-690, 2000
6. Lefebvre V, Peeters-Joris C, Vaes G: Production of collagens, collagenase and collagenase inhibitor during the dedifferentiation of articular chondrocytes by serial subcultures. *Biochim Biophys Acta* 1051;266-275, 1990
7. Yoon YM, Kim SJ, Oh CD, Ju JW, Song WK, Yoo YJ, Huh TL, Chun JS: Maintenance of differentiated phenotype of articular chondrocytes by protein kinase C and extracellular signal-regulated protein kinase. *J Biol Chem* 277;8412-8420, 2002
8. Amin AR, Attur M, Abramson SB: Nitric oxide synthase and cyclooxygenases: distribution regulation and intervention in arthritis. *Curr Opin Rheumatol* 11;202-209, 1999
9. Sandell LJ, Aigner I: Articular cartilage and changes in arthritis. An introduction: cell biology of osteoarthritis. *Arthritis Res* 3;107-113, 2001
10. Kim SJ, Ju JW, Oh CD, Yoon YM, Song WK, Kim JH, Yoo YJ, Bang OS, Kang SS, Chun JS: ERK-1/2 and p38 kinase oppositely regulate nitric oxide-induced apoptosis of chondrocytes in association with p53, caspase-3, and differentiation status. *J Biol Chem* 277;1332-1339, 2002
11. Kim SJ, Kim HG, Oh CD, Hwang SG, Song WK, Yoo YJ, Kang SS, Chun JS: p38 kinase-dependent and -independent inhibition of protein kinase C zeta and -alpha regulates nitric oxide-induced apoptosis and dedifferentiation of articular chondrocytes. *J Biol Chem* 277;30375-30381, 2002
12. Kim SJ, Hwang SG, Shin DY, Kang SS, Chun JS: p38 kinase regulates nitric oxide-induced apoptosis of articular chondrocytes by accumulating p53 via NF-kappa B-dependent transcription and stabilization by serine 15 phosphorylation. *J Biol Chem* 277;33501-33508, 2002
13. Hanke JH, Gardner JP, Dow RL, Changlianb PS, Brissette WH, Weringer EJ, Pollok BA, Connelly PA: Discovery of a novel, potent, and Src family-selective tyrosine kinase inhibitor: Study of Lck and FynT-dependent T cell activation. *J Biol Chem* 271;695-701, 1996
14. Yu SM, Cheong SW, Cho SR, Kim SJ: p38 kinase regulates nitric oxide-induced dedifferentiation and cyclooxygenase-2 expression of articular chondrocytes. *Immune Network* 6; 117-122, 2006
15. Rothberg KG, Heuser JE, Donzell WC, Ying YS, Glenney JR, Anderson RG: Caveolin, a protein component of caveolae membrane coats. *Cell* 68;673-682, 1992
16. Schwartz Z, Gilley RM, Sylvia VL, Dean DD, Boyan BD: The effect of prostaglandin E2 on costochondral chondrocyte differentiation is mediated by cyclic adenosine 3',5'-monophosphate and protein kinase C. *Endocrinology* 139; 1825-1834, 1998
17. Liu R, Liote F, Rose DM, Merz D, Terkeltaub R: Proline-rich tyrosine kinase 2 and Src kinase signaling transducer monosodium urate crystal-induced nitric oxide production and matrix metalloproteinase 3 expression in chondrocytes. *Arthritis Rheum* 50;247-258, 2004
18. Coe MR, Summers TA, Parsons SJ, Boskey AL, Balian G: Matrix mineralization in hypertrophic chondrocyte cultures. Beta glycerophosphate increases type X collagen messenger RNA and the specific activity of pp60c-Src kinase. *Bone Miner* 18;91-106, 1992
19. Yoon YM, Kim SJ, Oh CD, Ju JW, Song WK, Yoo YJ, Huh TL, Chun JS: Maintenance of dedifferentiated phenotype of articular chondrocytes by protein kinase C and extracellular signal-regulated protein kinase. *J Biol Chem* 277;8412-8420, 2002
20. Kim SJ, Hwang SG, Kim IC, Chun JS: Actin cytoskeletal architecture regulates nitric oxide-induced apoptosis, dedifferentiation, and cyclooxygenase-2 expression in articular chondrocytes via mitogen-activated protein kinase and protein kinase C pathways. *J Biol Chem* 278;42448-42456, 2003
21. Volonte D, Galbiati F, Pestell RG, Lisanti MP: Cellular stress induces the tyrosine phosphorylation of caveolin-1 [Tyr (14)] via activation of p38 mitogen-activated protein kinase and c-Src kinase. Evidence for caveolae, the actin cytoskeleton, and focal adhesions as mechanical sensors of osmotic stress. *J Biol Chem* 276;8094-8103, 2001
22. Abramson SB: The role of COX-2 produced by cartilage in arthritis. *osteoarthritis cartilage* 7;380-381, 1999
23. Gilroy DW, Tomlinson A, Greenslade K, Seed MP, Willoughby DA: The effects of cyclooxygenase-2 inhibitors on cartilage erosion and bone loss in a model of *Mycobacterium tuberculosis*-induced monoarticular arthritis in the rat. *Inflammation* 22;509-519, 1998
24. Myers LK, Kang AH, Postlethwaite AE, Rosloniec EF, Morham SG, Goorha S, Ballou LR: The genetic ablation of

- cyclooxygenase-2 prevents the development of autoimmune arthritis. *Arthritis Rheum* 43:2687-2693, 2000
25. Anderson GD, Hauser SD, McGarity KL, Bremer ME, Isakson PC, Gregory SA: Selective inhibition of cyclooxygenase (COX)-2 reverses inflammation and expression of COX-2 and interleukin 6 in rat adjuvant arthritis. *J Clin Invest* 97:2672-2679, 1996
26. Jaques C, Sautet A, Moldovan M, Thomas B, Humbert L, Berenbaum F: cyclooxygenase activity in chondrocytes from osteoarthritic and healthy cartilage. *Rev Rheum Engl Ed* 66:701-704, 1999
-