

p38 Kinase Regulates Nitric Oxide-induced Dedifferentiation and Cyclooxygenase-2 Expression of Articular Chondrocytes

Seon-Mi Yu¹, Seon-Woo Cheong², Sam-Rae Cho¹ and Song-Ja Kim¹

¹Department of Biological Sciences, Kongju National University, College of Natural Sciences, Gongju, ²Department of Biology, College of Natural Sciences, Changwon National University, Changwon, Korea

ABSTRACT

Background: Caveolin, a family of integral membrane proteins are a principal component of caveolae membranes. In this study, we investigated the effect of p38 kinase on differentiation and on inflammatory responses in sodium nitroprusside (SNP)-treated chondrocytes. **Methods:** Rabbit articular chondrocytes were prepared from cartilage slices of 2-week-old New Zealand white rabbits by enzymatic digestion. SNP was used as a nitric oxide (NO) donor. In this experiments measuring SNP dose response, primary chondrocytes were treated with various concentrations of SNP for 24 h. The time course of the SNP response was determined by incubating cells with 1 mM SNP for the indicated time period (0~24 h). The cyclooxygenase-2 (COX-2) and type II collagen expression levels were determined by immunoblot analysis, and prostaglandin E₂ (PGE₂) assay was used to measure the COX-2 activity. The tyrosine phosphorylation of caveolin-1 was determined by immunoblot analysis and immunostaining. **Results:** SNP treatment stimulated tyrosine phosphorylation of caveolin-1 and activation of p38 kinase. SNP additionally caused dedifferentiation and inflammatory response. We showed previously that SNP treatment stimulated activation of p38 kinase and ERK-1/-2. Inhibition of p38 kinase with SB203580 reduced caveolin-1 tyrosine phosphorylation and COX-2 expression but enhanced dedifferentiation, whereas inhibition of ERK with PD98059 did not affect caveolin-1 tyrosine phosphorylation levels, suggesting that ERK at least is not related to dedifferentiation and COX-2 expression through caveolin-1 tyrosine phosphorylation. **Conclusion:** Our results indicate that SNP in articular chondrocytes stimulates dedifferentiation and inflammatory response via p38 kinase signaling in association with caveolin-1 phosphorylation. (*Immune Network* 2006;6(3):117-122)

Key Words: SNP, caveolin-1, dedifferentiation, inflammatory response, p38 kinase

Introduction

Chondrocytes are differentiated from mesenchymal cells during embryo development (1,2). The phenotype of the differentiated chondrocyte is characterized by the synthesis, deposition, and maintenance of cartilage-specific extracellular matrix (ECM) molecules, including type II collagen and proteoglycans (3-5). Dedifferentiation is a major restriction in mass cell production for cell therapy or tissue engineering of destructive cartilage. Also, dedifferentiation of chondrocytes involves a gradual shift from the synthesis of type II to types I and III collagen. Nitric oxide

(NO) produced via inducible NO synthase in articular chondrocytes plays a central role in cartilage diseases such as osteoarthritis and rheumatoid arthritis. NO causes cartilage destruction by inducing apoptosis, dedifferentiation, and inflammatory responses such as cyclooxygenase (COX)-2 expression and prostaglandin E₂ (PGE₂) production in articular chondrocytes (6-9). We showed previously that direct production of NO by treatment with NO donor, sodium nitroprusside (SNP), causes dedifferentiation (10) and COX-2 expression (11) via MAP kinase signaling in rabbit articular chondrocytes. For example, NO-induced activation of extracellular signal-regulated protein kinase (ERK) promotes dedifferentiation, COX-2 expression, and inhibition of apoptosis, whereas NO-induced p38 kinase activation triggers apoptosis, COX-2 expression, and maintains differentiated phenotypes (10,11).

Correspondence to: Song-Ja Kim, Department of Biological Sciences, Kongju National University, 182, Shinkwan-dong, Gongju 314-701, Korea (Tel) 82-41-850-8507, (Fax) 82-41-850-8479 (E-mail) ksj85@kongju.ac.kr

Caveolin, a family of 21~24 kDa membrane protein that binds many different signaling molecules such as heterotrimeric G proteins, Src, Ras and calmodulin (12-16). Caveolin-1 is also directly involved in signaling cascades as a substrate of both serine and tyrosine kinases. Caveolin was first identified as a major tyrosine-phosphorylated protein in v-Src-transformed embryonic fibroblasts (17) and was demonstrated that tyrosine phosphorylation of caveolin-1 on tyrosine 14 was mediated by the activation of p38 kinase and Src tyrosine kinase (18). Phosphorylation of caveolin-1 on tyrosine is likely to be an intermediate step in a signaling cascade occurring within caveolae.

The present study investigate the roles of caveolin-1 tyrosine phosphorylation in the regulation of dedifferentiation and COX-2 expression of SNP-treated chondrocytes. The investigation additionally focuses on the characterization of the role of MAP kinase subtypes in dedifferentiation and COX-2 expression. We report here that p38 kinase regulates nitric oxide-induced dedifferentiation and COX-2 expression of articular chondrocytes in a caveolin-1 tyrosine phosphorylation-dependent pathway.

Materials and Methods

Monolayer culture of rabbit articular chondrocytes and experimental culture condition. Rabbit articular chondrocytes were released from cartilage slices of 2-week-old New Zealand white rabbits by enzymatic digestion. To summarize, after aseptic dissection cartilage slices were aseptically dissected and then dissociated enzymatically for 6 h in 0.2% collagenase type II (381 U/mg solid, Sigma) in PBS, and individual cells were then obtained by collecting the supernatant after brief centrifugation. The cells were resuspended in Dulbecco's modified Eagle's medium (DMEM, Gibco-BRL, Gaithersburg, MD) supplemented with 10% (v/v) fetal bovine-calf serum, 50 μ g/ml streptomycin, and 50 units/ml penicillin, after which and they were then plated on culture dishes at a density of 5×10^4 cells/cm². The medium was changed every 2 days after seeding, and cells reached confluence in approximately 5 days.

Determination of chondrocyte differentiation status. Loss of chondrocyte phenotype, dedifferentiation, was determined by examining the accumulation of sulfated glycosaminoglycan with Alcian Blue, as described previously (19,20) or by the expression of type II collagen studied by immunoblot analysis. To summarize, type II collagen expression was detected using antibodies purchased from Chemicon (Temecula, CA) by immunoblotting, as described below.

Immunoblot analysis. Whole cell lysates were prepared by extracting proteins using a buffer containing 50

mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Nonidet P-40, and 0.1% sodium dodecylsulfate, supplemented with protease inhibitors [10 μ g/ml leupeptin, 10 μ g/ml pepstatin A, 10 μ g/ml aprotinin and 1 mM of 4-(2-aminoethyl) benzenesulfonyl fluoride] and phosphatase inhibitors (1 mM NaF and 1 mM Na₃VO₄). The proteins were size-fractionated by SDS-polyacrylamide gel electrophoresis and transferred to a nitrocellulose membrane. The nitrocellulose sheet was then blocked with 3% non-fat dry milk in Tris-buffered saline. COX-2 was detected using antibody purchased from Cayman Chemical (Ann Arbor, MI). Caveolin-1, p-Caveolin-1, ERK-2 and p38 were detected using antibodies purchased from Santa Cruz Biotech. (Santa Cruz, CA). The bands were visualized using peroxidase-conjugated secondary antibodies and chemiluminescence.

Immunofluorescence microscopy. Tyrosine phosphorylation of caveolin-1 in rabbit articular chondrocytes was investigated using an indirect immunofluorescence microscopy, as described previously (20). Briefly, chondrocytes were fixed with 3.5% paraformaldehyde in PBS for 10 min at room temperature. The cells were permeabilized and blocked with 0.1% Triton X-100 and 5% fetal calf serum in PBS for 30 min. The fixed cells were washed and incubated for 1 h with antibody (10 μ g/ml) against anti-phospho-caveolin-1 antibodies and with fluorescein-conjugated goat anti-mouse antibodies for 30 min. Cells were then mounted with low-Fade, and observed under a fluorescence microscope.

p38 kinase assay. p38 kinase activity was determined by immune complex kinase assays as described previously (19,20). Briefly, cells were lysed in a buffer containing 20 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton-X 100, 2.5 mM sodium pyrophosphate, 1 mM β -glycerolphosphate and inhibitors of proteases and phosphatases, as described above. Total cell lysates were precipitated with polyclonal anti-p38 kinase antibody and the immune complexes collected using protein-A Sepharose beads. The beads were re-suspended in 20 μ l kinase reaction buffer containing 25 mM Tris-HCl, pH 7.5, 5 mM DTT, 0.1 mM sodium orthovanadate, 10 mM MgCl₂, 5 μ Ci [γ -³²P] ATP and 1 μ g of substrate activating transcription factor-2 (ATF-2) for p38 kinase (New England Biolabs, Beverly, MA). The kinase reaction was performed for 30 min at 30°C and phosphorylated ATF-2 was detected by autoradiography following gel electrophoresis.

PGE₂ assay. PGE₂ production was determined by measuring the levels of cellular and secreted PGE₂ using an assay kit (Amersham Pharmacia Biotech, NJ, UK). Briefly, chondrocytes were seeded in standard 96-well microtiter plates at 2×10^4 cells/well. Follo-




Figure 1. Tyrosine phosphorylation of caveolin-1 and p38 kinase in SNP-treated chondrocytes. A and B, rabbit articular chondrocytes were treated with 1 mM SNP for the indicated time periods (A) or for 24 h with the specified concentrations of SNP (B). Activation of Cav-1 (caveolin-1) was determined by immunoblot analysis, using antibody specific to activated p-Caveolin-1. p38 kinase activity was determined by immunocomplex kinase assay, using ATF-2 as a substrate. Expression of Caveolin-1 and p38 MAP kinase was determined by immunoblot analysis. The data in A represent the results of a typical experiment conducted four times, and B signifies the average values with standard deviation (n=4).




Figure 2. Dedifferentiation in SNP-treated chondrocytes. Chondrocytes were treated with the indicated concentrations of SNP for 24 h. The expression of type II collagen (collagen II) was determined immunoblot analysis (A). Accumulation of sulfated glycosaminoglycan in cells treated with the indicated concentrations of SNP for 24 h was quantified by Alcian Blue staining (B). The data represent the results of a typical experiment conducted at least three times with similar results.

wing addition of the indicated pharmacological reagents, total cell lysate was used to quantify the amount of PGE₂, according to the manufacturer's protocol. PGE₂ levels were calculated against a standard curve of PGE₂ and normalized against the amount of genomic DNA.

Data analyses and statistics. The results are expressed as the means ± S.E. values calculated from the specified number of determinations. A Student's test was used to compare individual treatments with their respective control values. A probability of $p < 0.05$ was taken as denoting a significant difference.




Figure 3. SNP treatment induces COX-2 expression in articular chondrocytes. Chondrocytes were treated with the indicated concentrations of SNP for 24 h (A, C) or with 1 mM SNP for specified time period (B). The expression of COX-2 and ERK-2 was determined immunoblot analysis (A, B), and production of PGE₂ was determined by using a PGE₂ assay kit (C). The data represent the results of a typical experiment conducted at least three times with similar results.

Results

Nitric-oxide induces phosphorylation of caveolin-1 and p38 kinase. Rabbit articular chondrocytes in primary culture were treated with the NO donor SNP. SNP caused a dramatic increased caveolin-1 tyrosine phosphorylation and p38 kinase activity as determined by immunoblotting and immunocomplex kinase assay, respectively, in a dose- and time-dependent manner, as shown in Fig. 1A and Fig. 1B. But expression of caveolin-1 slightly increased. These results indicate that NO induces caveolin-1 tyrosine phosphorylation and p38 kinase activation.

NO induces dedifferentiation, COX-2 expression, and PGE₂ production of articular chondrocytes. To examine the effects of SNP on articular cartilage chondrocyte on dedifferentiation, COX-2 expression, and PGE₂ production, primary cell cultures were treated with the indicated concentrations of SNP for 24 h or with 1 mM SNP for specified time periods. As expected, expression of type II collagen (Fig. 2A) and sulfated proteoglycan synthesis (Fig. 2B) were significantly reduced in a dose-dependent manner in SNP-treated chondrocytes. But, SNP increased protein levels of COX-2 in a time- and dose-dependent manner as determined by immunoblotting (Fig. 3A,B). Consistent with the induction of COX-2 expression, SNP stimulated PGE₂ production (Fig. 3C). These data indicate that NO not only causes dedifferentiation of articular chondrocytes but also stimulates COX-2 expression and PGE₂ production.

NO-induced activation of p38 kinase causes tyrosine phosphorylation of caveolin-1. To determine the association between caveolin-1 and p38 kinase activation, chondro-

cytes were treated with SNP in the presence and absence of SB203580, an inhibitor of p38 kinase and expression and tyrosine phosphorylation of caveolin-1 were examined. The addition of SB203580 to SNP-treated chondrocytes completely blocked the increased tyrosine phosphorylation of caveolin-1 in a dose dependent manner (Fig. 4A). But, caveolin-1 expression is slightly decreased by SB203580. In contrast to the effects of p38 kinase inhibition, blockade of ERK with PD98059, an inhibitor of ERK-1/-2 upstream kinase and did not affect tyrosine phosphorylation and expression of caveolin-1 (Fig. 4B). In an attempt to elucidate the mechanism of p38 kinase regulated caveolin-1 tyrosine phosphorylation, chondrocytes were treated with SNP in the absence or presence of




Figure 4. NO-induced activation of p38 kinase but not ERK-1/-2 causes tyrosine phosphorylation of caveolin-1. A and B, cells were treated with 1 mM SNP for 24 h in the presence of indicated concentrations of SB203580 (A) or PD98059 (B). Caveolin, p-Caveolin-1 and ERK-2 were detected by immunoblot analyses. The data represent the average values with standard deviation ($n=4$).

SB203580. SNP treatment resulted in strong staining of caveolin-1 tyrosine phosphorylation, and SB203580 abolished the effects of SNP (Fig. 5). This phenomenon is consistent with immunoblotting. These results indicate that NO-induced caveolin-1 tyrosine phosphorylation is regulated by p38 kinase activity, but not ERK-1/-2 activity.

p38 kinase regulates NO-induced dedifferentiation, COX-2 expression and PGE₂ production. A recent study demonstrated that SNP treatment of articular chondrocytes causes both dedifferentiation (10) and inflammatory response (11). To investigate the expression levels of type II collagen and COX-2, chondrocytes were treated with SNP in the presence and absence of SB203580. Consistent with our previous data (10,11), blockade of p38 kinase accelerated NO-induced decrease of type II collagen expression and COX-2 expression in a dose-dependent manner (Fig. 6A). Similar to the effects on the above results, inhibition of p38 kinase potentiated NO-induced decrease in the accumulation of sulfated proteoglycan (Fig. 6B) and PGE₂ production (Fig. 6C). Taken together, these results indicate that NO-induced dedifferentiation, COX-2 expression and PGE₂ production is regulated through p38 kinase pathway.

Discussion

Our previously studies indicated that NO caused apoptosis and dedifferentiation, which are mediated by MAP kinase subtypes ERK and p38 kinase (10). These MAP kinase play opposing roles, with activated ERK-1/-2 inducing dedifferentiation and inhibiting NO-induced apoptosis, while p38 kinase signaling maintains the differentiated status and induces apoptosis. We also previously demonstrated that NO-stimulates expression of inflammatory molecules such as COX-2 and PGE₂ in chondrocytes via ERK and p38 kinase (11,21).

Caveolin-1 is directly involved in signaling cascades as a substrate of both serine and tyrosine kinases.




Figure 5. NO-induced tyrosine phosphorylation of caveolin-1. Chondrocytes were untreated or treated with 20 μ M SB203580 and exposed to 1 mM SNP for 24 h. Tyrosine phosphorylated caveolin-1 (p-Caveolin-1) was immunostained, and photographs were taken with an immunofluorescence microscope. The data represent the results of a typical experiment conducted at least three times.




Figure 6. p38 kinase regulates dedifferentiation and COX-2 expression/PGE₂ production. A and C, articular chondrocytes were treated with 1 mM SNP for 24 h in the absence or presence of indicated concentrations of SB203580. Type II collagen (Collagen II) and COX-2 were detected by immunoblotting. Chondrocytes were untreated or treated with 20 μ M SB203580 and exposed to 1 mM SNP for 24 h. Accumulation of sulfated glycosaminoglycan was quantified by Alcian blue staining (B). Production of PGE₂ was determined by using a PGE₂ assay kit. The data represent the results of a typical experiment conducted at least three times.

Caveolin-1 is phosphorylated on tyrosine 14 by Src, Fyn and Abl in response to a number of stimuli, including insulin, angiotensin II, osmotic shock, and oxidative stress (22-24). Numerous studies have demonstrated that caveolin interacts with a number of signaling molecules that are thought to be enriched in caveolae, including certain small and heterotrimeric G proteins (13,25), endothelial nitric oxide synthase (26,27) and a subset of tyrosine kinase receptors (28). In fact, caveolin-1 is a preferred substrate for these tyrosine kinases in cells (29-32).

We first demonstrated signaling pathways involved in NO-induced dedifferentiation, COX-2 expression, and PGE₂ production in rabbit articular chondrocytes, focusing on the functional relationship between caveolin-1 and p38 kinase. Based on results obtained in this study and our previous reports (10,21,33), we propose the following pathway: NO activation of p38 kinase activates caveolin-1 tyrosine phosphorylation. This activation of caveolin-1 is necessary for NO-induced dedifferentiation, COX-2 expression and PGE₂ production. In this regard, p38 kinase have been proposed to interact with caveolin. This possibility will be the subject of future studies.

References

- Solursh M: Differentiation of cartilage and bone. *Curr Opin Cell Biol* 1;989-994, 1989
- Sandell LJ, Adler P: Developmental patterns of cartilage. *Front Biosci* 4;731-742, 1999
- Hauselman HJ, Fernandes RJ, Mok SS, Schmid TM, Block JA, Aydelotte MB, Kuettner KE, Thonar JMA: Phenotypic stability of bovine articular chondrocytes after long-term culture in alginate beads. *J Cell Sci* 107;17-27, 1994
- Reginato AM, Iozzo RV, Jimenez SA: Formation of nodular structures resembling mature articular cartilage in long-term primary cultures of human fetal epiphyseal chondrocytes on a hydrogel substrate. *Arthritis Rheum* 37;1338-1349, 1994
- Archer CW, McDowell J, Baileys MT, Stephens MD, Bentley G: Phenotypic modulation in sub-populations of human articular chondrocytes in vitro. *J Cell Sci* 97;361-371, 1990
- Sandell LJ, Aigner I: Articular cartilage and changes in arthritis. An introduction: cell biology of osteoarthritis. *Arthritis Res* 3;107-113, 2001
- Amin AR, Abramson SB: The role of nitric oxide in articular cartilage breakdown in osteoarthritis. *Curr Opin Rheumatol* 10;263-268, 1998
- Amin AR, Attur M, Abramson SB: Nitric oxide synthase and cyclooxygenases: distribution regulation and intervention in arthritis. *Curr Opin Rheumatol* 11;202-209, 1999
- Abramson SB, Attur M, Amin AR, Clancy R: Nitric oxide and inflammatory mediators in the perpetuation of osteoarthritis. *Curr Rheumatol Rep* 3;535-541, 2001
- Kim SJ, Ju JW, Oh CD, Yoon YM, Song WK, Kim JH, Yoo YJ, Bang OS, Kang SS, Chun JS: ERK-1/2 and p38 kinase oppositely regulate nitric oxide-induced apoptosis of chondrocytes in association with p53, caspase-3, and differentiation status. *J Biol Chem* 277;1332-1339, 2002
- Kim SJ, Chun JS: Protein kinase C- α and - ζ regulate nitric-oxide-induced NF- κ B activation that mediates cyclooxygenase-2 expression and apoptosis but not dedifferentiation in articular chondrocytes. *Biochem Biophys Res Commun* 303;206-211, 2003
- Li S, Okamoto T, Chun M, Sagicomo M, Casanova JE, Hansen SH, Nishimoto I, Lisanti MP: Evidence for a regulated interaction between heterotrimeric G proteins and caveolin. *J Biol Chem* 270;15693-15701, 1995
- Li S, Couet J, Lisanti MP: Src tyrosine kinases, G- α subunits, and H-Ras shear a common membrane-anchored scaffolding protein, caveolin. Caveolin binding negatively regulates the auto-activation of Src tyrosine kinases. *J Biol Chem* 271;29182-29190, 1996
- Song KS, Li S, Okamoto T, Quilliam LA, Sargiacomo M, Lisanti MP: Co-purification and direct interaction of Ras with caveolin, an integral membrane protein of caveolae microdomains. Detergent-free purification of caveolae microdomains. *J Biol Chem* 271;9690-9697, 1996

15. Michel JB, Feron O, Sase K, Prabhakar P, Michel T: Caveolin versus calmodulin. Counterbalancing allosteric modulators of endothelial nitric oxide synthase. *J Biol Chem* 272;25907-25912, 1997
16. Okamoto T, Schlegel A, Scherer PE, Lisanti MP: Caveolin, a family of scaffolding proteins for organizing "preassembled signaling complexes" at the plasma membrane. *J Biol Chem* 273;5419-5422, 1998
17. Glenny JR, Zokas L: Novel tyrosine kinase substrates from Rous sarcoma virus-transformed cells are present in the membrane skeleton. *J Cell Biol* 108;2401-2408, 1989
18. Volonte D, Galbiati F, Pestell RG, Lisanti MP: Cellular stress induces the tyrosine phosphorylation of caveolin-1 [Tyr(14)] via activation of p38 mitogen-activated protein kinase and c-Src kinase. Evidence for caveolae, the actin cytoskeleton, and focal adhesions as mechanical sensors of osmotic stress. *J Biol Chem* 276;8094-8103, 2001
19. Oh, CD, Chang SH, Yoon YM, Lee SJ, Lee YS, Kang SS, Chun JS: Opposing role of mitogen-activated protein kinase subtypes, Erk-1/-2 and p38, in the regulation of chondrogenesis of mesenchymes. *J Biol Chem* 275;5613-5619, 2000
20. Yoon YM, Kim SJ, Oh CD, Ju JW, Song WK, Yoo YJ, Huh TL, Chun JS: Maintenance of differentiated phenotype of articular chondrocytes by protein kinase C and extracellular signal-regulated protein kinase. *J Biol Chem* 277;8412-8420, 2002
21. Kim SJ, Hwang SG, Kim IC, Chun JS: Actin cytoskeletal architecture regulates nitric-oxide-induced apoptosis, dedifferentiation, and cyclooxygenase-2 expression in articular chondrocytes via mitogen-activated protein kinase and protein kinase C pathways. *J Biol Chem* 278;42448-42456, 2003
22. Mastick CC, Brady MJ, Saltiel AR: Insulin stimulates the tyrosine phosphorylation of caveolin. *J Cell Biol* 129;1523-1531, 1995
23. Mastick CC, Saltiel AR: Insulin-stimulated tyrosine phosphorylation of caveolin is specific for the differentiated adipocyte phenotype in 3T3-L1 cells. *J Biol Chem* 272;20706-20714, 1997
24. Ushio-Fukai M, Hilenski L, Santanam N, Becker P, Ma Y, Griendling K, Alexander R: Cholesterol depletion inhibits epidermal growth factor receptor transactivation by angiotensin II in vascular smooth muscle cells: role of cholesterol-rich microdomains and focal adhesions in angiotensin II signaling. *J Biol Chem* 276;48269-48275, 2001
25. Oh P, Schnitzer JE: Segregation of heterotrimeric G proteins in cell surface microdomains. G(q) binds caveolin to concentrate in caveolae, whereas G(i) and G(s) target lipid rafts by default. *Mol Biol Cell* 12;685-698, 2001
26. Garcia-Cardena G, Oh P, Liu J, Schnitzer JE, Sessa WC: Targeting of nitric oxide synthase to endothelial cell caveolae via palmitoylation: implications for nitric oxide signaling. *Proc Natl Acad Sci USA* 93;6448-6453, 1996
27. Feron O, Belhassen L, Kobzik L, Smith TW, Kelly RA, Michel T: Endothelial nitric oxide synthase targeting to caveolae. Specific interactions with caveolin isoforms in cardiac myocytes and endothelial cells. *J Biol Chem* 271;22810-22814, 1996
28. Liu P, Ying Y, Ko YG, Anderson RG: Localization of platelet-derived growth factor-stimulated phosphorylation cascade to caveolae. *J Biol Chem* 271;10299-10303, 1996
29. Glenny JR Jr: Tyrosine phosphorylation of a 22-kDa protein is correlated with transformation by Rous sarcoma virus. *J Biol Chem* 264;20163-20166, 1989
30. Ko YG, Liu P, Pathak RK, Craig LC, Anderson RGW: Early effects of pp60(v-src) kinase activation on caveolae. *J Cell Biochem* 71;524-535, 1998
31. Nomura R, Fujimoto T: Tyrosine-phosphorylated caveolin-1: immunolocalization and molecular characterization. *Mol Biol Cell* 10;975-986, 1999
32. Corley Mastick C, Sanguinetti AR, Knesek JH, Mastick GS, Newcomb LF: Caveolin-1 and a 29-kDa caveolin-associated protein are phosphorylated on tyrosine in cells expressing a temperature-sensitive v-Abl kinase. *Exp Cell Res* 266;142-154, 2001
33. Kim SJ, Kim HG, Oh CD, Hwang SG, Song WK, Yoo YJ, Kang SS, Chun JS: p38 kinase-dependent and -independent inhibition of protein kinase C zeta and alpha regulates nitric-oxide-induced apoptosis and dedifferentiation of articular chondrocytes. *J Biol Chem* 277;30375-30381, 2002