

NOTES ON THE BERGMAN PROJECTION TYPE OPERATOR IN \mathbb{C}^n

KI SEONG CHOI

ABSTRACT. In this paper, we will define the Bergman projection type operator P_r and find conditions on which the operator P_r is bound-ed on $L^p(B, d\nu)$. By using the properties of the Bergman projection type operator P_r , we will show that if $f \in L^p_a(B, d\nu)$, then $(1 - \|w\|^2)\nabla f(w) \cdot z \in L^p(B, d\nu)$. We will also show that if $(1 - \|w\|^2) \frac{\nabla f(w) \cdot z}{\langle z, w \rangle} \in L^p(B, d\nu)$, then $f \in L^p_a(B, d\nu)$.

1. Introduction

Throughout this paper, \mathbb{C}^n will be the Cartesian product of n copies of \mathbb{C} . For $z = (z_1, z_2, \dots, z_n)$ and $w = (w_1, w_2, \dots, w_n)$ in \mathbb{C}^n , the inner product is defined by $\langle z, w \rangle = \sum_{j=1}^n z_j \bar{w}_j$ and the norm by $\|z\|^2 = \langle z, z \rangle$.

Let B be the open unit ball in the complex space \mathbb{C}^n . Let ν be the Lebesgue measure in \mathbb{C}^n normalized by $\nu(B) = 1$. We let $L^2(B, d\nu)$ be the usual space of Lebesgue square-integrable complex valued functions on B . The Bergman space $L^2_a(B, d\nu)$ is defined to be the subspace of $L^2(B, d\nu)$ consisting of analytic functions.

The measure μ_r is the weighted Lebesgue measure:

$$d\mu_r(z) = c_r(1 - \|z\|^2)^r d\nu(z),$$

where $r > -1$ is fixed, and c_r is a normalization constant such that $\mu_r(B) = 1$.

If we equip $L^2_{a,r} = L^2_a(B, d\mu_r)$ with the norm $\|f\|_{2,r} = \sqrt{\int_B |f|^2 d\mu_r}$, then $L^2_{a,r}$ is a Banach space for each $r > -1$. The orthogonal projection

Received May 9, 2005. Revised June 8, 2005.

2000 Mathematics Subject Classification: 32H25, 32E25, 30C40.

Key words and phrases: Bergman space, Bergman projection.

operator from $L^2(B, d\mu_r)$ to $L^2_{a,r}$ is denoted by P and P is called the Bergman projection. The Bergman projection is used in many areas related with Hankel operators and Toeplitz operators (See [2, 6, 9, 14, 15]).

The space $L^2_{a,r}$ is a Hilbert space with the inner product

$$\langle f, g \rangle = \int_B f(z) \overline{g(z)} d\mu_r(z).$$

Fix a point $z \in B$. Since the functional e_z given by $e_z(f) = f(z)$, $f \in L^2_{a,r}$, is continuous, there exists a function $k_{r,z} \in L^2_{a,r}$ such that

$$f(z) = \int_B f(w) \overline{k_{r,z}(w)} d\mu_r(w)$$

by the Riesz representation theorem. The function $K_r(z, w) = \overline{k_{r,z}(w)}$ is called the weighted Bergman kernel. Also it is well known that

$$K_r(z, w) = \frac{1}{(1 - \langle z, w \rangle)^{r+n+1}}$$

(See [12]). It was shown in [10] that if $f \in L^1_{a,r}$, $r > -1$, then

$$f(z) = c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} f(w) d\nu(w).$$

Suppose $1 \leq p < +\infty$ and $r > 0$. Let $L^p_{a,r}$ be the subspace of $L^p(B, d\mu_r)$ consisting of analytic functions. Define the Bergman projection type operator P_r by

$$\begin{aligned} P_r f(z) &= \int_B f(w) \overline{k_{r,z}(w)} d\mu_r(w) \\ &= c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} f(w) d\nu(w). \end{aligned}$$

Since $P_r f = f$ for all analytic f in $L^1(B, d\mu_r)$, P_r is a projection from $L^1(B, d\mu_r)$ onto $L^1_a(B, d\mu_r)$. In general, P_r is not a bounded operator (See Corollary 6).

In section 2, we will find conditions on which the operator P_r is a bounded operator from $L^p(B, d\mu_r)$ onto $L^p_{a,r}$. In particular, we will

show that if $1 \leq p < +\infty$ and $r > 0$, then the operator P_r is a bounded projection from $L^p(B, d\nu)$ onto $L_a^p(B, d\nu)$.

Let N denote the set of natural numbers. A multi-index α is an ordered n -tuple $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ with $\alpha_j \in N, j = 1, 2, \dots, n$. For a multi-index α and $z \in \mathbb{C}^n$, set

$$\begin{aligned} |\alpha| &= \alpha_1 + \alpha_2 + \dots + \alpha_n, \\ \alpha! &= \alpha_1! \alpha_2! \dots \alpha_n!, \\ z^\alpha &= z_1^{\alpha_1} z_2^{\alpha_2} \dots z_n^{\alpha_n}. \end{aligned}$$

For all multi-indices α , we will write

$$\partial^\alpha f(w) = \frac{\partial^{|\alpha|} f(w)}{\partial w_1^{\alpha_1} \dots \partial w_n^{\alpha_n}}.$$

For a holomorphic function f , we will also write

$$\nabla f(w) \cdot z = \frac{\partial f(w)}{\partial w_1} z_1 + \dots + \frac{\partial f(w)}{\partial w_n} z_n.$$

Suppose $p \geq 1, |\alpha| \geq 1$ and $f \in L_a^p(B, d\nu)$. In section 3, we will show that

$$(1 - \|w\|^2)^{|\alpha|} \partial^\alpha f(w)$$

is in $L^p(B, d\nu)$. By using this result and the properties of the Bergman type projection in section 2, we will show that if $f \in L_a^p(B, d\nu)$, then $(1 - \|w\|^2) \nabla f(w) \cdot z \in L^p(B, d\nu)$. We will also show that if $(1 - \|w\|^2) \frac{\nabla f(w) \cdot z}{\langle z, w \rangle} \in L^p(B, d\nu)$, then $f \in L_a^p(B, d\nu)$.

2. Bergman projection type operator P_r

THEOREM 1. For $z \in B$, c is real, $t > -1$, define

$$I_{c,t}(z) = \int_B \frac{(1 - \|w\|^2)^t}{|1 - \langle z, w \rangle|^{n+1+c+t}} d\nu(w).$$

Then,

- (i) $I_{c,t}(z)$ is bounded in B if $c < 0$;
- (ii) $I_{0,t}(z) \sim -\log(1 - \|z\|^2)$ as $\|z\| \rightarrow 1^-$;
- (iii) $I_{c,t}(z) \sim (1 - \|z\|^2)^{-c}$ as $\|z\| \rightarrow 1^-$ if $c > 0$.

PROOF. See [12, Proposition 1.4.10]. \square

THEOREM 2. Suppose (X, μ) is a measure space and K is a measurable function on $X \times X$. Let T be the integral operator induced by K , that is,

$$Tf(x) = \int_X K(x, y)f(y)d\mu(y).$$

Suppose $1 < p < +\infty$ with $\frac{1}{p} + \frac{1}{q} = 1$. If there is a constant $c > 0$ and a positive measurable function h on X such that

$$\int_X |K(x, y)|h(y)^q d\mu(y) \leq ch(x)^q$$

for μ -almost every x in X and

$$\int_X |K(x, y)|h(x)^p d\mu(x) \leq ch(y)^p$$

for μ -almost every y in X , then T is bounded on $L^p(X, d\mu)$ with norm less than or equal to c .

PROOF. See [16, Theorem 3.2.2]. \square

THEOREM 3. If $f \in L^1_{a,r}$, $r > -1$, then

$$f(z) = c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} f(w) d\nu(w).$$

PROOF. See [10, Theorem 2]. \square

THEOREM 4. X and Y are Banach spaces. $\mathcal{L}(X, Y)$ is the set of bounded linear transformations from X to Y . If A^* is the adjoint of $A \in \mathcal{L}(X, Y)$, then:

- (1) $\|A^*\| = \|A\|$;
- (2) if A is invertible, then A^* is invertible and $(A^*)^{-1} = (A^{-1})^*$.

PROOF. See [7, Proposition 1.4]. \square

THEOREM 5. Suppose $1 \leq p < +\infty$ and $r > 0$. Then the operator P_r is a bounded projection from $L^p(B, d\nu)$ onto $L^p_a(B, d\nu)$.

PROOF. We first prove the case $p = 1$. If P_r is bounded on $L^p(B, d\nu)$,

$$\begin{aligned} & \langle P_r f, g \rangle \\ &= \int_B P_r f(z) \overline{g(z)} d\nu(z) \\ &= \int_B \int_B \frac{c_r (1 - \|w\|^2)^r f(w)}{(1 - \langle z, w \rangle)^{n+r+1}} d\nu(w) \overline{g(z)} d\nu(z) \\ &= \int_B c_r (1 - \|w\|^2)^r f(w) \int_B \frac{\overline{g(z)}}{(1 - \langle z, w \rangle)^{n+r+1}} d\nu(z) d\nu(w), \end{aligned}$$

where $g \in L^\infty(B)$. Let P_r^* be the adjoint of P_r under the usual integral pairing. Then above result shows that

$$P_r^* g(w) = c_r (1 - \|w\|^2)^r \int_B \frac{g(z)}{(1 - \langle w, z \rangle)^{n+r+1}} d\nu(z).$$

By Theorem 1, if $r > 0$

$$\sup_{w \in B} (1 - \|w\|^2)^r \int_B \frac{d\nu(z)}{(1 - \langle w, z \rangle)^{n+r+1}} < \infty.$$

This shows that if $r > 0$, then P_r^* is bounded on $L^\infty(B, d\nu)$. By Theorem 4, P_r is bounded on $L^1(B, d\nu)$ if $r > 0$.

Next assume $1 < p < +\infty$, $\frac{1}{p} + \frac{1}{q} = 1$. Let $h(z) = (1 - \|z\|^2)^{-\frac{1}{pq}}$.

$$\begin{aligned} & \int_B \frac{c_r (1 - \|w\|^2)^r}{|1 - \langle z, w \rangle|^{n+r+1}} (1 - \|w\|^2)^{-\frac{1}{p}} d\nu(w) \\ &= \int_B \frac{c_r (1 - \|w\|^2)^{r - \frac{1}{p}}}{|1 - \langle z, w \rangle|^{n+r+1}} d\nu(w) \\ &= c_r (1 - \|z\|^2)^{-\frac{1}{p}} \\ & \int_B \frac{c_r (1 - \|w\|^2)^r}{|1 - \langle z, w \rangle|^{n+r+1}} (1 - \|z\|^2)^{-\frac{1}{q}} d\nu(z) \\ &= c_r (1 - \|w\|^2)^r \int_B \frac{(1 - \|z\|^2)^{-\frac{1}{q}}}{|1 - \langle z, w \rangle|^{n+r+1}} d\nu(z) \\ &= c_r (1 - \|w\|^2)^r \int_B \frac{(1 - \|z\|^2)^{-\frac{1}{q}}}{|1 - \langle z, w \rangle|^{n+(r+\frac{1}{q})+1-\frac{1}{q}}} d\nu(z). \end{aligned}$$

Since $r > 0$, then $r + \frac{1}{q} > 0$. This shows that

$$\int_B \frac{c_r(1 - \|w\|^2)^r}{|1 - \langle z, w \rangle|^{n+r+1}} (1 - \|z\|^2)^{-\frac{1}{q}} d\nu(z) \leq c_r(1 - \|w\|^2)^{-\frac{1}{q}}.$$

By Theorem 2, P_r is bounded on $L^p(B, d\nu)$ if $r > 0$. If $f \in L^p_a(B, d\nu)$, then $f \in L^1(B, d\mu_r)$. Since $P_r f = f$, the operator P_r is a bounded projection from $L^p(B, d\nu)$ onto $L^p_a(B, d\nu)$. \square

COROLLARY 6. *If $rq \leq -1$ where $\frac{1}{p} + \frac{1}{q} = 1$, then P_r is not bounded.*

PROOF.

$$\begin{aligned} & \int_B \{P_r^* 1(w)\}^q d\nu(w) \\ &= \int_B \{c_r(1 - \|w\|^2)^r \int_B \frac{d\nu(z)}{(1 - \langle w, z \rangle)^{n+r+1}}\}^q d\nu(w) \\ &= \int_B c_r^q (1 - \|w\|^2)^{rq} \left\{ \int_B \frac{d\nu(z)}{(1 - \langle w, z \rangle)^{n+r+1}} \right\}^q d\nu(w). \end{aligned}$$

$\int_B \frac{d\nu(z)}{(1 - \langle w, z \rangle)^{n+r+1}} d\nu(w)$ is bounded by Theorem 1, but $\int_B (1 - \|w\|^2)^{rq} d\nu(w) = \infty$ since $rq \leq -1$. This shows that if $rq \leq -1$, then P_r is not bounded. \square

COROLLARY 7. *If $1 < p < +\infty$, then P is bounded on $L^p(B, d\nu)$.*

PROOF. This follows from Theorem 5. \square

3. Results related with P_r

Remember that, for all multi-indices $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$,

$$\partial^\alpha f(w) = \frac{\partial^{|\alpha|} f(w)}{\partial w_1^{\alpha_1} \dots \partial w_n^{\alpha_n}}.$$

For a multi-index $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ and $z = (z_1, z_2, \dots, z_n) \in \mathbb{C}^n$, set

$$\bar{z}^\alpha = \bar{z}_1^{\alpha_1} \bar{z}_2^{\alpha_2} \dots \bar{z}_n^{\alpha_n}.$$

LEMMA 8. Suppose $p \geq 1$, $|\alpha| \geq 1$ and $f \in L_a^p(B, d\nu)$. Then

$$(1 - \|z\|^2)^{|\alpha|} \partial^\alpha f(z)$$

is in $L^p(B, d\nu)$.

PROOF. We first prove the case $p = 1$. By Theorem 3,

$$f(z) = c_1 \int_B \frac{(1 - \|w\|^2)}{(1 - \langle z, w \rangle)^{n+2}} f(w) d\nu(w).$$

$$\begin{aligned} \partial^\alpha f(z) &= \frac{\partial^{|\alpha|} f(w)}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}} \\ &= c_1 \int_B \frac{(1 - \|w\|^2)(n+2) \dots (n+|\alpha|+1) f(w) \bar{w}^\alpha}{(1 - \langle z, w \rangle)^{n+2+|\alpha|}} f(w) d\nu(w). \end{aligned}$$

$$\begin{aligned} & \left| \int_B (1 - \|z\|^2)^{|\alpha|} \partial^\alpha f(z) d\nu(z) \right| \\ & \leq |c_1| \frac{(n+|\alpha|+1)!}{(n+1)!} \int_B (1 - \|z\|^2)^{|\alpha|} \\ & \quad \times \left| \int_B \frac{(1 - \|w\|^2) \bar{w}^\alpha}{(1 - \langle z, w \rangle)^{n+2+|\alpha|}} f(w) d\nu(w) \right| d\nu(z) \\ & \leq |c_1| \frac{(n+|\alpha|+1)!}{(n+1)!} \int_B (1 - \|w\|^2) |f(w)| \\ & \quad \times \int_B \frac{(1 - \|z\|^2)^{|\alpha|}}{|1 - \langle z, w \rangle|^{n+2+|\alpha|}} d\nu(z) d\nu(w) \\ & \leq |c_1| \frac{(n+|\alpha|+1)!}{(n+1)!} \int_B (1 - \|w\|^2) (1 - \|w\|^2)^{-1} |f(w)| d\nu(w) \\ & \leq |c| \int_B |f(w)| d\nu(w). \end{aligned}$$

This shows that if f is in $L_a^1(B, d\nu)$, then $(1 - \|z\|^2)^{|\alpha|} \partial^\alpha f(z)$ is in $L^1(B, d\nu)$.

Next assume $1 < p < +\infty$. By Theorem 3,

$$f(z) = \int_B \frac{f(w)}{(1 - \langle z, w \rangle)^{n+1}} d\nu(w).$$

$$\begin{aligned}
& (1 - \|z\|^2)^{|\alpha|} \partial^\alpha f(z) \\
&= (1 - \|z\|^2)^{|\alpha|} \frac{(n + |\alpha|)!}{n!} \int_B \frac{\bar{w}^\alpha f(w)}{(1 - \langle z, w \rangle)^{n+1+|\alpha|}} d\nu(w) \\
&= (1 - \|z\|^2)^{|\alpha|} \frac{(n + |\alpha|)!}{n!} \int_B \frac{f_1(w)}{(1 - \langle z, w \rangle)^{n+1+|\alpha|}} d\nu(w) \\
&= \frac{1}{c_{|\alpha|}} \frac{(n + |\alpha|)!}{n!} c_{|\alpha|} (1 - \|z\|^2)^{|\alpha|} \int_B \frac{f_1(w)}{(1 - \langle z, w \rangle)^{n+1+|\alpha|}} d\nu(w) \\
&= \frac{(n + |\alpha|)!}{n!} \frac{1}{c_{|\alpha|}} P_{|\alpha|}^* f_1(z),
\end{aligned}$$

where $f_1(w) = \bar{w}^\alpha f(w)$. Since $P_{|\alpha|}$ is bounded on $L^q(B, d\nu)$ (where $\frac{1}{p} + \frac{1}{q} = 1$) by Theorem 4, $P_{|\alpha|}^*$ is bounded on $L^p(B, d\nu)$ and

$$(1 - \|w\|^2)^{|\alpha|} \partial^\alpha f(w)$$

is in $L^p(B, d\nu)$. □

THEOREM 9. *Suppose that $p \geq 1$. If $f \in L_a^p(B, d\nu)$, then $(1 - \|w\|^2) \nabla f(w) \cdot z \in L^p(B, d\nu)$.*

PROOF. By Lemma 8, $(1 - \|w\|^2) \frac{\partial f(w)}{\partial w_i} \in L^p(B, d\nu)$ for $i = 1, \dots, n$.

$$\begin{aligned}
& (1 - \|w\|^2) |\nabla f(w) \cdot z| \\
&\leq (1 - \|w\|^2) \sqrt{\left(\frac{\partial f(w)}{\partial w_1}\right)^2 + \dots + \left(\frac{\partial f(w)}{\partial w_n}\right)^2} \\
&\leq (1 - \|w\|^2) \left(\left|\frac{\partial f(w)}{\partial w_1}\right| + \dots + \left|\frac{\partial f(w)}{\partial w_n}\right| \right).
\end{aligned}$$

This shows that $(1 - \|w\|^2) \nabla f(w) \cdot z \in L^p(B, d\nu)$. □

THEOREM 10. *Suppose $z \in B$ and $\nabla f(w) \cdot z \in L^1(B, d\mu_r)$. Then*

$$f(z) = f(0) + \frac{c_r}{n+r} \int_B \frac{(1 - \|w\|^2)^r \nabla f(w) \cdot z}{\langle z, w \rangle (1 - \langle z, w \rangle)^{n+r}} d\nu(w).$$

PROOF. See [10, Theorem 3]. □

THEOREM 11. Suppose $1 \leq p < +\infty$. If $(1 - \|w\|^2) \frac{\nabla f(w) \cdot z}{\langle z, w \rangle} \in L^p(B, d\nu)$, then $f \in L_a^p(B, d\nu)$.

PROOF. Note that $(1 - \|w\|^2) \frac{\nabla f(w) \cdot z}{\langle z, w \rangle} \in L^p(B, d\nu)$ implies $(1 - \|w\|^2) \nabla f(w) \cdot z \in L^p(B, d\nu)$. We first consider the case $p = 1$. Since $\nabla f(w) \cdot z \in L^1(B, d\mu_2)$, by Theorem 10,

$$\begin{aligned} & f(z) - f(0) \\ &= \frac{c_2}{n+2} \int_B \frac{(1 - \|w\|^2)^2 \nabla f(w) \cdot z}{\langle z, w \rangle (1 - \langle z, w \rangle)^{n+2}} d\nu(w) \\ &= \frac{c_2}{n+2} \frac{1}{c_1} \int_B \frac{c_1 (1 - \|w\|^2) (1 - \|w\|^2) \nabla f(w) \cdot z}{(1 - \langle z, w \rangle)^{n+2} \langle z, w \rangle} d\nu(w) \\ &= \frac{c_2}{n+2} \frac{1}{c_1} (P_1 F_z)(z), \end{aligned}$$

where $F_z(w) = \frac{(1 - \|w\|^2) \nabla f(w) \cdot z}{\langle z, w \rangle}$. Since $F_z \in L^1(B, d\nu)$, $P_1 F_z \in L_a^1(B, d\nu)$ by Theorem 5. This shows that $f \in L_a^1(B, d\nu)$.

Now suppose that $1 < p < +\infty$. Since $(1 - \|w\|^2) \nabla f(w) \cdot z \in L^p(B, d\nu)$ and $p > 1$, $(1 - \|w\|^2) \nabla f(w) \cdot z \in L^1(B, d\nu)$. This implies that $\nabla f(w) \cdot z \in L^1(B, d\mu_1)$. By Theorem 10,

$$\begin{aligned} f(z) - f(0) &= \frac{c_1}{n+1} \int_B \frac{(1 - \|w\|^2) \nabla f(w) \cdot z}{\langle z, w \rangle (1 - \langle z, w \rangle)^{n+1}} d\nu(w) \\ &= \frac{c_1}{n+1} \int_B \frac{1}{(1 - \langle z, w \rangle)^{n+1}} \frac{(1 - \|w\|^2) \nabla f(w) \cdot z}{\langle z, w \rangle} d\nu(w) \\ &= \frac{c_1}{n+1} (P F_z)(z), \end{aligned}$$

where $F_z(w) = \frac{(1 - \|w\|^2) \nabla f(w) \cdot z}{\langle z, w \rangle}$. Since $F_z \in L^p(B, d\nu)$, $P F_z \in L_a^p(B, d\nu)$ by Corollary 7. This shows that $f \in L_a^p(B, d\nu)$. \square

References

- [1] J. Anderson, J. Clunie and Ch. Pommerenke, *On Bloch functions and normal functions*, J. Reine Angew. Math. **270** (1974), 12–37.
- [2] J. Arazy, S. D. Fisher, and J. Peetre, *Hankel operators on weighted Bergman spaces*, Amer. J. Math. **110** (1988), 989–1054.

- [3] S. Axler, *The Bergman spaces, the Bloch space and commutators of multiplication operators*, Duke Math. J. **53** (1986), 315–332.
- [4] D. Bekolle, C. A. Berger, L. A. Coburn, and K. H. Zhu, *BMO in the Bergman metric on bounded symmetric domain*, J. Funct. Anal. **93** (1990), 310–350.
- [5] K. S. Choi, *Lipschitz type inequality in Weighted Bloch spaces \mathcal{B}_q* , J. Korean Math. Soc. **39** (2002), no. 2, 277–287.
- [6] ———, *Little Hankel operators on Weighted Bloch spaces in \mathbb{C}^n* , Commun. Korean Math. Soc. **18** (2003), no. 3, 469–479.
- [7] J. B. Conway, *A course in Functional Analysis*, Springer Verlag, New York (1985).
- [8] K. T. Hahn, *Holomorphic mappings of the hyperbolic space into the complex Euclidean space and Bloch theorem*, Canadian J. Math. **27** (1975), 446–458.
- [9] K. T. Hahn and E. H. Youssfi, *M-harmonic Besov p-spaces and Hankel operators in the Bergman space on the unit ball in \mathbb{C}^n* , Manuscripta Math **71** (1991), 67–81.
- [10] K. T. Hahn and K. S. Choi, *Weighted Bloch spaces in \mathbb{C}^n* , J. Korean Math. Soc. **35** (1998), no. 1, 177–189.
- [11] L. Hwa, *Harmonic analysis of functions of several complex variables in the classical domains*, Amer. Math. Soc. providence, R. I. **6** (1963).
- [12] W. Rudin, *Function theory in the unit ball of \mathbb{C}^n* , Springer Verlag, New York (1980).
- [13] R. M. Timoney, *Bloch functions of several variables*, J. Bull. London Math. Soc. **12** (1980), 241–267.
- [14] K. H. Zhu, *Duality and Hankel operators on the Bergman spaces of bounded symmetric domains*, J. Funct. Anal. **81** (1988), 260–278.
- [15] ———, *Multipliers of BMO in the Bergman metric with applications to Toeplitz operators*, J. Funct. Anal. **87** (1989), 31–50.
- [16] ———, *Operator theory in function spaces*, Marcel Dekker, New York (1990).

Department of Information Security
Konyang university
Nonsan 320-711, Korea
E-mail: ksc@konyang.ac.kr