

Mini Review

Seaweed Biotechnology and Biologically Active Substances

Yong-Ki Hong*

Department of Biotechnology and Bioengineering, Pukyong National University, Nam-gu, Busan 608-737, Korea

Abstract Seaweed biotechnology is a multidisciplinary subject to produce food, pharmaceuticals, chemicals, and environmental remediation materials from seaweed resources. It uses various techniques of cell culture, enzyme reaction and genetic manipulation to increase the production efficiency of useful seaweeds or their products. Firstly, an overview of key topics will be introduced in the fields of seaweed tissue culture, strain improvement, genetic analysis briefly as basic techniques. Secondly, some biologically active substances such as anti-inflammatory and antifouling substances that have been screened in my laboratory will be focused.

Key words : Antifouling, anti-inflammation, biologically active substance, seaweed biotechnology

Introduction

One quarter of the world's drugs come from natural sources, primarily from microorganisms and plants. As terrestrial resources become over explored, attention has turned to the marine environment as an alternative source of novel bioactive metabolites. Compounds isolated from marine organisms have shown potent anti-cancer, antiviral, antibacterial, anti-inflammatory or pain-killing activity [16]. Obstacle to the drug development of marine natural products is the lack of sufficient material for comprehensive pharmacological evaluation [8]. More often than not, compounds are isolated in mg quantities, and their structures are elucidated and published. For any pharmaceutical lead from a marine source, supply issues will always be a problem. Unless supply can be addressed in an economically feasible fashion, the dream of new effective drugs from the sea will falter. Thus, one of candidates for pharmaceutical source would be seaweed that is abundant or easily aquaculturable. The research in this laboratory is exploring abundant seaweeds that produce biologically active substances. In particular, our survey efforts in this area have led to two main themes in studies: (1) screening, isolation and biological analysis of biologically ac-

tive substances, such as phospholipase A₂ inhibitor, antifouling agent, and microalgal growth enhancer; (2) tissue culture, mutant selection, and genetic analysis to overcome supply issues of useful substances. These studies are highly interdisciplinary in nature and draw on diverse methodologies in marine biotechnology.

Basic biotechniques for seaweed

Tissue culture

For the tissue culture of seaweed, callus and blade formation depended on the gelling agents used under axenic culture conditions [9]. Procedures were developed for the axenic isolation of *Chlorella* and monospores from the red seaweed *Porphyra yezoensis* [3]. A spectrophotometric quantification method was optimized to evaluate its utility in seaweed tissue viability tests using the enzymatic reduction of colorless 2,3,5-triphenyltetrazolium chloride to a colored triphenylformazan [14].

Strain improvement

Putative transgenic *P. yezoensis* was obtained under the conditions optimized by the particle bombardment.

* Corresponding author

Phone: +82-51-620-6182, Fax: +82-51-620-6180

E-mail: ykhong@pknu.ac.kr

To make new varieties of *P. yezoensis*, monospores are the most useful cells in maintaining and culturing [13]. Making mutants for overproduction of essential amino acids are in progress.

Genetic analysis

A rapid and economical method of DNA and RNA extraction from seaweed was developed by the use of lithium chloride [6]. Complete sequence of the 18S rDNA was amplified and sequenced from species of the aquaculturable *Porphyra* in Korea. A pollutant (pine needle ash)-responding gene for glutaredoxin was isolated from the seaweed *P. yezoensis* using differential display technique [10].

Biologically active substances

Anti-inflammatory substance

A number of seaweed species are used as traditional medicine and health care belief as well as food in various regions of the world. Almost all Korean women, even immigrated to foreign countries [15], eat the brown seaweed *Undaria pinnatifida* (known as Miyok) soup after childbirth for a month because of the belief that it cleans the blood and increases milk production. Use of the *U. pinnatifida* for curing fever, urination, lump or swelling is recorded in an oriental medical textbook Donguibogam published in 1613 [4]. It is also known to contain ingredients that contract the uterus after childbirth [7]. For herbal medicine in China, it is used to treat urinary diseases and dropsy [17]. All of these symptoms are related to anti-inflammation reaction. Thus, to evaluate medicinal activities of the *U. pinnatifida*, we have measured anti-inflammatory activities of methanol extract against mouse ear edema, analgesic, antipyretic, and toxicity [11]. The methanol extract showed IC₅₀ value of 10.3 mg/mL against mouse ear edema induced by phorbol myristate acetate. Also in analgesic test, the extract showed inhibitory effect on acetic acid-induced writhing response with IC₅₀ of 0.48 g per kg-body weight. The extract elicited antipyretic action when tested in yeast-induced hyperthermia. Of the common 37 seaweed compared, *U. pinnatifida* showed the strongest suppression (Table 1).

The brown seaweed *Ishige okamurae* showed a potent inhibitory activity against bacterial phospholipase A₂ from *Vibrio mimicus* [5]. The purified compound inhibited the PLA₂ with IC₅₀ of 1.86 µg/mL and KI

Table 1. Biological activities in the methanol-soluble extract from seaweeds*

Species	1	2	3	4	5
CHLOROPHYTA					
<i>Codium fragile</i>	-	-	ND	-	-
<i>Enteromorpha compressa</i>	ND	ND	+	-	ND
<i>Enteromorpha linza</i>	-	-	+	+	-
<i>Monostroma nitidum</i>	-	-	ND	ND	-
<i>Ulva pertusa</i>	+	-	-	-	-
PHAEOPHYTA					
<i>Colpomenia bullosa</i>	-	+	ND	-	-
<i>Colpomenia sinuosa</i>	-	-	ND	+	-
<i>Costaria costata</i>	ND	+	ND	-	ND
<i>Ecklonia cava</i>	-	+	ND	-	-
<i>Endrachne binghamiae</i>	-	-	ND	ND	-
<i>Hizikia fusiformis</i>	-	-	ND	-	-
<i>Ishige okamurae</i>	ND	ND	ND	-	+
<i>Ishige sinicola</i>	+	+	ND	-	ND
<i>Kjellmaniella crassifolia</i>	-	-	ND	ND	-
<i>Sargassum confusum</i>	-	-	ND	-	-
<i>Sargassum horneri</i>	+	-	+	-	-
<i>Sargassum sagamianum</i>	-	-	ND	-	+
<i>Sargassum thunbergii</i>	-	-	ND	+	-
<i>Scytosiphon lomentaria</i>	-	+	ND	ND	-
<i>Undaria pinnatifida</i>	-	-	-	+	-
RHODOPHYTA					
<i>Carpopeltis affinis</i>	-	-	-	ND	-
<i>Carpopeltis cornea</i>	ND	ND	+	-	ND
<i>Chondrus ocellatus</i>	+	-	+	-	-
<i>Corallina pilulifera</i>	+	-	ND	-	-
<i>Gigartina intermedia</i>	-	-	ND	ND	-
<i>Gracilaria verrucosa</i>	ND	ND	-	+	ND
<i>Grateloupia prolongata</i>	-	-	ND	ND	-
<i>Grateloupia turuturu</i>	-	-	-	ND	-
<i>Hypnea charoides</i>	-	-	ND	-	-
<i>Pachymeniopsis elliptica</i>	+	-	+	+	-
<i>Porphyra yezoensis</i>	-	-	ND	-	-
<i>Symphycladia latiuscula</i>	-	-	ND	-	-

* (1) A “+” symbol indicates attachment inhibition of spores from the fouling seaweed *Enteromorpha prolifera* with 5 µL extract (200 µg) in 1 mL seawater. (2) A “+” symbol indicates inhibition of foot repulsive reaction of the mussel *Mytilus edulis* with 10 µL extract (40 µg). (3) A “+” symbol indicates inhibition of seaweed spore settlement with 1 µL extract (4 mg) from the crustose coralline *Lithophyllum yezoense* in 200 µL seawater. (4) A “+” symbol indicates anti-inflammatory activity with 10 µL (400 µg) against mouse ear edema induced by 0.2 µg phorbol myristate acetate. (5) A “+” symbol indicates inhibition of bacterial phospholipase A₂ with 15 µL (600 µg) in 3.1 mL enzyme reaction mixture. ND, not determined.

of 3.89 µg/mL competitively. It showed more than twice stronger than a standard rutin compound (Fig. 1). Topical application reduced mouse ear edema with IC₅₀ of 3.57 mg/mL (Table 1).




Fig. 1. Structure of 7-methoxy-9-methylhexadeca-4,8-dienoic acid (MMHDA) isolated from the brown seaweed *Ishige okamurae*. The compound strongly inhibits PLA₂ activity *in vitro* and has potent anti-inflammatory activity *in vivo*.

Antifouling substances

Three antifouling active compounds of L-pyrroglutamic acid (PGA), triethyl citrate (TEC), and di-*n*-octylphthalate (DNOP) were isolated from the brown seaweed *Ishige okamurae* (Fig. 2). Approximately 2.8 mg PGA, 1.7 mg TEC, and 2.0 mg DNOP were isolated from 600 g of *I. okamurae* powder. The concentrations of PGA, TEC, and DNOP required to cause foot repulsion in 50% of mussels (RD₅₀) were 9, 26, and 0.08 mM, respectively [1]. The PGA, TEC, and DNOP concentrations required to inhibit 50% attachment of algal spores (ID₅₀) were 24, 50, and 0.1 mM, respectively. These compounds showed stable antifouling activities against mussel and algal spore attachment (Table 1).

A study was made to investigate possible formation by the crustose coralline alga *Lithophyllum yessoense* of multiple allelopathic-related substances against the settlement and germination of spores of various seaweeds [12]. Spore settlement of 14 species was inhibited over 90% by solvent extracts and conditioned seawater. The germination of spores from 13 species was also inhibited by the extracts and conditioned seawater (Table 1). Isolation of the main active compound is in progress.

Microalgal growth enhancer

Cell growth of the marine microalga *Isochrysis galbana* was regulated by the addition of seaweed extracts in the culture medium [2]. Methanol-soluble extracts from 27 species of seaweed showed growth activation only from *Enteromorpha linza*, and growth inhibition from *Ishige foliacea* and *Sargassum sagamianum* (Table 1). Water-soluble extracts from *Grateloupia turuturu* and *Monostrom nitidum* showed growth activation, while none of the seaweed showed growth inhibition. From results of growth activation of extracts on *I. galbana*, the water extract of *M. nitidum* was the most effective up to two-fold increase in cell density with the addition of 1 mg mL⁻¹ of extract to the medium. The cell growth rate was increased from 0.52




Fig. 2. Structures of L-pyrroglutamic acid (A), triethyl citrate (B), di-*n*-octylphthalate (C) isolated from the brown seaweed *Ishige okamurae*. These compounds showed potent antifouling activities against mussel and algal spore attachment.

to 0.65 d⁻¹. Cell size, gross biochemical compositions, fatty acid compositions, and digestion efficiency by shellfish differed marginally between cultures of *I. galbana* grown with and without the *M. nitidum* aqueous extract. This extract has also enhanced the growth of other feed microalgae tested, including *Dunaliella salina*.

Acknowledgements

This research was supported by a grant (B-2004-01) from the Marine Bioprocess Research Center of the Marine Bio 21 Project funded by MOMAF, Korea.

References

1. Cho, J. Y., Choi, J. S., Kang, S. E., Kim, J. K., Shin, H. W. and Hong, Y. K. 2005. Isolation of antifouling active pyrroglutamic acid, triethyl citrate, and di-*n*-octylphthalate from the brown seaweed *Ishige okamurae*. *J. Appl. Phycol.* **17**, 431-435.
2. Cho, J. Y., Jin, H. J., Lim, H. J., Whyte, J. N. C. and Hong, Y. K. 1999. Growth activation of the microalga *Isochrysis galbana* by the aqueous extract of the seaweed *Monostroma nitidum*. *J. Appl. Phycol.* **10**, 561-567.
3. Choi, J. S., Cho, J. Y., Jin, L. G., Jin, H. J. and Hong, Y. K. 2002. Procedures for the axenic isolation of chonchocelis and monospores from the red seaweed *Porphyra*

- yezoensis*. *J. Appl. Phycol.* **14**, 115-121.
4. Donguibogam committee. 1999. Translated Donguibogam. Bubinmunwha Publisher, Seoul, pp 2198.
 5. Gyawali, Y. P., Choi, J. S., Khan, M. N. A., Ahn, S. H., Kong, I. S. and Hong, Y. K. 2005. Isolation of PLA₂ inhibitor from the brown seaweed *Ishige okamurae*. *Proc. Kor. Soc. Phycol.*, **2005**, 48.
 6. Hong, Y. K., Kim, S. D., Polne-Fuller, M. and Gibor, A. 1995. DNA extraction conditions from *Porphyra perforata* using LiCl. *J. Appl. Phycol.* **7**, 101-107.
 7. Huh, K., Song, J. W. and Choi, J. W. 1992. Studies on uterus contraction of the components of *Undaria pinnatifida*. *Kor. J. Pharmacogn.* **23**, 146-152.
 8. Jensen, P. R. and Fenical, W. 2000. Marine microorganisms and drug discovery: Current status and future potential. In: Fusetani N. (ed), *Drugs from the Sea*, Karger, Basel, pp 6-29.
 9. Jin, H. J., Seo, G. M., Cho, Y. C., Hwang, E. K., Sohn, C. H. and Hong, Y. K. 1997. Gelling agents for tissue culture of the seaweed *Hizikia fusiformis*. *J. Appl. Phycol.* **9**, 489-493.
 10. Jin, L. G., Choi, J. S., Jin, H. J., Shin, H. W. and Hong, Y. K. 2002. Isolation of pollutant (pine needle ash)-responding genes from the seaweed *Porphyra yezoensis* tissue. *Fish. Sci.* **68**, S1044-S1047.
 11. Khan, M. N. A., Gyawali, Y. P., Yoon, S. J., Choi, J. S., Kang, S. E. and Hong, Y. K. 2005. Anti-inflammatory activities of the methanol extract from the brown seaweed *Undaria pinnatifida*. *Proc. Kor. Soc. Phycol.* **2005**, 47.
 12. Kim, M. J., Choi, J. S., Kang, S. E., Cho, J. Y., Jin, H. J., Chun, B. S. and Hong, Y. K. 2004. Multiple allelopathic activity of the crustose coralline alga *Lithophyllum yessoense* against settlement and germination of seaweed spores. *J. Appl. Phycol.* **16**, 175-179.
 13. Mizuta, H., Yasui, H. and Saga, N. 2003. A simple method to mass produce monospores in the thallus of *Porphyra yezoensis* Ueda. *J. Appl. Phycol.* **15**, 345-349.
 14. Nam, B. H., Jin, H. J., Kim, S. K. and Hong, Y. K. 1998. Quantitative viability of seaweed tissues assessed with 2,3,5-triphenyltetrazolium chloride. *J. Appl. Phycol.* **10**, 31-36.
 15. Park, K. J. and Peterson, L. M. 1991. Beliefs, practices, and experiences of Korean women in relation to childbirth. *Health Care Women Int.* **12**, 261-269.
 16. Smit, A. J. 2004. Medicinal and pharmaceutical uses of seaweed natural products: A review. *J. Appl. Phycol.* **16**, 245-262.
 17. Tseng, C. K. and Chang, C. F. 1984. Chinese seaweeds in herbal medicine. *Hydrobiologia* **116/117**, 152-154.