

Role of MAP kinase on MMP-13 expression in rat periodontal ligament cells

Chan Gil Chung¹, De-Zhe Cui², Hyun-Ju Chung^{1,2}, Young Joon Kim^{1,2}

Department of Periodontology School of Dentistry¹, Dental Science Research Institute², Chonnam National University

I. Introduction

The pathways of periodontal tissue destruction during the course of destructive periodontal diseases have been fairly well described at the cellular level¹⁾. According to the current paradigm of infectious etiology, the events associated with tissue destruction could be most briefly summarized as follows: an external stimulus, usually toxins and antigens from bacteria of the dental biofilm, can exert direct effects on the periodontal tissues, but acts mainly by triggering a complex array of inflammatory and immune reactions that ultimately leads to the loss of supporting apparatus of affected teeth²⁾.

There is evidence supporting a direct participation of PDL cells on pathologic breakdown of extracellular matrix from both min-

eralized and non-mineralized periodontal tissues. Chang et al.³⁾ showed synthesis and secretion of matrix metalloproteinase 2 (MMP-2, type IV collagenase, gelatinase A) as well as indication of some synthesis of MMP-9 (also a type IV collagenase, gelatinase B) by PDL cells. Domeij et al.⁴⁾ also found similar results in gingival fibroblasts, demonstrating increase in secretion of both MMP-1 and MMP-3 by these cells after stimulation with IL-1 β , TNF- α and epidermal growth factor (EGF). By using specific pharmacological inhibitors, they also report indirect evidence of the involvement of both tyrosine kinase and p38 MAP kinase on the signaling pathways for the proinflammatory stimuli in these cells.

Collagenase 3 (or MMP-13) is a recently described protein in the MMP family and although type II collagen is the main sub-

* corresponding author: Young Joon Kim, Dept. of Periodontology, School of Dentistry, Chonnam National University, 5, Hak 1 dong, Dong-gu, Gwangju, 501-746, Korea. E-mail : youngjun@chonnam.ac.kr

strate target for this enzyme, it also degrades collagen types I, III and V, which are all major components of cartilage and bone⁵⁾. In periodontal inflammation, several resident cell types have been shown to immunoreactive to MMP-13, including fibroblasts and macrophages. MMP-13 has been shown to exist at higher levels in periodontal infected tissues than in healthy control tissues⁶⁾. In peri-implantitis models of bone loss, implants having >3 mm of bone loss had higher levels of MMP-13 from the periimplant fluid at these sites⁷⁾. Thus, MMP-13 expression may be related not only to collagen degradation, but also directly or indirectly to bone resorption.

It was shown recently that proinflammatory TNF- α increases MMP-13 mRNA expression in human PDL cells, with the maximal effect observed with a concentration of 10 ng/ml⁸⁾.

The study of the signal transduction pathway leading to expression of MMP-13 mRNA may give further insight into the possible cross-talk by proteins involved in signal transduction, which may also activate other genes related to bone resorption.

Proinflammatory cytokines stimulate multiple intracellular signaling pathways, including the p38 MAP kinase pathways, NF- κ B pathway, extracellular regulated kinase (ERK), and stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK)^{9,10)}. Although much attention has focused on the NF- κ B pathway of cytokine gene activation, relatively less attention has been spent on p38 MAP kinase which is also involved in

inflammation and environmental stress induced signaling^{11, 12)}. The p38 kinases constitute a distinct MAP Kinase subfamily that plays a role in adaptation, homeostasis, and stress responses. In recent years, the MAP kinase system has been shown to mediate, in part, the actions of proinflammatory cytokines as well as acting a principal mediator of the inflammatory response to LPS in osteoblastic cells¹³⁾. The role of p38 MAP Kinase in MMP-13 gene regulation is becoming more evident¹⁴⁾, although the mechanisms of gene regulation are not entirely clear.

PDL cells play a potential central role in tissue destruction associated with periodontal disease¹⁵⁾. The relative lack of data related to signal transduction pathways of both inflammatory and infectious external signals in these cells, and the relevance of metalloproteinases, specifically of MMP-13.

This study was accomplished as an initial step towards a better understanding of the role of p38 MAP Kinase on the expression of MMP-13 secreted by PDL cells. Thus, the purpose of this study is to investigate the role of p38 MAP Kinase on MMP-13 mRNA expression through RT-PCR and western blot assays.

II. Material and Method

1. Cell culture

Rat periodontal ligament fibroblasts (PDL) obtained as described by Matsuda et al.¹⁶⁾ were cultured in DMEM (Invitrogen, U.S.A) medium, supplemented with 10%

heat-inactivated fetal bovine serum (FBS; Invitrogen, U.S.A), 100 U/ml penicillin and 100 U/ml streptomycin (Penicillin-Streptomycin, GibcoBRL, U.S.A), at 37°C in a humidified atmosphere of 5% CO₂ in air. Cells were routinely deinduced in culture media without FBS for 6-8 hours prior to stimulation with proinflammatory cytokines.

2. Reverse transcription-polymerase chain reaction (RT-PCR) analysis

Cells were grown to near confluency in 6-well dishes, deinduced and then treated with the following proinflammatory and other substances: recombinant human interleukin 1 β (IL-1 β ; 1 nM), recombinant human tumour necrosis factor- α (TNF- α 10 nM), recombinant human bone morphogenic protein-7 (BMP-7, 100 ng/ml), parathyroid hormone (PTH, 10 nM). A negative control represented by the vehicle used to dilute these substances also were included.

After the treatment periods, total RNA were collected with Trizol[®] reagent (Invitrogen, U.S.A), according to the instructions provided by the manufacturer. Briefly, approximately 1 ml of Trizol[®] was added to each well to cell lysis and this cell lysate was homogenized by passing several times through a pipette. This solution was transferred to eppendorf tubes, incubated at room temperature (RT) for 15-30 min, and then 0.2 ml of chloroform per ml of Trizol[®] was added. After vigorously shaking the tubes and centrifugation at 10,000 xg for 15 min at 2-8°C, the aqueous phase was collected and transferred to a fresh tube. Total RNA was pre-

cipitated by adding isopropyl alcohol, washed in 75% ethanol and redissolved in RNase-free water for storage at -70°C until use.

Using spectrophotometry, 5 μ g of RNA was quantitated (read at an absorbance of 260 nm) and used for cDNA synthesis with oligo (dT) 12-18 primer by reverse transcriptase(RT) reactions (RT-PCR primers: MMP-13 328bp S5-GACTTCACGATGGCATTGCTG-3, A5-GCATCAACCTGCTGAGGATGC-3 GAPDH 418bp S5-ATTCCATGGCACCGTCAAGGCT-3, A5-TCAGGTCCACCACTGACACGTT-3).

Approximately 2 μ l of RT product was used as a template for the polymerase chain reaction (PCR).

The PCR products were checked by 2% agarose gel electrophoresis, stained by ethidium bromide. Semi-quantitative PCR was performed. Semi-quantitative differences of MMP-13 expression were normalized by GAPDH.

3. mRNA stability Experiments

To determine the relative role of p38 MAP Kinase on IL-1 β -induced MMP-13 mRNA stability, the rate of MMP-13 mRNA decay was determined in the presence and absence of the p38 inhibitor, SB203580 (10 μ M). PDL cells were pre-treated with SB203580 (10 μ M) for 60 minutes, then treated with IL-1 β (1 nM) for 16-18 hours. Subsequently, actinomycin D (2 μ g/ml), DNA polymerase inhibitor, was added to prevent further transcription. Total RNA was harvested and analyzed by RT-PCR as described above after 0, 60, 120, 240 minutes treated with actinomycin D.

4. Western blot analysis

To analyze the cell signaling pathways associated with the different stimuli, the cells were treated with the same substances (IL-1 β , TNF- α , BMP-7 and PTH) for shorter time periods. For this experiment, PDL cells were grown to near confluency in 6-well dishes and de-induced as described previously.

After treatment, whole cell lysates were collected for Western blotting analysis of both active (phosphorylated) and inactive (non-phosphorylated) MAP kinase. Western blotting was conducted as described in the literature. Briefly, after removal of media by aspiration, cells were washed with ice-cold PBS (pH 7.4), and 100 μ l of SDS-sample buffer was added to each well with removal of cells by scraping the wells with a disposable cell scraper. The cell lysate was transferred to eppendorf tubes, boiled at 100°C for 5 min, placed on ice, sonicated for 10-15 seconds (three times) and stored at -20°C until use.

Following sample preparation (by boiling at 100°C for 5 min and placing on ice), SDS-PAGE was performed using pre-stained molecular weight markers to verify electro-transfer and biotinylated protein markers to determine molecular weights. After running the gel, proteins were blotted to a nitrocellulose membrane.

The immunoblotting protocol was performed as described by the supplier of the commercial antibody kits. The nitrocellulose membrane was blocked for non-specific binding, washed and incubated with the primary mouse anti-human antibodies (for MAP kin-

ases: Cell Signaling, U.S.A) using gentle agitation overnight at 4°C. After three washes, the membrane was incubated with the secondary anti-mouse antibodies conjugated to Horseradish peroxidase (Cell Signaling, U.S.A) and anti-biotin HRP-conjugated antibody (to detect biotinylated protein markers) with gentle agitation for 1 hour at room temperature. After three consecutive washes, the membrane was incubated with the enzyme substrate for 1 minute, wrapped in plastic film and exposed to X-ray film (2 mins).

III. Results

1. MMP-13 mRNA was expressed by proinflammatory cytokines and PTH

In semi-quantitative RT-PCR analysis, MMP-13 was induced to different extents by IL-1 β , TNF α , rh BMP-7, and PTH. When levels of GAPDH were used to normalize the data, IL-1 β , TNF α and PTH-induced MMP-13 expression was increase to 320%, 180%, 380%, respectively. But BMP-7 had little effect on MMP-13 expression (120%) (Figure 1).

2. p38 was required for IL- β -induced MMP-13 mRNA induction

MMP-13 mRNA gene expression was determined by RT-PCR. The result was shown that p38 is necessary dor IL-1 β -induced MMP-13 mRNA synthesis. When PDL cells were pre-treated with SB203580, the p38 MAPK inhibitor, and then stimulated with





Figure 1. Semiquantitative RT-PCR of rat periodontal ligament cells showing steady state MMP-13 mRNA levels. (lane 1:DNA ladder, 2: control, 3: IL-1 β (1 nM), 4: TNF α (10 nM), 5: BMP-7(50 ng/ml), 6: PTH(10 nM))

IL-1 β , the result was shown to approximately 40% reduction in IL-1 β -induced MMP-13 mRNA levels compared to IL-1 β treated cells only (from 3.5 fold to 2.0 fold). There was no effect on PTH pre-treated cells (Figure 2).

3. p38 was needed to mediate IL-1 β -induced MMP-13 mRNA stability.

To determine if p38 MAP Kinase regulated IL-1 β induced MMP-13 stability, we used Actinomycin D. The results indicate



Time(hr)	0	1	2	4	0	1	2	4
IL-1 β	+	+	+	+	+	+	+	+
SB	-	-	-	-	+	+	+	+

Figure 2. Semiquantitative RT-PCR of rat periodontal ligament cells showing steady state MMP-13 mRNA levels. (lane 1:DNA ladder, 2: control, 3: IL-1 β (1 nM), 4: PTH, 5: SB203580(10mM) + IL-1, 6: SB203580(10mM) + PTH)




Figure 3. MMP-13 mRNA stability experiment using Actinomycin D in PDL cells. P38 MAP kinase mediates IL-1 β -induced MMP-13 stability. * indicates half lives of MMP-13 mRNA in the presence of SB203580.

that IL-1 β increased the half-life of MMP-13, with 120 minutes, but in the presence of SB203580, the half life was decreased to 60 minutes following normalization to GAPDH (Figure 3).

4. p38 MAP Kinase and JNK were activated by IL-1 β

Western blot analysis was utilized to examine whether IL-1 β , TNF α , BMP-7 and PTH activated MAP Kinase. Western blot analysis of antibodies indicated that IL-1 β




Figure 4. Western blot analysis of phosphorylated and unphosphorylated p38 MAP Kinase. (lane 1:MW, 2: control, 3: IL-1 β (1 nM), 4: TNF- α (10 nM), 5: BMP-7 (50 ng/ml), 6: PTH (10 nM))




Figure 5. Western blot analysis of phosphorylated and unphosphorylated JNK. JNK is activated by IL-1 β . (lane 1:MW, 2: control, 3: IL-1 β (1 nM), 4: TNF- α (10 nM), 5: BMP-7 (50ng/ml), 6: PTH (10 nM))

activates both phosphorylated form of p38 MAP Kinase and JNK MAP Kinase, but not ERK MAP Kinase. Neither BMP-7 nor PTH activates p38, JNK, and ERK MAP Kinase (Figure 4~6).

5. Activation of p38 MAP Kinase was inhibited with treatment with SB203580

Western blot analysis of antibodies indicated that anti-phospho p38 and non-phospho p38 MAP Kinase are activated by IL-1 β in PDL cells. Importantly, the p38 in-




Figure 6. Western blotanalysis of phosphorylated and unphosphorylated ERK. (lane 1:MW, 2: control, 3: IL-1 β (1 nM), 4: TNF- α (10 nM), 5: BMP-7 (50 ng/ml), 6: PTH (10 nM))




Figure 7. Western blot analysis of phosphorylated and unphosphorylated p38 MAP Kinase. p38 MAP Kinase is activated by IL-1 β and subsequent p38 inhibition by treatment with SB203580. (lane 1:MW, 2: control, 3: IL-1 β (1 nM), 4: SB203580 (10 mM) only, 5: SB203580 (10 mM)+ IL-1)

hibitor, SB203580 was able to block IL-1 β -induced p38 MAP Kinase (Figure 7).

IV. Discussion

Proinflammatory cytokine production in periodontal disease, as well as other chronic inflammatory bone diseases, such as rheumatoid arthritis, results in a net bone loss that ultimately negatively affects host function¹⁷⁾. Complex cytokine networking and cellular cross-talk dictate cellular responses involved in bone remodeling. Several media-

tors of bone resorption and remodeling have been identified in response to proinflammatory cytokines, such as IL-1, lipopolysaccharide, as well as other bone-resorptive agents¹⁸⁾.

In fact, there is evidence supporting a direct participation of PDL cells on pathologic breakdown of extracellular matrix from both mineralized and non-mineralized periodontal tissues^{15, 19)}. In most systems, collagenase expression is not constitutive but is induced by variety of agents, including cytokines, tumour promoters, and bone-resorbing agents. Bone resorptive agents, IL-1, IL-6, PTH, and TNF- α enhance the expression of collagenase 3 in mouse calvaria, primary rat osteoblastic cells, and other osteoblastic model systems^{14, 20, 21)}.

Several studies showed that the synthesis of MMP-13 by rat osteoblasts is regulated by systemic hormones and by cytokines. But there are few reports that the expression of MMP-13 by various proinflammatory cytokines in PDL cells. Thus we examined the effects of various stimuli, including proinflammatory cytokines such as IL-1 β , TNF- α , and BMP-7, and PTH on MMP-13 in PDL cells. In this study, MMP-13 steady state gene expression in PDL cells was determined by semi-quantitative RT-PCR. MMP-13 was induced to different extents by IL-1 β , TNF- α , BMP-7 and PTH. IL-1 β and PTH stimulated MMP-13 in PDL cells. However, in contrast to IL-1 β and PTH, BMP-7-induced MMP-13 expression were weak. This result is consistent with results obtained other investigators^{20, 22, 23)} where PTH enhanced expression of MMP-13 but

bone morphogenic proteins suppressed expression of MMP-13 in rat osteoblasts. In the present study, TNF- α also induced expression of MMP-13. This result is consistent with Kirkwood's report²⁰⁾ that TNF- α -induced MMP-13 expression in rat osteoblasts was more enhanced than untreated control.

The signaling cascade from the external membrane to the nucleus is extremely complex and can involve a great number of different proteins, depending on the external factor being considered. Nevertheless, some components of the signaling cascade have been extensively studied, such as the cytoplasmic protein kinases collectively known as the mitogen-activated protein (MAP) kinases, which up to now includes the following kinases: SAPK/JNK, ERK1/2 and p38¹³⁾.

In recent years, the MAP Kinase system has been shown to mediate, in part, the actions of pro-inflammatory cytokines, although the mechanisms of gene regulation are not entirely clear. Moreover, there are many questions unanswered regarding the role for specific MAP Kinase involvement in bone metabolism. No studies to date have directly addressed the role of p38 MAP Kinase in periodontal disease models. Understanding p38 MAP kinase-mediated signaling pathways is essential to aid in generating therapeutics capable of treating diseases mediated by MMP-13 and other mediators. In this study, through semi-quantitative RT-PCR in PDL cells, IL-1 β -induced MMP-13 expression was reduced in the presence of specific p38 MAP Kinase inhibitor, SB203580 by 50%. However in con-

trast IL-1 β , PTH-induced MMP-13 repression was not observed in the presence of SB203580. This result showed that p38 MAP Kinase was involved in IL-1 β -induced MMP-13 steady state mRNA levels. From this result, although a number of cytokines and hormones can induce MMP-13 expression, they do not always activate the same signaling events that mediate these effects. For example, PTH-induced MMP-13 expression requires activation of protein kinase C and protein kinase A pathway. In contrast, MMP-13 induction by IL-1 required nuclear factor (NF) κ B translocation, p38 MAP kinase activity in chondrocytes²⁴⁾. Thus different signaling pathways may be activated in MMP-13 expression. This result also showed that specific inhibition of p38 MAP Kinase prevents MMP-13 induction by IL-1 β , indicating the importance of this pathway in MMP-13 gene expression in PDL cells.

In figure 3 to 5, it was shown that IL-1 β only induced phosphorylation of p38 MAP Kinase and c-Jun N-terminal Kinase (JNK) in PDL cells by Western blot analysis. It implicates that IL-1 β -induced MMP-13 requires p38 MAP Kinase activity and JNK activity. This result is consistent with Mengshol's study²⁴⁾ that IL-1-induced-MMP-13 expression required NF- κ B, p38 and JNK activation in chondrocytes. In this study, TNF- α was not activate p38 MAP kinase nor ERK. According to Johansson et al²⁵⁾, TNF- α stimulation of MMP-13 expression in keratinocytes led to activation of ERK, JNK, p38 activation, but there is also a report²⁶⁾ that ERK activation did not mediate MMP-13 ex-

pression in fibroblast-like synoviocytes. A possible explanation of different results may be different intracellular mechanism which affect MMP-13 expression. Thus, although a number of cytokines and hormones can induce MMP-13 expression, they do not always activate the same signaling events in different cell lines.

Through further Western blot analysis, it was shown that SB203580 inhibited the ability of IL-1 β to induce p38 phosphorylation in PDL cells. Specific inhibition of p38 MAP kinase prevents MMP-13 induction by IL-1 β , indicating the importance of this pathway in MMP-13 gene expression.

The mechanism of p38 regulation of MMP-13 mRNA expression could occur at the transcriptional or post-transcriptional level. To determine whether p38 stabilized MMP-13 mRNA by post-transcriptional regulation, mRNA degradation studies were performed. SB203580 altered IL-1 β -induced MMP-13 mRNA decay rates in PDL cells. It means that in PDL cells, p38 MAP kinase appears to mediate its effects on enhancing MMP-13 gene expression mainly through the stabilization of MMP-13 mRNA. As shown in Figure 3, p38 MAP kinase mediates IL-1 β -induced MMP-13 mRNA stability. But SB203580 reduces the half life of the MMP-13 from approximately 120 minutes to less than 60 minutes in IL-1 β -stimulated cells in the presence of Actinomycin D, an inhibitor of DNA-primed RNA polymerase, to arrest transcription.

This study demonstrates that activation of MMP-13 gene expression is induced by IL-1 β , TNF- α and PTH in PDL cells, but only

IL-1 β -induced MMP-13 activation through p38 MAP kinase. It implicates that p38 MAP kinase pathway is a central signal pathway which controls IL-1 β -induced MMP-13 expression.

Further studies will be needed to elucidate that MAP Kinases pathway would be a central signalling pathway controlling IL-1 β -induced MMP-13 expression at the gene promoter level. Understanding signalling mechanism which contributes towards MMP-13 expression may provide therapeutic targets in the management of inflammatory bone diseases such as periodontitis and rheumatoid arthritis

V. Conclusion

Matrix metalloproteinases (MMPs) play an important role both in the maintenance and degradation of extracellular matrix within the periodontium. MMP-13 has been shown to positively correlate with periodontal disease. Mitogen activated protein (MAP) Kinases play an important role in the induction of pro-inflammatory cytokines such as IL-1 β and TNF- α . This study is accomplished as an initial step towards a better understanding of the role of MMP-13 in the pathogenesis of periodontal disease and possible regulation of MMP-13 through MAP Kinase signaling by different stimuli. MMP-13 mRNA expression in PDL cells was determined by semi-quantitative RT-PCR and MAP Kinases expression were determined by Western blot.

The results are as follows;

1. In semi-quantitative RT-PCR analysis, IL-1 β , TNF- α , and PTH-induced MMP-13 expression was increase to 320%, 180%, 380%, respectively. But rh BMP-7 had little effect on MMP-13 expression (120%).
2. When PDL cells were pre-treated with SB203580 (10 μ M), the p38 inhibitor, IL-1 β -induced MMP-13 mRNA level was reduced by 40% in compared to IL-1 β -treated cells only. There was no effect on PTH pre-treated cells.
3. IL-1 β increased the half-life of MMP-13 mRNA with 120 minutes, but in the presence of SB203580, the half life was decreased to 60 minutes.
4. In western blot analysis, IL-1 β activates both phosphorylated form of p38 MAP Kinase and JNK, but not of ERK. Neither BMP-7 nor PTH activates p38, JNK, and ERK MAP Kinase. IL-1 β -induced p38 MAP Kinase phosphorylation is inhibited by SB203580.

This results suggest that p38 MAP Kinase pathway would be a central signal pathway which controls IL-1 β -induced MMP-13 expression.

VI. References

1. Page RC. and Schroeder HE. Pathogenesis of inflammatory periodontal disease. A summary of current work. Lab. Invest 1976;34:235-249.
2. Dennison DK. and Van Dyke TE. The acute inflammatory response and the role of phagocytic cells in periodontal

- health and disease. *Periodontol.* 2000; 14:54-78.
3. Chang Y-C, Yang S-F, Lai C-C, Liu J-Y, Hsieh YS. Regulation of matrix metalloproteinase production by cytokines, pharmacological agents and periodontal pathogens in human periodontal ligament fibroblast cultures. *J. Periodont. Res.* 2002 ;37:96-203.
 4. Domeij H, Yucel-Lindberg T, Moder T. Signal pathways involved in the production of MMP-1 and MMP-3 in human gingival fibroblasts. *Eur J Oral Sci.* 2002;110: 302-306.
 5. Freije J M P, Diez-Itza I, Balbin M et al. Molecular cloning and expression of collagenase-3, a novel human matrix metalloproteinase produced by breast carcinomas. *J Biol Chem.* 1994;269: 16766-16773.
 6. Tervahartiala T, Pirila E, Ceponis A et al. The in vivo expression of the collagenolytic matrix metalloproteinases (MMP-2, -8, -13, and -14) and matrilysin (MMP-7) in adult and localized juvenile periodontitis. *J Dent Res.* 2000;79:1969-1977.
 7. Ma J, Kitti U, Teronen O, Sorsa T et al. Collagenases in different categories of peri-implant vertical bone loss. *J Dent Res.* 2000;79:1870-1873.
 8. Nishikawa M, Yamaguchi Y, Yoshitake K, Saeki Y. Effects of TNFalpha and prostaglandin E2 on the expression of MMPs in human periodontal ligament fibroblasts. *J Periodontal Res.* 2002;37: 167-76.
 9. Gonzalez G.A. and Montminy M.R. Cyclic AMP stimulates somatostatin gene transcription by phosphorylation of CREB at serine 133. *Cell* 1989;59:675-680.
 10. Haskill S et al. Characterization of an immediate-early gene induced in adherent monocytes that encodes I kappa B-like activity. *Cell* 1991;65:1281-1289.
 11. Han Z, Boyle DL, Chang L et al. c-Jun N-terminal kinase is required for metalloproteinase expression and joint destruction in inflammatory arthritis. *J Clin Invest.* 2001;108:73-81.
 12. Patil C, Zhu X, Rossa C Jr, Kim YJ, Kirkwood KL. p38 MAPK regulates IL-1beta induced IL-6 expression through mRNA stability in osteoblasts. *Immunol Invest.* 2004;33:213-233.
 13. Seger R. and Krebs EG. The MAPK signaling cascade. *FASEB* 1995;9:726-735.
 14. Lee HS, Miao LH, Chen CH et al. Differential role of p38 in IL-1alpha induction of MMP-9 and MMP-13 in an established liver myofibroblast cell line. *J Biomed Sci.* 2003;10:757-765.
 15. Ejeil AL, Gaultier F, Igondjo-Tchen S et al. Are cytokines linked to collagen breakdown during periodontal disease progression? *J Periodontol* 2003;74:196-201.
 16. Matsuda N, Kumar M, Ramakrishnan PR et al. Evidence for up-regulation of epidermal growth factors on rat periodontal ligament fibroblastic cells associated with stabilization of phenotype in vitro. *Arch Oral Biol.* 1993;38: 559-569.
 17. Dayer J.M. and H. Fenner. The role of cytokines and their inhibitors in arthritis. *Baillieres Clin Rheumatol.* 1992;6:485-516.
 18. Horwood N.J et al. Osteotropic agents regulate the expression of osteoclast differentiation factor and osteoprotegerin

- in osteoblastic stromal cells. *Endocrinology* 1998;139:4743-6.
19. Lekic P. and McCulloch CA. Periodontal ligament cell population: the central role of fibroblasts in creating a unique tissue. *Anat Rec.* 1996;245:327-341.
 20. Kirkwood K, Martin T, Agnello K, Kim YJ. Differential regulation of MMP-13 by chemical modified tetracyclines in osteoblasts. *J Int Acad Periodontol.* 2004;6:39-46.
 21. Shlopov BV, Stuart JM, Gumanovskaya ML, Hasty KA. Regulation of cartilage collagenase by doxycycline. *J Rheumatol.* 2001;28:835-42.
 22. Canalis E, Rydziel S, Delany A, Varghese S, Jeffrey J. Insulin-like growth factors inhibit interstitial collagenase synthesis in bone cell cultures. *Endocrinology* 1995;136: 1348-1354.
 23. Gazzo E, Rydziel S, Canalis E. Skeletal bone morphogenetic proteins suppress the expression of collagenase-3 by rat osteoblasts. *Endocrinology.* 1999;140: 562-567.
 24. Mengshol JA, Vincenti MP, Brinckerhoff CE. IL-1 induces collagenase-3 (MMP-13) promoter activity in stably transfected chondrocytic cells: requirement for Runx-2 and activation by p38 MAPK and JNK pathways. *Nucleic Acids Res.* 2001;29: 4361-4372.
 25. Johansson N, Ala-aho R, Uitto V et al. Expression of collagenase-3 (MMP-13) and collagenase-1 (MMP-1) by transformed keratinocytes is dependent on the activity of p38 mitogen-activated protein kinase. *J Cell Sci.* 2000;113:227-235.
 26. Pillinger MH, Rosenthal PB, Tolani SN et al. Cyclooxygenase-2-derived E prostaglandins down-regulate matrix metalloproteinase-1 expression in fibroblast-like synoviocytes via inhibition of extracellular signal-regulated kinase activation. *J Immunol.* 2003;171:6080-6089

백서 치주인대세포에서의 MMP-13 mRNA에 대한 MAP kinase의 역할

정찬길¹, 최득철², 정현주^{1,2}, 김영준^{1,2}

전남대학교 치의학전문대학원 치주과학교실¹ 및 치의학연구소²

Matrix metalloproteinases (MMPs)는 치주조직내에 존재하는 세포외기질의 유지와 분해에 중요한 역할을 담당하고 있으며 이중 MMP-13은 치주질환의 진행과 깊은 관계가 있다고 알려져 있다.

이번 연구는 치주질환의 진행에 있어서 MMP-13의 활성화에 대한 mitogen activated protein(MAP) Kinase의 역할을 구명하기 위해 시행되었다. 백서 치주인대세포에서의 MMP-13 mRNA의 발현은 RT-PCR에 의하여, 그리고 MAP Kinase의 발현은 Western blot에 의하여 측정하였다.

Interleukin-1 β (IL -1 β), Tumor necrosis factor α (TNF- α)와 parathyroid hormon(PTH)는 MMP-13 mRNA 발현을 각각 320%, 180%, 380% 증가시켰으나 bone morphogenetic protein-7(BMP-7)은 MMP-13 mRNA의 발현을 증가시키지 않았다. p38 MAP Kinase 억제제인 SB203580은 IL -1 β 유도 MMP-13의 발현을 약 40%정도 억제시켰으나, PTH-유도 MMP-13 mRNA의 발현은 억제하지 못했다. IL -1 β 는 MMP-13 mRNA의 반감기를 약 2시간 정도로 증가시켰으나, p38 MAP Kinase 억제제로 전처리한 경우에는 반감기가 60분으로 줄어들었다. IL-1 β 는 p38 MAP kinase와 JNK의 인산화 활성을 증가시켰으나 PTH, TNF- α 와 BMP-7은 p38, JNK, ERK의 활성을 증가시키지 못했다.

이상의 연구결과는 p38 MAP Kinase가 백서 치주인대세포에서의 MMP-13 mRNA 발현을 조절하는데 중요한 역할을 담당함을 시사하였다.