

Notes

Synthesis and Metal Binding Properties of Anthraquinone Bridged with Coumarin

Byung Ju Ryu, Eun Jin Cho, and Kye Chun Nam*

Department of Chemistry, Chonnam National University, Gwangju 500-757, Korea. *E-mail: kenam@chonnam.ac.kr


Received June 21, 2007

Key Words : Anthraquinone, Coumarin, Metal binding

Molecular switching in macrocyclic polyether systems has been the subject of intense study during the past 30 years. Especially, photochemical switching has been well-studied by Shinkai, Ueno, Takagi, Tabushi, and others.¹ In addition, changes in pH,² light³ and oxidation-reduction chemistry⁴ have all been explored as switching mechanisms.

The lariat ethers,⁵ podands having nitroaromatic sidearms⁶ and similar species derived from anthraquinones have been concerned with electrochemical switching. Especially the anthraquinone system is an interesting one for the study of reducible and switchable lariat ethers⁷ because its anion is stable for months in water so long as oxygen is excluded.⁸ Although podands (noncyclic polyethylene glycols) generally exhibit poor cation binding, when combined with the semiquinone nucleus, very strong cation complexation is observed for the radical anion.^{5,9} The anthraquinone podands can be used for electrochemically switched cation transport in bulk organic membranes^{7b} and both monoanion and dianion species can be formed and detected.¹⁰ Both one-electron and two-electron reduction can lead to enhanced cation binding in anthraquinone-derived podands and lariat ethers.^{7a} Anthraquinones differ significantly from the nitroaromatic systems in their ability to undergo discrete one or two electron reduction and also anthraquinone substituted podands exhibit surprising geometrical effects during the electrochemical switching process.^{7b} Gokel and co-workers¹⁰ have reported two armed podands based on the anthraquinone ring systems, which showed the cation binding enhancement by cyclic voltammetry. 1,8-Anthraquinone derived crown ether having a total of six oxygen donors in the macrocycling is an exceptionally strong cation binder not only when reduced, but in its neutral form as well.

Anthraquinone plays an important role in the various photochemical and colorimetric sensor systems. In addition, coumarins¹¹ are interesting chromophores by their photochemical and photophysical properties, and have been used to convert crown ethers into fluorescent probes for alkali metal ions. Herein we synthesized a new cation receptor **4** which combined coumarin and anthraquinone with ether units, and studied the binding properties of cation guests.

Scheme 1. Synthesis of receptor **4**.

Results and Discussion


The receptor **4** was prepared by the reaction of 1,8-dihydroxyanthraquinone and 6,7-bis[2-(2-iodoethoxy)ethyl]oxy-4-methylcoumarin **3** in the presence of potassium carbonate in 42% yield. Iodo derivative **3** was obtained from **2**, which was synthesized from the reaction of **1** and toluene-4-sulfonic acid 2-(2-chloroethoxy)ethyl ester, in order to increase the reactivity by treating with NaI.

The ¹H NMR spectrum of **4** shows two doublets and a triplet at δ 7.85, 7.62 and δ 7.28 for six anthraquinone protons and three singlets at δ 7.09, 6.84 and 6.13 for three coumarin protons. Also two multiplets at δ 4.30–3.90 for sixteen ether protons and a singlet at δ 2.34 for methyl protons are observed.

The alkali metal binding properties were investigated from two phase extraction experiment where aqueous solutions of the picrate salts are shaken with methylene chloride solution of ligand. The amount of metal ion was determined from

Table 1. Percentage extraction^a (%E) of alkali picrates from water into CH₂Cl₂

Ligand	% Extraction				
	Li ⁺	Na ⁺	K ⁺	Rb ⁺	Cs ⁺
4	0	0	13	16	15



^a 1×10^{-3} M receptor solution in CH₂Cl₂ was shaken (15 hrs.) with 1×10^{-1} M picrate salt solution in H₂O.




Figure 1. ¹H NMR spectra of **4** (a) without cation (b) with 10 eq. of LiClO₄ (c) with 10 eq. of NaClO₄ (d) with 10 eq. of KClO₄ (e) with 10 eq. of RbClO₄ (f) with 10 eq. of CsClO₄ in CD₃CN/DMSO-*d*₆ (v/v = 3:2)

picrate absorption in the UV spectrum. Table 1 showed the percent extraction of metal picrate salts by ligand **4**. The ligand **4** extracted 16% of the rubidium picrate from aqueous solution into the methylene chloride solution of the host, and also 13% of the potassium and 15% cesium were extracted. However, no extraction of lithium and sodium ions were observed.

The cation binding properties of **4** were examined by ¹H NMR experiments in CD₃CN/DMSO-*d*₆ (v/v = 3:2). The addition of 10 equivalent of KClO₄, RbClO₄ and CsClO₄ caused a small downfield shift of all peaks. However, the other alkali ions such as LiClO₄ and NaClO₄ were not changed as shown in Figure 1.

A clear selectivity between relatively large cations such as potassium, rubidium and cesium ions was not observed. Even though a binding site of receptor **4** can be an equivalent of 22-crown-7 which could prefer the binding of the larger cation such as cesium, a rigid anthraquinone moiety might prevent the adjustment for the large cation. But as expected from the extraction study, no changes of chemical shifts were observed in the presence of lithium and sodium cations, indicating that sodium and lithium cations are too small for the effective binding with **4**. Therefore, a clear binding selectivity between relatively large cations such as potassium, rubidium and cesium ions and small cations such as sodium and lithium were observed. Optical properties of **4** in the presence of alkali cations were investigated, but any noticeable changes were not observed.

Experimental

6,7-Bis[2-(2-chloroethoxy)ethyl]loxy-4-methylcoumarin (2). To a stirred solution of 0.5 g (2.6 mmol) of 6,7-dihydroxy-4-methylcoumarin **1** and 2.16 g (15.6 mmol) of K₂CO₃ in 100 ml. of CH₃CN, 1.45 g (5.2 mmol) of toluene-4-sulfonic acid 2-(2-chloro-ethoxy)-ethyl ester prepared by the reaction of 2-(2-chloroethoxy)ethanol with *p*-toluene-sulfonyl chloride was added and the reaction mixture was refluxed for 8 h. The solvent was removed and residue was taken up in CHCl₃ (100 ml.) and washed with 0.1 N HCl (100 ml.) and water. The organic phase was separated and dried with MgSO₄ and the crude products was purified by column chromatography (eluent, CHCl₃:*n*-Hexane:Ethyl acetate = 2:1:1) to yield 0.7 g (67%) of white powder **2**. ¹H NMR (CDCl₃) δ 7.09 (s, 1H, ArH), 6.84 (s, 1H, ArH), 6.16 (d, 1H, ArH, *J* = 1.1 Hz), 4.25-3.67 (m, 16H, -CH₂CH₂-), 2.39 (d, 3H, CH₃, *J* = 1.1 Hz).

6,7-Bis[2-(2-iodoethoxy)ethyl]loxy-4-methylcoumarin (3). To a stirred solution of 0.30 g (0.74 mmol) of 6,7-Bis[2-(2-iodoethoxy)ethyl]loxy-4-methylcoumarin and 1.10 g (7.34 mmol) of NaI in 30 mL of CH₃CN was added and the reaction mixture was stirred for 2 days. The solvent was removed and residue was taken up in CHCl₃ (100 ml.) and washed with water. The organic phase was separated and dried with MgSO₄ and evaporated to yield 0.20 g (69%) of powder **3**. ¹H NMR (CDCl₃) δ 7.08 (s, 1H, ArH), 6.86 (s, 1H, ArH), 6.16 (s, 1H, ArH), 4.24-4.21 (m, 4H, -CH₂CH₂-), 3.96-3.84 (m, 8H, -CH₂CH₂-), 3.33-3.27 (m, 4H, -CH₂CH₂-), 2.39 (s, 3H, CH₃).

1,8-(Etoxyethoxy-4-methylcoumarin)-anthraquinone (4). To a solution of 1,8-dihydroxyanthraquinone (0.20 g, 0.83 mmol) and K₂CO₃ (1.20 g, 8.70 mmol) in 60 ml. of THF/DMF (2:1), 6,7-bis[2-(2-iodoethoxy)ethyl]loxy-4-methylcoumarin **3** (0.50 g, 0.85 mmol) was added and the reaction mixture was stirred and refluxed for 3 days. The solvent was removed and residue was taken up in MeOH (50 mL). The residue was filtered and treated with 0.1 N HCl and extracted with CHCl₃. The organic phase was separated and dried with MgSO₄ and evaporated to yield 0.20 g (42%) of yellow powder **4**. ¹H NMR (CDCl₃) δ 7.85 (d, 2H, ArH, *J* = 7.5 Hz), 7.62 (t, 2H, ArH, *J* = 8.1 Hz), 7.28 (d, 2H, ArH), 7.09 (s, 1H, ArH), 6.84 (s, 1H, ArH), 6.13 (s, 1H, ArH), 4.29-4.17 (m, 16H, -CH₂CH₂-), 2.34 (s, 3H, -CH₃).

Acknowledgment. This work was supported by the Ministry of Education of Korea (BK 21 project).

References

- (a) Shinkai, S.; Ogawa, T.; Kusano, Y.; Manabe, O.; Kikukawa, K.; Goto, T.; Masuda, T. *J. Am. Chem. Soc.* **1982**, *104*, 1960. (b) Youhei, I.; Paul, K.; Yasushi, O.; Yoshinori, T.; Hiroyasu, Y.; Akira, H. *J. Am. Chem. Soc.* **2007**, *129*, 6396. (c) Roger, J.; Coulston, H. O.; Stephen, F. L.; Christopher, J. E. *J. Am. Chem. Soc.* **2006**, *128*, 14750.
- (a) Bissell, R. A.; Cordova, E.; Kaifer, A. E.; Stoddart, J. F. *Nature*

- 1994, 369, 133. (b) Badjic, J. D.; Balzani, V.; Credi, A.; Silvi, S.; Stoddart, J. F. *Science* **2004**, 303, 1845.
3. (a) Brouwer, A. M.; Frochet, C.; Gatti, F. G.; Leigh, D. A.; Mottier, L.; Paolucci, F.; Roffia, S.; Wurpel, G. W. H. *Science* **2001**, 291, 2124. (b) Shinkai, S.; Nakaji, T.; Nishida, Y.; Ogawa, T.; Manabe, O. *J. Am. Chem. Soc.* **1980**, 102, 5860. (c) Tomatsu, I.; Hashidzume, A.; Harada, A. *J. Am. Chem. Soc.* **2006**, 128, 2226.
4. (a) Armaroli, N.; Balzani, V.; Collin, J.-P.; Gavina, P.; Sauvage, J.-P.; Ventura, B. *J. Am. Chem. Soc.* **1999**, 121, 4397. (b) Liu, Y.; Flood, A. H.; Stoddart, J. F. *J. Am. Chem. Soc.* **2004**, 126, 9150.
5. (a) Gouloumis, A.; Lawson, R. C.; Vazquez, P.; Echegoyen, L.; Torres, T. *Tetrahedron* **2002**, 58, 961. (b) Awasthy, A.; Bhatnagar, M.; Tomar, J.; Sharma, U. *Biomorg. Chem. Appl.* **2006**, 16.
6. (a) Bako, P.; Mako, A.; Keglevich, G.; Menyhart, D. K.; Sefesik, T.; Fekete, J. *J. Incl. Phenom.* **2006**, 55, 295. (b) Torun, L.; Robison, T. W.; Krzykawski, J.; Purkiss, D. W.; Bartsch, R. A. *Tetrahedron* **2005**, 61, 8345.
7. (a) Gustowski, D. A.; Delgado, M.; Gatto, V. J.; Echegoyen, L.; Gokel, G. W. *J. Am. Chem. Soc.* **1986**, 108, 7553. (b) Barzegar, M.; Mousavi, M. F.; Khajehsharifi, H.; Shamsipur, M.; Sharghi, H. *IEEE Sensors J.* **2005**, 5, 392.
8. (a) Stone, E. W.; Maki, A. H. *J. Chem. Phys.* **1962**, 36, 1944. (b) Stauf, J. *Photochem. Photobiol.* **1965**, 4, 1199.
9. (a) Licheng, S.; Jörg-von, G.; Jens, S.; Harry, K. *Tetrahedron* **1995**, 51, 3535. (b) Rahman, M. A.; Kwon, N.; Won, M.; Hyun, M.; Shim, Y. *Anal. Chem.* **2004**, 76, 3660.
10. Delgado, M.; Gustowski, D. A.; Yoo, H. K.; Gatto, V. J.; Gokel, G. W.; Echegoyen, L. *J. Am. Chem. Soc.* **1988**, 110, 119.
11. Taziaux, D.; Soumillion, J.-Ph.; Habib-Jirwan, J.-L. *J. Photochem. Photobiol. A: Chem.* **2004**, 162, 599.
-