

Effects of Proton on Lysolipid-induced Actions in OGR1-subfamily GPCRs

Sung-Mee LIM¹ and Dong-Soon IM^{2*}

¹Department of Food Science & Technology, Tongmyong University, Busan 608-711, Republic of Korea and ²Laboratory of Pharmacology, College of Pharmacy (BK21 Project) and Research Institute of Drug Development, Pusan National University, Busan 609-735, Republic of Korea

(Received February 9, 2007; Accepted March 15, 2007)

Abstract – Lysolipids such as lysophosphatidylcholine (LPC), sphingosylphosphorylcholine (SPC), galactosylsphingosine (psychosine) have been matched as ligands for OGR1-subfamily G-protein-coupled receptors (GPCR), consisted of OGR1, GPR4, G2A, and TDAG8. Recently, those members of GPCRs have been reported as proton-sensing GPCRs. We used Jurkat T cells, which express four members of OGR1 subfamily GPCRs endogenously to investigate effects of proton on lysolipid-induced several cellular events. We found no significant effect of proton on the lysolipid-induced Ca²⁺ increase and ROS production in Jurkat T cells. Further investigation is necessary to clarify the relationship of lysolipid and proton on the OGR1-subfamily GPCRs.

Keywords □ lysolipid; proton, G-protein-coupled receptor, sphingosylphosphorylcholine, lysophosphatidylcholine, psychosine

INTRODUCTION

G-protein-coupled receptor is the largest gene family of human genome. Their importance is glaringly obvious by the fact that more than 50% of drugs on the market are either agonists or antagonists on GPCRs (Im 2002). Many chemical messengers such as hormones and neurotransmitters use their own GPCRs as signal receivers on the target cell membrane. Therefore, positive or negative modulation of GPCRs with drugs has been successful tools to treat many diseases such as allergy, gastric ulcer, and hypertension (Fredriksson et al. 2003; Im 2002, 2004). GPCR activations are evoked by stimuli as diverse as light, Ca²⁺, odorants, amino acids, nucleotides, proteins, polypeptides, steroids, and fatty acid derivatives.

The completion of the human genome project has identified about 865 GPCR genes (Perez 2005). Except sensory receptors, 367 GPCRs have been considered as receptors for endogenous ligands in the human genome (Vassilatis et al. 2003). OGR1 subfamily is composed of four members (OGR1, GPR4, G2A, and TDAG8) and has previously been identified as receptors for lysolipids; sphingosylphosphorylcholine (SPC),

lysophosphatidylcholine (LPC) and galactosylsphingosine (psychosine) (Im *et al.* 2001; Kabarowski *et al.* 2001; Xu *et al.* 2000; Zhu *et al.* 2001). And their importance has been reported as like G2A deficient mouse developed an autoimmune syndrome similar to systemic lupus erythematosus (SLE) (Le *et al.* 2001), and therapeutic application of LPC for sepsis was proposed in relation with G2A receptor (Yan *et al.* 2004). Recently, Kim *et al.* reported that GPR4 plays a critical role in SPC-induced angiogenesis and SPC transactivates VEGF receptor 2 in endothelial cells (Kim *et al.* 2005). Furthermore, Ikeno *et al.* reported that secretory phospholipase A₂ induce neurite outgrowth in PC12 cells through LPC generation and activation of G2A receptor (Ikeno *et al.* 2005).

Nevertheless, confirmation of the ligand matching of OGR1 subfamily with lysolipids has not been fully supported by publications of independent research groups (Im 2004, 2005). A series of publications propose that extracellular proton could be an activator of the OGR1 subfamily GPCRs (Ishii *et al.* 2005; Ludwig *et al.* 2003; Murakami *et al.* 2004; Radu *et al.* 2005; Wang *et al.* 2004). More than two independent groups reported proton-sensing properties of OGR1, GPR4, and TDAG8 (Ishii *et al.* 2005; Ludwig *et al.* 2003; Mogi *et al.* 2005; Murakami *et al.* 2004; Radu *et al.* 2005; Wang *et al.* 2004). In the case of G2A, constitutive activation at pH 7.4 has been observed in many transfected cells by many research groups, however, pH-

*Corresponding author

Tel: +82-51-510-2817, Fax: +82-51-513-6754

E-mail: imds@pusan.ac.kr

dependent activation was supported only by one group and was not fully reproduced by another group (Murakami *et al.* 2004; Radu *et al.* 2005).

Reactive oxygen species (ROS) exert numerous effects on cell functions, including induction of growth and regulation of kinase activity (Finkel 2000; Larsson and Cerutti 1988; Schreck *et al.* 1991). Oxidative stress plays a prominent role in the functioning of the immune system at both physiological and pathological levels, and also is known to be associated with reduced or abnormal immune function with aging (Lieber 1998), AIDS (Dobmeyer *et al.* 1997), diabetes (Dandona *et al.* 1996), smoking-related pathologies (McAllister-Sistilli *et al.* 1998), and some autoimmune and inflammatory diseases (Frenkel *et al.* 1992).

Calcium is a ubiquitous second messenger controlling a broad range of cellular functions, and an increase of intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) plays an important role in cellular functions such as cell proliferation and insulin secretion (Himmel *et al.* 1998; Lipskaia and Lompre 2004).

Lysophospholipids regulate a variety of biological processes including cell proliferation, tumor cell invasiveness, and inflammation (Moolenaar 1999; Spiegel and Milstien 1995). Jurkat T cell line is homogeneous T lymphocytes and possess membrane markers that characterize normal T lymphocytes (Konikova *et al.* 1992). In the previous study, LPC was found to induce ROS generation and $[\text{Ca}^{2+}]_i$ increase in human lymphocytic Jurkat T cells (Im *et al.* 2006). Furthermore, four members of OGR1 subfamily GPCRs are expressed endogenously (Im *et al.* 2006). Because all four members of OGR1 subfamily GPCR were detected at mRNA level in Jurkat T cells, we applied this cell line as a model.

MATERIALS AND METHODS

Materials

1-palmitoyl (C16:0) LPC, 1-oleoyl lysophosphatidic acid (LPA), sphingosyl phosphorylcholine (SPC), sphingosine 1-phosphate (S1P), and galactosyl sphingosine (psychosine, Psy) were purchased from Avanti Polar Lipids (Alabaster, AL, USA); fura 2/acetoxymethyl ester was from Calbiochem (Darmstadt, Germany). 2',7' dichlorofluorescein diacetate (DCFDA) were from Sigma-Aldrich (St. Louis, MO, USA).

Cell culture

Jurkat T cells were maintained in RPMI 1640 containing 10 % (v/v) fetal bovine serum, 100 units/ml penicillin, 50 $\mu\text{g}/\text{ml}$

streptomycin, 2 mM glutamine, and 1 mM sodium pyruvate at 37°C in a humidified 5% CO incubator.

Measurement of ROS

Jurkat T cells were sedimented, resuspended in HEPES-buffered media (HBM) consisting of 20 mM HEPES (pH 7.4), 103 mM NaCl, 4.8 mM KCl, 1.2 mM KH_2PO_4 , 1.2 mM MgSO_4 , 0.5 mM CaCl_2 , 25 mM NaHCO_3 and 15 mM glucose, and then incubated for 20 min with 10 μM DCFDA at pH 7.4. Then, the cells were used for the measurements. Fluorescence was measured in DCFDA-loaded cells. Fluorescence emission at 520 nm wavelength from 488 nm excitation wavelength was measured every 0.1 sec by an F4500 fluorescent spectrophotometer (Hitachi, Japan). The extracellular pH was adjusted by addition of pre-calculated amounts of 1 N HCl for each pH like 7.1, 6.8, and 6.5. After 15 sec, fluorescent intensity was recorded at the first 60 sec, 60 sec after lipid addition and 10 min later. Individual histogram was made by the percent of control level, and ROS was measured more than twice (Im *et al.* 2006).

Measurement of intracellular Ca^{2+} concentration

Different pHs of HBM were adjusted just before the experiments by addition of 1 N HCl. The Jurkat T cells were sedimented, resuspended in the HBM (pH 7.4), and then incubated for 40 min with 5 μM fura 2/acetoxymethyl ester at pH 7.4. Fura 2-loaded cells were washed twice with each HBM with different pH and resuspended in the same media. Fluorescence emission at 510 nm wavelength from two excitation wavelengths (340 nm and 380 nm) were measured every 0.1 sec (F4500, Hitachi, Japan). The ratio of fluorescence intensities from the two wavelengths was monitored as an estimate of intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) (Lee *et al.* 2006).

Data presentation and statistical analysis

Representative traces for intracellular Ca^{2+} concentration were chosen from 3-5 separate experiments and are shown in Figs 1 and 2. In Fig 3, results from two separate experiments were shown by the percent of control level.

RESULTS

Effect of extracellular pH on LPC-induced $[\text{Ca}^{2+}]_i$ increase

Because all four members of OGR1 subfamily GPCR were detected at mRNA level in Jurkat T cells, we applied this cell line as a model to study functions of endogenous OGR1 GPCRs. Recent data show that LPC elevates the intracellular




Fig. 1. Effect of proton on LPC-induced increase of intracellular Ca^{2+} concentration. After loading Fura-2/AM into Jurkat T cells ($1.5 \times 10^6/\text{cells/ml}$) described in Method section, cells suspended in HBM with different pHs without BSA were divided into 2 ml cuvette. Each trace was monitored and saved for F4500 fluorescent spectrophotometer. Representative Ca^{2+} traces with $20 \mu\text{M}$ of LPC in Jurkat T cells at different extracellular pH. LPC was added at the arrow indicated. The data shown are representative of three independent experiments.

Ca^{2+} concentration in Jurkat T cells (Im *et al.* 2006; Legradi *et al.* 2004). As shown in Fig. 1, $20 \mu\text{M}$ concentration of LPC sharply increased $[\text{Ca}^{2+}]_i$ in Jurkat T cells. The effect of extracellular pH on LPC-induced increase of $[\text{Ca}^{2+}]_i$ was studied. As shown in Fig 1, the increase was not influenced by lowering extracellular pH upto 6.5. Further, we tested effects of other lysolipids including SPC, Psy, LPA, and S1P. Those lipids did not increase the intracellular Ca^{2+} concentration in different

extracellular pHs (Fig. 2).

Effect of extracellular pH on LPC stimulated ROS generation in Jurkat T cells

Previously, we reported that LPC rapidly and significantly increased reactive oxygen species (ROS) level in Jurkat T cells (Im *et al.* 2006). We measured changes of ROS levels by lysolipids and influences of extracellular pH on the effect. As




Fig. 2. Effect of lysolipid on intracellular Ca^{2+} concentration at different extracellular pH. Representative Ca^{2+} traces with $20 \mu\text{M}$ of SPC, S1P, Psy, or LPA in Jurkat T cells at different extracellular pH. Each lipid was added at the arrow indicated. The data shown are representative of three independent experiments.




Fig. 3. Effect of proton on lysolipid-induced increase of ROS generation. After loading DCFDA into Jurkat T cells (1.5×10^6 cells/ml) as described in Method section, cells suspended in HBM without BSA were divided into 2 ml cuvette. Each trace was monitored and saved for F4500 fluorescent spectrophotometer. After adjusting the pH by HCl, fluorescent intensity was monitored at the first 60 sec, 60 sec after lipid addition and 10 min later. The first ten points of each step of times were averaged, and individual histogram was made by the percent of control level. Open column means ROS level after lysolipid addition and filled column means ROS level 10 min after lysolipid addition.

shown in Fig. 3, LPC increased ROS most prominently and the increase was followed by LPA and SIP. Effect of proton on lysolipid-induced ROS generation was tested by changing extracellular pH (Fig. 3). However, decrease of extracellular pH, in other word increase of proton in the media, did not affect on the increase of ROS by each lipid (Fig. 3).

DISCUSSION

In the present study, we were not able to find any effect of proton on lysolipid-mediated responses such as Ca^{2+} increase and ROS generation. Proton itself also did not induce any change in Ca^{2+} increase in Jurkat T cells (data not shown). In the previous study, we concluded that LPC-induced Ca^{2+} increase and ROS generation may not be mediated through GPCRs in Jurkat T cells. In this study, we found no effect of proton on lysolipid-induced responses. Even though four members of OGR1 subfamily GPCRs are expressed in Jurkat T cells, no significant change was observed by lysolipid and proton. Three possibilities could be considered; 1) different signalings of OGR1 subfamily GPCRs triggered by proton in the

Jurkat T cells to signalings of proton reported in overexpression system of the GPCRs, 2) dysfunction of four OGR1 subfamily GPCRs in the Jurkat T cells, and 3) the idea of proton-sensing GPCR of OGR1 subfamily GPCRs may not be correct as like lysolipid.

ACKNOWLEDGMENTS

This work was supported by the Grand-in-aid for 2005 Korea-Japan Joint Research Project from Korea Science and Engineering Foundation (DSI, F01-2005-000-10046-0).

REFERENCES

- Chang YJ, Lee YK, Lee EH, Park JJ, Chung SK and Im DS. Structure-activity relationships of dimethylsphingosine (DMS) derivatives and their effects on intracellular pH and Ca^{2+} in the U937 monocyte cell line. *Arch Pharm Res* **29**, 657-665, 2006.
- Dandona P, Thusu K, Cook S, Snyder B, Makowski J, Armstrong D and Nicotera T. Oxidative damage to DNA in diabetes mellitus. *Lancet* **347**, 444-445, 1996.
- Dobmeyer TS, Findhammer S, Dobmeyer JM, Klein SA, Raffel B, Hoelzer D, Helm EB, Kabelitz D and Rossol R. Ex vivo

- induction of apoptosis in lymphocytes is mediated by oxidative stress: role for lymphocyte loss in HIV infection. *Free Radic Biol Med* **22**, 775-785, 1997.
- Finkel T. Redox-dependent signal transduction. *FEBS Lett* **476**, 52-54, 2000.
- Fredriksson R, Lagerstrom MC, Lundin LG and Schiöth HB. The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. *Mol Pharmacol* **63**, 1256-1272, 2003.
- Frenkel K, Khasak D, Karkoszka J, Shupack J and Stiller M. Enhanced antibody titers to an oxidized DNA base in inflammatory and neoplastic diseases. *Exp Dermatol* **1**, 242-247, 1992.
- Himmel HM, Meyer zu Heringdorf D, Windorfer B, van Koppen CJ, Ravens U and Jakobs KH. Guanine nucleotide-sensitive inhibition of L-type Ca^{2+} current by lysosphingolipids in RINm5F insulinoma cells. *Mol Pharmacol* **53**, 862-869, 1998.
- Ikeno Y, Konno N, Cheon SH, Bolchi A, Ottonello S, Kitamoto K and Arioka M. Secretory phospholipase A_2 induces neurite outgrowth in PC12 cells through lysophosphatidylcholine generation and activation of G2A receptor. *J Biol Chem* **280**, 28044-28052, 2005.
- Im DS. Orphan G protein-coupled receptors and beyond. *Jpn J Pharmacol* **90**, 101-106, 2002.
- Im DS. Discovery of new G protein-coupled receptors for lipid mediators. *J Lipid Res* **45**, 410-418, 2004.
- Im DS. Two ligands for a GPCR, proton vs lysolipid. *Acta Pharmacol Sin* **26**: 1435-1441, 2005.
- Im DS, Heise CE, Nguyen T, O'Dowd BF and Lynch KR. Identification of a molecular target of psychosine and its role in globoid cell formation. *J Cell Biol* **153**, 429-434, 2001.
- Im YJ, Lee YK, Chung HY and Im DS. Multiple actions of lysophosphatidylcholine in human Jurkat T cells. *Acta Pharmacol Sin* **27**, 700-707, 2006.
- Ishii S, Kihara Y and Shimizu T. Identification of T cell death-associated gene 8 (TDAG8) as a novel acid sensing G-protein-coupled receptor. *J Biol Chem* **280**, 9083-9087, 2005.
- Jatana M, Giri S and Singh AK. Apoptotic positive cells in Krabbe brain and induction of apoptosis in rat C6 glial cells by psychosine. *Neurosci Lett* **330**, 183-187, 2002.
- Kabarowski JH, Zhu K, Le LQ, Witte ON and Xu Y. Lysophosphatidylcholine as a ligand for the immunoregulatory receptor G2A. *Science* **293**, 702-705, 2001.
- Kim HS, Lee EH, Ko SR, Choi KI, Park JH and Im DS. Effects of ginsenosides Rg3 and Rh2 on the proliferation of prostate cancer cells. *Arch Pharm Res* **27**, 429-435, 2004.
- Kim KS, Ren J, Jiang Y, Ebrahim Q, Tipps R, Cristina K, Xiao YJ, Qiao J, Taylor KL, Lum H, Anand-Apte B and Xu Y. GPR4 plays a critical role in endothelial cell function and mediates the effects of sphingosylphosphorylcholine. *Faseb J* **19**, 819-821, 2005.
- Konikova E, Babusikova O, Kusenda J and Glasova M. Detection of cytoplasmic and surface membrane markers in cells of some human hematopoietic cell lines. *Neoplasma* **39**, 337-342, 1992.
- Larsson R and Cerutti P. Oxidants induce phosphorylation of ribosomal protein S6. *J Biol Chem* **263**, 17452-17458, 1988.
- Le LQ, Kabarowski JH, Weng Z, Satterthwaite AB, Harvill ET, Jensen ER, Miller JF and Witte ON. Mice lacking the orphan G protein-coupled receptor G2A develop a late-onset autoimmune syndrome. *Immunity* **14**, 561-571, 2001.
- Lee EH, Lee YK, Im YJ, Kim JH, Okajima F and Im DS. Dimethylsphingosine regulates intracellular pH and Ca^{2+} in human monocytes. *J Pharmacol Sci* **100**, 289-296, 2006.
- Legradi A, Chitu V, Szukacsov V, Fajka-Boja R, Szekely Szucs K and Monostori E. Lysophosphatidylcholine is a regulator of tyrosine kinase activity and intracellular Ca^{2+} level in Jurkat T cell line. *Immunol Lett* **91**, 17-21, 2004.
- Lieber MR. Warner-Lambert/Parke-Davis Award Lecture. Pathological and physiological double-strand breaks: roles in cancer, aging, and the immune system. *Am J Pathol* **153**, 1323-1332, 1998.
- Lipskaia L and Lompre AM. Alteration in temporal kinetics of Ca^{2+} signaling and control of growth and proliferation. *Biol Cell* **96**, 55-68, 2004.
- Ludwig MG, Vanek M, Guerini D, Gasser JA, Jones CE, Junker U, Hofstetter H, Wolf RM and Seuwen K. Proton-sensing G-protein-coupled receptors. *Nature* **425**, 93-98, 2003.
- McAllister-Stylli CG, Caggiola AR, Knopf S, Rose CA, Miller AL and Donny EC. The effects of nicotine on the immune system. *Psychoneuroendocrinology* **23**, 175-187, 1998.
- Mogi C, Tomura H, Tobo M, Wang JQ, Damirin A, Kon J, Komachi M, Hashimoto K, Sato K and Okajima F. Sphingosylphosphorylcholine antagonizes proton-sensing ovarian cancer G-protein-coupled receptor 1 (OGR1)-mediated inositol phosphate production and cAMP accumulation. *J Pharmacol Sci* **99**, 160-167, 2005.
- Moolenaar WH. Bioactive lysophospholipids and their G protein-coupled receptors. *Exp Cell Res* **253**, 230-238, 1999.
- Murakami N, Yokomizo T, Okuno T and Shimizu T. G2A is a proton-sensing G-protein coupled receptor antagonized by lysophosphatidylcholine. *J Biol Chem* **279**, 42484-42491, 2004.
- Perez DM. From Plants to Man: The GPCR "Tree of Life". *Mol Pharmacol* **67**, 1383-1384, 2005.
- Radu CG, Nijagal A, McLaughlin J, Wang L and Witte ON. Differential proton sensitivity of related G protein-coupled receptors T cell death-associated gene 8 and G2A expressed in immune cells. *Proc Natl Acad Sci U S A* **102**, 1632-1637, 2005.
- Schreck R, Rieber P and Baeuerle PA. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *Embo J* **10**, 2247-2258, 1991.
- Spiegel S and Milstien S. Sphingolipid metabolites: members of a new class of lipid second messengers. *J Membr Biol* **146**, 225-237, 1995.
- Sueyoshi N, Maehara T and Ito M. Apoptosis of Neuro2a cells induced by lysosphingolipids with naturally occurring stereochemical configurations. *J Lipid Res* **42**, 1197-1202, 2001.
- Vassilatis DK, Hohmann JG, Zeng H, Li F, Ranchalis JE, Mortrud MT, Brown A, Rodriguez SS, Weller JR, Wright AC, Bergmann JE and Gaitanaris GA. The G protein-coupled receptor repertoires of human and mouse. *Proc Natl Acad Sci U S A* **100**, 4903-4908, 2003.
- Wang JQ, Kon J, Mogi C, Tobo M, Damirin A, Sato K, Komachi M, Malchinkhuu E, Murata N, Kimura T, Wakamatsu K, Koizumi H, Uede T, Tsujimoto G, Kurose H, Sato T, Harada A, Misawa N, Tomura H and Okajima F. TDAG8 is a proton-sensing and psychosine-sensitive G-protein-coupled receptor. *J Biol Chem* **279**, 45626-45633, 2004.
- Xu Y, Zhu K, Hong G, Wu W, Baudhuin LM, Xiao Y and Dameron DS. Sphingosylphosphorylcholine is a ligand for ovarian cancer G-protein-coupled receptor 1. *Nat Cell Biol* **2**, 261-267, 2000.
- Yan JJ, Jung JS, Lee JE, Lee J, Huh SO, Kim HS, Jung KC, Cho JY, Nam JS, Suh HW, Kim YH and Song DK. Therapeutic effects of lysophosphatidylcholine in experimental sepsis. *Nat*

Med **10**, 161-167, 2004.

Zaka M and Wenger DA. Psychosine-induced apoptosis in a mouse oligodendrocyte progenitor cell line is mediated by caspase activation. *Neurosci Lett* **358**, 205-209, 2004.

Zhu K, Baudhuin LM, Hong G, Williams FS, Cristina KL,

Kabarowski JH, Witte ON and Xu Y. Sphingosylphosphorylcholine and lysophosphatidylcholine are ligands for the G protein-coupled receptor GPR4. *J Biol Chem* **276**, 41325-41335, 2001.