

NEW RELATIONSHIPS INVOLVING THE MEAN CURVATURE OF SLANT SUBMANIFOLDS IN S -SPACE-FORMS

LUIS M. FERNÁNDEZ AND MARÍA BELÉN HANS-ÜBER

ABSTRACT. Relationships between the Ricci curvature and the squared mean curvature and between the shape operator associated with the mean curvature vector and the sectional curvature function for slant submanifolds of an S -space-form are proved, particularizing them to invariant and anti-invariant submanifolds tangent to the structure vector fields.

1. Introduction

In words of B.-Y. Chen, to “find simple relationships between the main extrinsic invariant and the main intrinsic invariants of a submanifold” is one of the basic problems in the theory of submanifolds [7]. In this way, he established a relationship between sectional curvature function and the shape operator for submanifolds in real space-forms [7] and another relationship between the Ricci curvature and the squared mean curvature [8]. Corresponding relationships have been proved in [14] for slant submanifolds of Sasakian space-forms.

Slant immersions in complex geometry were defined by B.-Y. Chen as a natural generalization of both holomorphic and totally real immersions [9]. Recently, A. Lotta has introduced the notion of slant immersion of a Riemannian manifold into an almost contact metric manifold [12] and slant submanifolds of Sasakian manifolds have been studied in [2]. For a general view about slant submanifolds, the survey written by A. Carriazo [3] can be consulted. On the other hand, for manifolds with an f -structure, D. E. Blair has introduced S -manifolds as the analogue of the Kaehler structure in the almost complex case and of Sasakian structure in the almost contact case [1] and we have defined and begun with the study of slant submanifolds in such S -manifolds [4, 5, 6].

The purpose of this paper is to obtain similar relationships to Chen’s ones mentioned above, generalizing and improving in some sense the ones proved

Received November 24, 2005.

2000 *Mathematics Subject Classification.* 53C25, 53C40.

Key words and phrases. S -space-form, slant immersion, mean curvature vector, Ricci curvature, shape operator.

The first author is partially supported by the PAI project (Junta de Andalucía, Spain, 2005).

in [14], for slant submanifolds in S -space-forms. To this end, after reviewing, for later use, necessary details about S -manifolds and slant submanifolds in Section 2, we devote Section 3 to get an inequality between Ricci curvature and squared mean curvature vector and discuss the equality case. Finally, in Section 4 we establish an inequality between the shape operator associated with the mean curvature vector and the sectional curvature function for slant submanifolds of S -space forms. In both cases, we particularize these inequalities for invariant and anti-invariant submanifolds tangent to the structure vector fields.

2. Preliminaries

Let (\widetilde{M}, g) be a Riemannian manifold and denote by $\mathcal{X}(\widetilde{M})$ the Lie algebra of vector fields in \widetilde{M} . Then \widetilde{M} is said to be a metric f -manifold if there exist a $(1, 1)$ tensor field f , s global unit vector fields ξ_1, \dots, ξ_s (called structure vector fields) and s 1-forms η_1, \dots, η_s on \widetilde{M} such that

$$f^2 X = -X + \sum_{\alpha=1}^s \eta_\alpha(X) \xi_\alpha, \quad g(X, \xi_\alpha) = \eta_\alpha(X),$$

$$f \xi_\alpha = 0, \quad \eta_\alpha \circ f = 0$$

and

$$g(fX, fY) = g(X, Y) - \sum_{\alpha=1}^s \eta_\alpha(X) \eta_\alpha(Y),$$

for any $X, Y \in \mathcal{X}(\widetilde{M})$ and $\alpha = 1, \dots, s$. Let F denote the fundamental 2-form in \widetilde{M} given by $F(X, Y) = g(X, fY)$, for any $X, Y \in \mathcal{X}(\widetilde{M})$. The tensor field f is said to be normal if $[f, f] + 2 \sum_{\alpha} \xi_\alpha \otimes d\eta_\alpha = 0$, where $[f, f]$ is the Nijenhuis torsion of f . Thus \widetilde{M} is called an S -manifold if the structure is normal and $F = d\eta_\alpha$, for any $\alpha = 1, \dots, s$.

Given an S -manifold \widetilde{M} , a plane section π in $T_p \widetilde{M}$ is called an f -section if it is spanned by X and fX , where X is a unit tangent vector field orthogonal to the distribution \mathcal{M} spanned by the structure vector fields. The sectional curvature $K(\pi)$ of an f -section π is called f -sectional curvature. An S -manifold is said to be an S -space-form if it has constant f -sectional curvature c and then, it is denoted by $\widetilde{M}(c)$. As examples of S -space-forms, we can mention $\mathbf{R}^{2m+s}(-3s)$ with its standard S -structure given in [10] and $\mathcal{H}^{2m+s}(4-3s)$, where

$$\mathcal{H}^{2m+s} = \{(x_1, \dots, x_s) \in S^{2m+1} \times \dots \times S^{2m+1} / \pi(x_1) = \dots = \pi(x_s)\},$$

being $\pi : S^{2m+1} \rightarrow \mathbf{P}C^m$ the Hopf fibration (see [1] for more details). More interesting examples of S -space-forms can be found in [10].

Now, let M be a submanifold isometrically immersed in an S -manifold \widetilde{M} . Let $\mathcal{X}(M)$ be the Lie algebra of vector fields in M and $T^\perp M$ the set of all vector fields normal to M . We denote by σ the second fundamental form of M and by A_V the shape operator associated with any $V \in T^\perp M$. They are related by

the equation $g(\sigma(X, Y), V) = g(A_V X, Y)$, for any $X, Y \in \mathcal{X}(M)$ and any $V \in T^\perp M$. The mean curvature vector H is defined by $H = (1/\dim(M))\text{trace}(\sigma)$. The submanifold is said to be minimal if H vanishes identically and it is said to be totally geodesic if $\sigma(X, Y) = 0$, for any $X, Y \in \mathcal{X}(M)$. Moreover, the relative null space of M is defined by:

$$\mathcal{N} = \{X \in \mathcal{X}(M) : \sigma(X, Y) = 0, \text{ for all } Y \in \mathcal{X}(M)\}.$$

For any $X \in \mathcal{X}(M)$, we put $fX = TX + NX$, where TX (resp. NX) is the tangential (resp. normal) component of fX . The submanifold M is said to be invariant if N is identically zero, that is, if $fX \in \mathcal{X}(M)$, for any $X \in \mathcal{X}(M)$ and it is said to be anti-invariant if T is identically zero, that is, if $fX \in T^\perp M$, for any $X \in \mathcal{X}(M)$.

From now on, we suppose that the structure vector fields are tangent to M and we denote by $n + s$ (resp. $2m + s$) the dimension of M (resp. \widetilde{M}). Hence, if we denote by \mathcal{L} the orthogonal distribution to \mathcal{M} in $\mathcal{X}(M)$, we can write the orthogonal direct decomposition $\mathcal{X}(M) = \mathcal{L} \oplus \mathcal{M}$.

It is well-known that

$$(2.1) \quad \sigma(X, \xi_\alpha) = -NX,$$

for any $X \in \mathcal{X}(M)$ and any $\alpha = 1, \dots, s$. In particular, $\sigma(\xi_\alpha, \xi_\beta) = 0$, for any $\alpha, \beta = 1, \dots, s$.

Now, given a local orthonormal basis

$$\{e_1, \dots, e_n, e_{n+1}, \dots, e_{2m}, \xi_1, \dots, \xi_s\}$$

of $\mathcal{X}(\widetilde{M})$, such that $\{e_1, \dots, e_n\}$ is a local orthonormal basis of \mathcal{L} , we can write the mean curvature vector H and the squared norms of T and σ by:

$$(2.2) \quad H = \frac{1}{n + s} \sum_{i=1}^n \sigma(e_i, e_i),$$

$$(2.3) \quad \|T\|^2 = \sum_{i,j=1}^n g^2(e_i, Te_j),$$

$$(2.4) \quad \|\sigma\|^2 = \sum_{i=1}^n \|\sigma(e_i, e_i)\|^2 + 2 \sum_{1 \leq i < j \leq n} \|\sigma(e_i, e_j)\|^2 + 2 \sum_{i=1}^n \sum_{\alpha=1}^s \|\sigma(e_i, \xi_\alpha)\|^2.$$

The curvature tensor field R of a submanifold M of an S -space-form $\widetilde{M}(c)$ satisfies

$$\begin{aligned} R(X, Y, Z, W) &= g(\sigma(X, W), \sigma(Y, Z)) - g(\sigma(X, Z), \sigma(Y, W)) \\ &\quad + \sum_{\alpha, \beta} (g(fX, fW)\eta_\alpha(Y)\eta_\beta(Z) - g(fX, fZ)\eta_\alpha(Y)\eta_\beta(W)) \\ &\quad + g(fY, fZ)\eta_\alpha(X)\eta_\beta(W) - g(fY, fW)\eta_\alpha(X)\eta_\beta(Z) \end{aligned}$$

$$(2.5) \quad \begin{aligned} & + \frac{c + 3s}{4} (g(fX, fW)g(fY, fZ) - g(fX, fZ)g(fY, fW)) \\ & + \frac{c - s}{4} (F(X, W)F(Y, Z) - F(X, Z)F(Y, W) - 2F(X, Y)F(Z, W)), \end{aligned}$$

for any $X, Y, Z, W \in \mathcal{X}(\widetilde{M})$ (see, for references, [1, 11]).

The scalar curvature τ of M is defined by

$$(2.6) \quad \tau = \frac{1}{2} \sum_{i \neq j} K(e_i \wedge e_j) + \sum_{i=1}^n \sum_{\alpha=1}^s K(e_i \wedge \xi_\alpha),$$

where $K(X \wedge Y)$ denotes the sectional curvature of M associated with the plane section spanned by $X, Y \in \mathcal{X}(M)$.

From (2.1)-(2.6), we obtain the following relationship between the scalar curvature and the mean curvature of M [4]:

$$(2.7) \quad 2\tau = (n + s)^2 \|H\|^2 - \|\sigma\|^2 + n(n - 1) \frac{c + 3s}{4} + 2ns + \frac{3(c - s)}{4} \|T\|^2.$$

If for each nonzero vector $X \in T_pM - \mathcal{M}_p$, we consider the angle $\theta(X)$ between fX and T_pM , then the submanifold is said to be θ -slant [6] if such angle is a constant, which is independent on the choice of $p \in M$ and $X \in T_pM - \mathcal{M}_p$. The angle θ of a slant immersion is called the *slant angle* of the immersion. Invariant and anti-invariant submanifolds tangent to the structure vector fields are slant submanifolds with slant angle $\theta = 0$ and $\theta = \pi/2$, respectively. A slant immersion which is neither invariant nor anti-invariant is called a proper slant immersion. Different examples of slant submanifolds of S -manifolds are given in [6].

Finally, in [6], we have also proved that a θ -slant submanifold M of a metric f -manifold \widetilde{M} satisfies

$$g(TX, TY) = \cos^2 \theta g(fX, fY),$$

for any $X, Y \in \mathcal{X}(M)$. Moreover, it is easy to show that, using a local orthonormal basis $\{e_1, \dots, e_{n+s}\}$ of $\mathcal{X}(M)$,

$$(2.8) \quad \sum_{j=1}^{n+s} g^2(e_i, fe_j) = \cos^2 \theta (1 - \sum_{\alpha=1}^s \eta_\alpha^2(e_i)),$$

for any $i = 1, \dots, s$.

3. The Ricci curvature for slant immersions

Let M be an $(n + s)$ -dimensional isometrically immersed submanifold in a $(2m + s)$ -dimensional S -space-form $\widetilde{M}(c)$, tangent to the structure vector fields. In this section, we want to study the Ricci curvature of unit vector fields in M , normal to the structure vector fields, when M is a slant submanifold and to relate them with the mean curvature vector. For this reason, we shall assume that $n \geq 2$, because it is known that there are not proper slant submanifolds of

dimension $1 + s$ ([5]). Throughout the section, we consider local orthonormal basis of $\mathcal{X}(\widetilde{M}(c))$,

$$(3.1) \quad \{e_1, \dots, e_n, e_{n+1}, \dots, e_{2m}, e_{2m+1} = \xi_1, \dots, e_{2m+s} = \xi_s\},$$

such that $\{e_1, \dots, e_n\}$ is a local orthonormal basis of \mathcal{L} . First, we have the following general result:

Theorem 3.1. *Let M be an $(n + s)$ -dimensional submanifold $\widetilde{M}(c)$, tangent to the structure vector fields. Then,*

$$(3.2) \quad 4\text{Ric}(U) \leq (n + s)^2 \|H\|^2 + (n - 1)(c + 3s) + \|TU\|^2(3c + s),$$

for any unit vector field $U \in \mathcal{L}$.

Proof. We choose a local orthonormal basis of $\mathcal{X}(\widetilde{M}(c))$ as in (3.1) and such that $e_1 = U$. Then:

$$(3.3) \quad \begin{aligned} \|\sigma\|^2 &= \frac{1}{2}(n + s)^2 \|H\|^2 + 2 \sum_{1 \leq i < j \leq n} \|\sigma(e_i, e_j)\|^2 + 2 \sum_{i=1}^n \sum_{\alpha=1}^s \|\sigma(e_i, \xi_\alpha)\|^2 \\ &+ \frac{1}{2} \sum_{i=1}^n \|\sigma(e_i, e_i)\|^2 - \sum_{1 \leq i < j \leq n} g(\sigma(e_i, e_i), \sigma(e_j, e_j)). \end{aligned}$$

Thus, from (2.7) and (3.3), we have:

$$(3.4) \quad \begin{aligned} \frac{1}{4}(n + s)^2 \|H\|^2 &= \tau - \frac{1}{8}n(n - 1)(c + 3s) - \frac{3}{8}\|T\|^2(c - s) - ns \\ &+ \sum_{1 \leq i < j \leq n} \|\sigma(e_i, e_j)\|^2 + \sum_{i=1}^n \sum_{\alpha=1}^s \|\sigma(e_i, \xi_\alpha)\|^2 \\ &+ \frac{1}{4} \sum_{i=1}^n \|\sigma(e_i, e_i)\|^2 - \frac{1}{2} \sum_{1 \leq i < j \leq n} g(\sigma(e_i, e_i), \sigma(e_j, e_j)). \end{aligned}$$

On the other hand, since from (2.6),

$$\tau = \text{Ric}(U) + \sum_{2 \leq i < j \leq n} K(e_i \wedge e_j) + \sum_{i=2}^n \sum_{\alpha=1}^s K(e_i \wedge \xi_\alpha)$$

and, by using (2.5), we get

$$\begin{aligned} \sum_{2 \leq i < j \leq n} K(e_i \wedge e_j) &= \frac{1}{8}n(n - 1)(c + 3s) + \frac{3}{4} \left(\frac{\|T\|^2}{2} - \|TU\|^2 \right) (c - s) \\ &+ \sum_{2 \leq i < j \leq n} (g(\sigma(e_i, e_i), \sigma(e_j, e_j)) - \|\sigma(e_i, e_j)\|^2) \end{aligned}$$

and

$$\sum_{i=2}^n \sum_{\alpha=1}^s K(e_i \wedge \xi_\alpha) = (n - 1)s - \sum_{i=2}^n \sum_{\alpha=1}^s \|\sigma(e_i, \xi_\alpha)\|^2,$$

then, substituting into (3.4) and taking into account (2.1), we obtain:

$$\begin{aligned}
 \text{Ric}(U) &= \frac{1}{4}(n+s)^2\|H\|^2 + \frac{1}{4}(n-1)(c+3s) + \frac{1}{4}\|TU\|^2(3c+s) \\
 &\quad - \frac{1}{2} \sum_{2 \leq i < j \leq n} g(\sigma(e_i, e_i), \sigma(e_j, e_j)) + \frac{1}{2} \sum_{i=2}^n g(\sigma(U, U), \sigma(e_i, e_i)) \\
 (3.5) \quad &\quad - \sum_{i=2}^n \|\sigma(U, e_i)\|^2 - \frac{1}{4} \sum_{i=1}^n \|\sigma(e_i, e_i)\|^2.
 \end{aligned}$$

Now, if we put $\sigma_{ij}^r = g(\sigma(e_i, e_j), e_r)$, for any $i, j = 1, \dots, n$ and $r = n+1, \dots, 2m$, (3.5) becomes

$$\begin{aligned}
 \text{Ric}(U) &= \frac{1}{4}(n+s)^2\|H\|^2 + \frac{1}{4}(n-1)(c+3s) + \frac{1}{4}\|TU\|^2(3c+s) \\
 (3.6) \quad &\quad - \sum_{r=n+1}^{2m} \left[\frac{1}{4} \left(\sigma_{11}^r - \sum_{i=2}^n \sigma_{ii}^r \right)^2 + \sum_{i=2}^n (\sigma_{1i}^r)^2 \right],
 \end{aligned}$$

which completes the proof. □

Observe that, if we also put $\sigma_{1(2m+\alpha)}^r = g(\sigma(U, \xi_\alpha), e_r) = -g(NU, e_r)$, for any $\alpha = 1, \dots, s$ and $r = n+1, \dots, 2m$, we have

$$\begin{aligned}
 \text{Ric}(U) &= \frac{1}{4}(n+s)^2\|H\|^2 + \frac{1}{4}n(c+3s) + \frac{1}{4}(3\|TU\|^2 - 1)(c-s) \\
 (3.7) \quad &\quad - \sum_{r=n+1}^{2m} \left[\frac{1}{4} \left(\sigma_{11}^r - \sum_{i=2}^n \sigma_{ii}^r \right)^2 + \sum_{i=2}^n (\sigma_{1i}^r)^2 + \sum_{\alpha=1}^s (\sigma_{1(2m+\alpha)}^r)^2 \right]
 \end{aligned}$$

and, consequently:

$$(3.8) \quad 4\text{Ric}(U) \leq (n+s)^2\|H\|^2 + n(c+3s) + (3\|TU\|^2 - 1)(c-s).$$

When $s = 1$, this upper bound is the one obtained in [14] for submanifolds of Sasakian space-forms tangent to the structure vector field. However, it is easy to show that it is worse (in the sense of higher) than the one of (3.2). Moreover, both upper bounds are equal if and only if $\|NU\|^2 = 0$, that is, if and only if, from (2.1), $\sigma(U, \xi_\alpha) = 0$, for any $\alpha = 1, \dots, s$. If it is the case, their common value is $(n+s)^2\|H\|^2 + n(c+3s) + 2(c-s)$. Then, we can prove the following theorem.

Theorem 3.2. *Let M be an $(n+s)$ -dimensional minimal submanifold of $\widetilde{M}(c)$, tangent to the structure vector fields. Then, a unit vector field U in \mathcal{L} satisfies the equality case of (3.8) if and only if U lies in the relative null space of M . Moreover, in this case:*

$$4\text{Ric}(U) = n(c+3s) + 2(c-s).$$

Proof. If $U \in \mathcal{L}$ is a unit vector field satisfying the equality case of (3.8), then it also satisfies the equality case of (3.2) and so, $\sigma(U, \xi_\alpha) = 0$, for any $\alpha = 1, \dots, s$. Furthermore, choosing a local orthonormal basis of $\mathcal{X}(\widetilde{M}(c))$ as in (3.1) such that $e_1 = U$, from (3.7) we get $\sigma_{1i}^r = 0$, for any $i = 2, \dots, n, r = n + 1, \dots, 2m$ and

$$\sigma_{11}^r = \sum_{i=2}^n \sigma_{ii}^r,$$

for any $r = n + 1, \dots, 2m$. But, since $H = 0$,

$$\sigma_{11}^r = - \sum_{i=2}^n \sigma_{ii}^r,$$

for any $r = n + 1, \dots, 2m$, that is, $\sigma_{11}^r = 0$. Thus, $U \in \mathcal{N}$.

Conversely, if $U \in \mathcal{N}$, choosing a local orthonormal basis of $\mathcal{X}(\widetilde{M}(c))$ as in (3.1) with $e_1 = U$, we have that $\sigma_{1i}^r = 0$ and $\sigma_{1(2m+\alpha)}^r = 0$, for any $i = 1, \dots, n, \alpha = 1, \dots, s, r = n + 1, \dots, 2m$. Again, since $H = 0$, we obtain that $\sigma_{22}^r + \dots + \sigma_{nn}^r = 0$, for any $r = n + 1, \dots, 2m$. Then, from (3.7) we get the equality case of (3.8). Finally, from (2.1) we complete the proof. \square

Now, observe that if the equality case of (3.8) holds for all unit vector fields $U \in \mathcal{L}$, the equality case of (3.2) is true for these vector fields too. Thus, from (2.1), $NU = 0$ for any $U \in \mathcal{L}$ and M is an invariant submanifold. Then, it is easy to show that it is minimal. Consequently, making use of Theorem 3.2, $U \in \mathcal{N}$, for any $U \in \mathcal{L}$ and M is totally geodesic. The converse result is a straightforward computation. So, we have proved the following corollary of Theorem 3.2.

Corollary 3.1. *Let M be an $(n+s)$ -dimensional minimal submanifold of $\widetilde{M}(c)$, tangent to the structure vector fields. Then, the equality case of (3.8) holds for all unit vector field in \mathcal{L} if and only if M is a totally geodesic submanifold.*

Next, we are going to study the equality case of (3.2). To this end, we recall that an $(n+s)$ -dimensional submanifold of an S -manifold, tangent to the structure vector fields, is said to be a totally f -geodesic submanifold (resp., totally f -umbilical) if the distribution \mathcal{L} is totally geodesic (resp., totally umbilical), that is, if $\sigma(X, Y) = 0$ (resp., $\sigma(X, Y) = g(X, Y)V$, being $V = \frac{n+s}{n}H$), for any $X, Y \in \mathcal{L}$ [13]. Then, we can prove the following theorem.

Theorem 3.3. *Let M be an $(n+s)$ -dimensional ($n \geq 2$) submanifold of $\widetilde{M}(c)$, tangent to the structure vector fields. Then, the equality case of (3.2) holds for all unit vector field in \mathcal{L} if and only if either M is a totally f -geodesic submanifold or $n = 2$ and M is a totally f -umbilical submanifold.*

Proof. If the equality case of (3.2) is true for any unit vector field $U \in \mathcal{L}$, then, by choosing local orthonormal basis of $\mathcal{X}(\widetilde{M}(c))$ as in (3.1) and since e_1 can be

chosen to be any arbitrary unit vector fields in \mathcal{L} , from (3.6) we get

$$2\sigma_{ii}^r = \sigma_{11}^r + \dots + \sigma_{nn}^r, \quad i = 1, \dots, n,$$

$$\sigma_{ij}^r = 0, \quad i \neq j,$$

for any $r = n + 1, \dots, 2m$. Thus, we have two cases, namely either $n = 2$ or $n > 2$. In the first case, $\sigma_{11}^r = \sigma_{22}^r$, for any r and M is a totally f -umbilical submanifold, while in the second case $\sigma_{ii}^r = 0, i = 1, \dots, n$ and M is a totally f -geodesic submanifold. The converse part is a straightforward computation. \square

The above results correspond to that one proved by B.-Y. Chen in [8] for submanifolds in real space-forms. Moreover, they imply the following theorem for a slant submanifold isometrically immersed in an S -space-form.

Theorem 3.4. *Let M be an $(n + s)$ -dimensional $(n \geq 2)$ θ -slant submanifold of an S -space-form $\widetilde{M}(c)$. Then:*

(i) *For each unit vector field $U \in \mathcal{L}$, we have:*

$$(3.9) \quad 4\text{Ric}(U) \leq (n + s)^2 \|H\|^2 + (n - 1)(c + 3s) + \cos^2 \theta(3c + s).$$

(ii) *The equality case of (3.9) holds for all unit vector field in \mathcal{L} if and only if either M is a totally f -geodesic submanifold or $n = 2$ and M is a totally f -umbilical submanifold.*

Proof. For any unit vector field $U \in \mathcal{L}$, by using a local orthonormal basis of $\mathcal{X}(\widetilde{M}(c))$ as in (3.1), such that $e_1 = U$, we get from (2.8) that

$$\|TU\|^2 = \cos^2 \theta$$

and so, from (3.2) we have (3.9). The rest of the proof is similar to that of Theorem 3.3. \square

In particular, if M is an invariant submanifold (then, it is a minimal manifold too), we have:

Theorem 3.5. *Let M be an $(n + s)$ -dimensional $(n \geq 2)$ invariant submanifold of an S -space-form $\widetilde{M}(c)$ tangent to the structure vector fields. Then:*

(i) *For each unit vector field $U \in \mathcal{L}$, we have:*

$$(3.10) \quad 4\text{Ric}(U) \leq n(c + 3s) + 2(c - s).$$

(ii) *The equality case of (3.10) holds for all unit vector field in \mathcal{L} if and only if either M is a totally f -geodesic submanifold or $n = 2$ and M is a totally f -umbilical submanifold.*

Finally, if M is an anti-invariant submanifold, we obtain:

Theorem 3.6. *Let M be an $(n + s)$ -dimensional $(n \geq 2)$ anti-invariant submanifold of an S -space-form $\widetilde{M}(c)$ tangent to the structure vector fields. Then:*

(i) *For each unit vector field $U \in \mathcal{L}$, we have:*

$$(3.11) \quad 4\text{Ric}(U) \leq (n + s)^2 \|H\|^2 + (n - 1)(c + 3s).$$

- (ii) *The equality case of (3.11) holds for all unit vector field in \mathcal{L} if and only if either M is a totally f -geodesic submanifold or $n = 2$ and M is a totally f -umbilical submanifold.*

These results improve those ones proved in [14] for slant submanifolds of a Sasakian space-form (case $s = 1$).

4. The shape operator in slant submanifolds

Let M be an $(n+s)$ -dimensional θ -slant submanifold of a $(2m+s)$ -dimensional S -space-form $\widetilde{M}(c)$. Let $p \in M$ and a number

$$b > \frac{c + 3s}{4} + \frac{3(c - s)}{4} \cos^2 \theta$$

such that the sectional curvature of M , $K \geq b$ at p . Throughout this section, we consider an orthonormal basis of the tangent space $T_p(M)$,

$$\{e_1, \dots, e_n, e_{n+1}, \dots, e_{n+s}, e_{n+s+1}, \dots, e_{2m+s}\}$$

with $\{e_1, \dots, e_{n+s}\}$ being an orthonormal basis of $T_p(M)$, e_{n+s+1} parallel to the mean curvature vector at p and e_1, \dots, e_{n+s} diagonalizing the shape operator A_{n+s+1} . As above, we put $\sigma_{ij}^r = g(\sigma(e_i, e_j), e_r)$, for any $i, j = 1, \dots, n + s$ and $r = n + s + 1, \dots, 2m + s$. Then, we have

$$(4.1) \quad A_{n+s+1} = \begin{pmatrix} a_1 & 0 & \cdots & 0 \\ 0 & a_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{n+s} \end{pmatrix},$$

where $a_i = \sigma_{ii}^{n+s+1}$, $i = 1, \dots, n + s$. Moreover, if $r = n + s + 2, \dots, 2m + s$, $A_r = (\sigma_{ij}^r)$, $i, j = 1, \dots, n + s$ and so,

$$(4.2) \quad \text{trace} A_r = \sum_{i=1}^{n+s} \sigma_{ii}^r = 0,$$

since

$$\sum_{i=1}^{n+s} \sigma_{ii}^r = g((n + s)H, e_r) = 0,$$

for any $r = n + s + 2, \dots, 2m + s$ because H is parallel to e_{n+s+1} .

Now, for $i \neq j$, $i, j = 1, \dots, n + s$, we define

$$u_{ij} = a_i a_j = \sigma_{ii}^{n+s+1} \sigma_{jj}^{n+s+1} = u_{ji}$$

and from (2.5), (2.8) and by using (4.1), we obtain, for $X = Z = e_i$, $Y = W = e_j$:

$$(4.3) \quad u_{ij} \geq b - \frac{c + 3s}{4} - \frac{3(c - s)}{4} \cos^2 \theta - \sum_{r=n+s+2}^{2m+s} (\sigma_{ii}^r \sigma_{jj}^r - (\sigma_{ij}^r)^2).$$

Taking into account these formulas, we can prove the following technical lemma for u_{ij} , $i, j = 1, \dots, n + s$, $i \neq j$.

Lemma 4.1. (i) For any fixed $i \in \{1, \dots, n + s\}$:

$$\sum_{i \neq j} u_{ij} \geq (n + s - 1) \left(b - \frac{c + 3s}{4} - \frac{3(c - s)}{4} \cos^2 \theta \right) > 0.$$

(ii) $u_{ij} \neq 0$.

(iii) For distinct $i, j, k \in \{1, \dots, n + s\}$:

$$a_i^2 = \frac{u_{ij}u_{ik}}{u_{jk}}.$$

(iv) For a fixed k , $1 \leq k \leq \left\lfloor \frac{n + s}{2} \right\rfloor$:

$$\sum_{i=1}^k \sum_{j=k+1}^{n+s} u_{ij} \geq k(n - k + s) \left(b - \frac{c + 3s}{4} - \frac{3(c - s)}{4} \cos^2 \theta \right).$$

(v) $u_{ij} > 0$.

Proof. First, from (4.2) and (4.3), a straightforward computation gives (i). Now, if $u_{ij} = 0$, then $a_i = 0$ or $a_j = 0$. If $a_i = 0$, we have $u_{ik} = 0$, for any $k \in \{1, \dots, n + s\}$, $k \neq i$. Thus,

$$\sum_{j \neq i} u_{ij} = 0,$$

which contradicts to (i). Now, (iii) is direct from the definition of u_{ij} , u_{ik} and u_{jk} . Again, a straightforward computation using (4.2) and (4.3) gives (iv). Finally, if we suppose that $u_{1(n+s)} < 0$, from (iii) we get $u_{1i}u_{i(n+s)} < 0$, for $1 < i < n + s$. Without loss of generality, we may assume

$$u_{12}, \dots, u_{1k}, u_{(k+1)(n+s)}, \dots, u_{(n+s-1)(n+s)} > 0,$$

$$u_{1(k+1)}, \dots, u_{1(n+s-1)}, u_{2(n+s)}, \dots, u_{k(n+s)} < 0,$$

for some k such that:

$$\left\lfloor \frac{n + s - 1}{2} + 1 \right\rfloor \leq k \leq n + s - 1.$$

If $k = n + s - 1$, then $u_{1(n+s)} + u_{2(n+s)} + \dots + u_{(n+s-1)(n+s)} < 0$, which contradicts (i). Consequently, $k < n + s - 1$. From (iii), we obtain

$$a_{n+s}^2 = \frac{u_{i(n+s)}u_{t(n+s)}}{u_{it}} > 0,$$

where $2 \leq i \leq k$, $k + 1 \leq t \leq n + s - 1$ and so, $u_{it} < 0$. This implies that

$$\sum_{i=1}^k \sum_{t=k+1}^{n+s} u_{it} = \sum_{i=2}^k \sum_{t=k+1}^{n+s-1} u_{it} + \sum_{i=1}^k u_{i(n+s)} + \sum_{t=k+1}^{n+s} u_{1t} < 0,$$

which contradicts (iv) and completes the proof. □

In the following theorem, we establish a relationship between the shape operator associated with the mean curvature vector and the sectional curvature for slant submanifolds in an S -space-form.

Theorem 4.1. *Let M be an $(n + s)$ -dimensional θ -slant submanifold isometrically immersed in a $(2m + s)$ -dimensional S -space-form $\widetilde{M}(c)$. If at a point $p \in M$ there exists a number*

$$b > \frac{c + 3s}{4} + \frac{3(c - s)}{4} \cos^2 \theta$$

such that the sectional curvature of M , $K \geq b$ at p , then the shape operator associated with the mean curvature vector satisfies

$$A_H > \frac{n + s - 1}{n + s} \left(b - \frac{c + 3s}{4} - \frac{3(c - s)}{4} \cos^2 \theta \right) I_{n+s},$$

at p , where I_{n+s} is the identity map.

Proof. Let $p \in M$ and a number

$$b > \frac{c + 3s}{4} + \frac{3(c - s)}{4} \cos^2 \theta$$

such that the sectional curvature $K \geq b$ at p . We choose an orthonormal basis of $T_p(\widetilde{M})$ as in the beginning of the section. Then, from the above lemma, we observe that a_1, \dots, a_{n+s} have the same sign. We assume that $a_j > 0$, for any $j \in \{1, \dots, n + s\}$. Thus:

$$\begin{aligned} \sum_{j \neq i} u_{ij} &= a_i(a_1 + \dots + a_{n+s}) - a_i^2 \\ (4.4) \quad &\geq (n + s - 1) \left(b - \frac{c + 3s}{4} - \frac{3(c - s)}{4} \cos^2 \theta \right). \end{aligned}$$

From (4.1) and (4.4), we get, for any $i = 1, \dots, n + s$:

$$\begin{aligned} a_i(n + s)|H| &\geq (n + s - 1) \left(b - \frac{c + 3s}{4} - \frac{3(c - s)}{4} \cos^2 \theta \right) + a_i^2 \\ &> (n + s - 1) \left(b - \frac{c + 3s}{4} - \frac{3(c - s)}{4} \cos^2 \theta \right). \end{aligned}$$

This completes the proof. □

For the particular cases of invariant and anti-invariant submanifolds (cases $\theta = 0$ and $\theta = \pi/2$, respectively), we have the following two theorems.

Theorem 4.2. *Let M be an $(n + s)$ -dimensional invariant submanifold isometrically immersed in a $(2m + s)$ -dimensional S -space-form $\widetilde{M}(c)$ tangent to the structure vector fields. If at a point $p \in M$ there exists a number $b > c$*

such that the sectional curvature of M , $K \geq b$ at p , then the shape operator associated with the mean curvature vector satisfies at p :

$$A_H > \frac{n+s-1}{n+s}(b-c)I_{n+s}.$$

Theorem 4.3. *Let M be an $(n+s)$ -dimensional anti-invariant submanifold isometrically immersed in a $(2m+s)$ -dimensional S -space-form $\widetilde{M}(c)$ tangent to the structure vector fields. If at a point $p \in M$ there exists a number*

$$b > \frac{c+3s}{4}$$

such that the sectional curvature of M , $K \geq b$ at p , then the shape operator associated with the mean curvature vector satisfies at p :

$$A_H > \frac{n+s-1}{n+s} \left(b - \frac{c+3s}{4} \right) I_{n+s}.$$

These results should be compared with those ones proved in [14] for Sasakian space-forms (case $s = 1$).

References

- [1] D. E. Blair, *Geometry of manifolds with structural group $U(n) \times \mathcal{O}(s)$* , J. Differential Geometry **4** (1970), 155–167.
- [2] J. L. Cabrerizo, A. Carriazo, L. M. Fernández, and M. Fernández, *Slant submanifolds in Sasakian manifolds*, Glasg. Math. J. **42** (2000), no. 1, 125–138.
- [3] A. Carriazo, *New developments in slant submanifolds theory*, Applicable Mathematics in the Golden Age (Edited by J. C. Misra), 339–356, Narosa Publishing House (New Delhi, 2002).
- [4] A. Carriazo, L. M. Fernández, and M. B. Hans-Uber, *B. Y. Chen's inequality for S -space-forms: applications to slant immersions*, Indian J. Pure Appl. Math. **34** (2003), no. 9, 1287–1298.
- [5] ———, *Minimal slant submanifolds of the smallest dimension in S -manifolds*, Rev. Mat. Iberoamericana **21** (2005), no. 1, 47–66.
- [6] ———, *Some slant submanifolds of S -manifolds*, Acta Math. Hungar. **107** (2005), no. 4, 267–285.
- [7] B.-Y. Chen, *Mean curvature and shape operator of isometric immersions in real-space-forms*, Glasgow Math. J. **38** (1996), no. 1, 87–97.
- [8] ———, *Relations between Ricci curvature and shape operator for submanifolds with arbitrary codimensions*, Glasg. Math. J. **41** (1999), no. 1, 33–41.
- [9] ———, *Slant immersions*, Bull. Austral. Math. Soc. **41** (1990), no. 1, 135–147.
- [10] I. Hasegawa, Y. Okuyama, and T. Abe, *On p -th Sasakian manifolds*, J. Hokkaido Univ. Ed. Sect. II A **37** (1986), no. 1, 1–16.
- [11] M. Kobayashi and S. Tsuchiya, *Invariant submanifolds of an f -manifold with complemented frames*, Kodai Math. Sem. Rep. **24** (1972), 430–450.
- [12] A. Lotta, *Slant submanifolds in contact geometry*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **39** (1996), 183–198.
- [13] L. Ornea, *Generic CR -submanifolds in S -manifolds*, Stud. Cerc. Mat. **36** (1984), no. 5, 435–443.
- [14] M. M. Tripathi, J.-S. Kim, and S.-B. Kim, *Mean curvature and shape operator of slant immersions in a Sasakian space form*, Balkan J. Geom. Appl. **7** (2002), no. 1, 101–111.

LUIS M. FERNÁNDEZ
DEPARTMENT OF GEOMETRY AND TOPOLOGY
FACULTY OF MATHEMATICS
UNIVERSITY OF SEVILLA
APARTADO DE CORREOS 1160
41080-SEVILLA, SPAIN
E-mail address: `lmfer@us.es`

MARÍA BELÉN HANS-UBER
DEPARTMENT OF GEOMETRY AND TOPOLOGY
FACULTY OF MATHEMATICS
UNIVERSITY OF SEVILLA
APARTADO DE CORREOS 1160
41080-SEVILLA, SPAIN
E-mail address: `lmfer@us.es`

