

Status of *Haemaphysalis* tick infestation in domestic ruminants in Iran

Sadegh RAHBARI*, Sedigheh NABIAN, Parviz SHAYAN
and Hamid Reza HADDADZADEH

Department of Parasitology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

Abstract: The geographical distribution and ecological preferences of *Haemaphysalis* in domestic animals in Iran were studied 4 times a year from April 2003 to March 2005. A total of 1,622 ixodid tick specimens were collected from 3 different zones. Among them, 108 (6.7%) *Haemaphysalis* ticks, consisting of 6 species, were identified; *H. punctata* (3.4%), *H. parva* (0.5%), *H. sulcata* (0.6%), *H. choldokovskyi* (1.7%), *H. concinna* (0.06%) and *Haemaphysalis* sp. (0.6%). *H. punctata* was the most abundant species, whereas *H. concinna* was the rarest species collected in humid and sub-humid zones on cattle, sheep and goats. *H. choldokovskyi* was principally collected from sheep and goats grazed in cold mountainous areas. The infested areas consisted of Caspian Sea (Guilan, Mazandaran, Golestan, and central provinces), mountainous (Azarbaiejan, Ardebil, Kohgilouyeh, and Kordestan) and semi-desert (Khorasan, Semnan, Kerman, Sistan, and Baluchestan) zones. The Caspian Sea zone (23.6%) was the most highly infested region. The results show that various species of *Haemaphysalis* ticks infest domestic ruminants in Iran and each tick species show characteristic geographical distributions.

Key words: *Haemaphysalis*, tick, domestic ruminants, Iran

INTRODUCTION

Ticks (Ixodidae) play a significant role as a vector of pathogens of domestic animals in Iran. The major losses caused by ticks are related to transmission of babesiosis, theileriosis, and anaplasmosis in ruminants. The distribution of tick species that are able to infest animals in Iran is briefly reviewed on the basis of published records. The tick studies were started by Delpy (1936) in Iran. Later, Abbasian (1961) and Mazlum (1971) described a list of adult ticks collected from domestic animals in different regions. Filipova et

al. (1976) presented data for 642 ixodid tick larvae and nymphs taken from small mammals, chiefly rodents in different zoogeographical zones of Iran. Hoogstraal and Wassef (1979) and Hoogstraal and Valdez (1980) studied ixodid ticks parasitizing wild sheep and goat in Iran with focusing on maintaining natural foci of many hazardous diseases of humans. Rahbari (1995) published ecological aspects of various species of ticks encountering domestic animals in northwest of Iran. Razmi et al. (2002) published a list of tick species of domestic animals in northeast of Iran. However, there still seems to be a gap in our knowledge about the distribution of tick species in Iran. Therefore, the objective of this study was to determine the species of *Haemaphysalis* ticks infesting ruminants and its geographical distribution in Iran.

• Received 12 September 2006, accepted after revision 2 March 2007.

• This study was supported by the Research Council of Tehran University and Iranian Veterinary Organization, Iran.

*Corresponding author (e-mail: srahbari@ut.ac.ir)




Fig. 1. Four different geographical zones of Iran.

MATERIALS AND METHODS

Epidemiological studies on parasitic diseases of animals in Iran have been divided into 4 ecological zones (Skermann and Hillard, 1966). These localities are shown in Fig. 1. The tick specimens were collected from animals which grazed in open rangeland pastures in 4 ecological zones, which consist of Caspian Sea zone in the north, mountainous area extended from northwest to southeast, Persian Gulf lowlands, and semi-desert area in central part of Iran. Tick sampling was carried out randomly on 629 sheep, 336 goats and 151 cattle, and from the whole body of each animal. Collected ticks were counted and preserved in 70% alcohol. The speciation was done by using the

identification key of Delpy (1938) and Walker et al. (2003).

RESULTS

Tick collection was started from the late April of 2003, and continued almost to the middle of March 2005. During this period, a total of 2,170 ixodid tick specimens were collected from cattle, sheep, goats, and camels in 4 different zones. *Haemaphysalis* ticks were not found from Persian Gulf zone. From the other 3 zones, a total of 1,622 ixodid ticks, including 108 (6.7%) *Haemaphysalis* ticks, were collected. The infested areas for *Haemaphysalis* consisted of Caspian Sea areas (Guilan, Mazandaran, and Golestan provinces), mountainous areas (Azarbaiejan, Ardebil, Kohgilouyeh, and Kordestan provinces) and semi-desert zones (Khorasan, Semnan, Kerman, Sistan, and Baluchestan provinces). The Caspian Sea zone was the most highly infested region (23.6%).

Species diversity of *Haemaphysalis* ticks among ixodid tick population in 3 zones was summarized in Table 1. A total of 6 species of *Haemaphysalis* were identified on ruminants, when they were reared on pastures in Iran; *H. punctata* (3.4%), *H. parva* (0.5%), *H. sulcata* (0.6%), *H. choldokovskyi* (1.7%), *H. concinna* (0.06%), and *Haemaphysalis* sp. (0.6%). *H. punctata* was the most abundant species, whereas *H. concinna* was the rarest species collected in humid and sub-humid zones on cattle, sheep and goats. *H. choldokovskyi* was principally collected from sheep and goats grazed in cold mountainous areas.

Table 1. The distribution of *Haemaphysalis* species in 3 different geographical zones

Geographical zone	No. of ixodid ticks	<i>Haemaphysalis</i> (Total)		<i>H. sulcata</i>		<i>H. punctata</i>		<i>H. parva</i>		<i>H. concinna</i>		<i>H. choldokovskyi</i>		<i>Haemaphysalis</i> sp. ^{a)}	
		No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
I	140	33	23.6	0	0.0	22	15.7	2	1.4	1	0.7	8	5.7	0	0.0
II	608	28	4.6	0	0.0	25	4.1	4	0.7	0	0.0	14	2.3	2	0.3
III	874	47	5.4	9	1.0	8	0.9	2	0.2	0	0.0	5	0.6	7	0.8
Total	1,622	108	6.7	9	0.6	55	3.4	8	0.5	1	0.06	27	1.7	9	0.6

^{a)}Species undetermined.

DISCUSSION

The land exploitation of these last decades has dramatically reduced the diversity of Iranian environment and significantly modified the distribution and the abundance of the tick species, which strongly adapted to domestic animals. If this trend continues, it is possible to hypothesize that some new records of tick species will gradually replace most of the others. This hypothesis seems to be confirmed by continued observations on tick populations in any content. However, it is attempted to compare the obtained results in this study with the past documented reports.

The tick species, *H. choldokovskyi*, is commonly found in sheep pastured in surroundings of Caspian Sea, mountainous, and semi-desert zones in Iran. There is no report of attempts to isolate any pathogenic agent from this species, and there is also a gap of knowledge about its biological aspects. Delpy (1938) concluded that this species is distributed in high altitude territories (20°-60°E, 30°-45°N). *H. parva* is a rare species encountered in Iran, and found in Caspian Sea, mountainous, and semi-desert zones. The immature stages are frequently found on small rodents, such as social vole (Filipova et al., 1976). The adults are frequently found on sheep and goats (Mazlum, 1971). Carnivora are also the host for adults (Hoogstraal and Wassef, 1979). Hoogstraal and Valdez (1980) previously identified this species from wild sheep, and they believed that the range of this species extends to Italy and also to some parts of Libya. This tick has been known to transmit *Theileria sergenti* and Crimean-Congo hemorrhagic fever virus (Shchelkanov et al., 2005).

The species, *H. punctata*, was recorded throughout rocky mountain slopes of Caspian Sea zone and in parts of the mountainous area in wooded, brushy locations (Mazlum, 1971). But in recent years, this species has been expanding its range in most provinces in Iran. The larvae feed on small animals, such as the great gerbil, and the nymphs also feed on small mammals and birds (Filipova et al., 1976), while we found that the adults prefer cattle and sheep. This species is well known as the vector of ovian babesiosis

(Lewis and Herbert, 1980) and *Babesia major* strains (Yin et al., 1996). It has also been demonstrated that *H. punctata* carries *Rickettsia siberica* (Chen et al., 1998) and causes tick paralysis (Harwood and Maurice, 1979).

Another species, *H. sulcata*, is widely distributed in Iran. It is commonly found from northeast to southeast in semi-desert zones. Grebenyuk (1966) showed its wide distribution in India, southern USSR, and from southwestern Asia to the western Mediterranean area. Recently, Bouattour et al. (1999) reported *H. sulcata* from humid and sub-humid zones on cattle and sheep in Tunisia. The larvae feed on a variety of rodents (Filipova et al., 1976), whereas the nymphs feed on many small and large animals. Adults are usually found on larger animals, such as wild and domestic sheep, goats, cattle, horses, and camel (Hoogstraal and Valdez, 1980). This tick is known to be a vector for *Anaplasma ovis* (Walker et al., 2003).

The rarest species, *H. concinna*, is found in the east of the Caspian Sea zone to southern mountainous areas. This tick is less commonly encountered than the others. It is, however, relatively common in sheep pasture regions. This tick is highly host-specific for wood mice, but occasionally found on Persian jirds, turkestan rats, and house mice (Filipova et al., 1976). Delpy (1938) found the adult ticks on sheep, cattle and horses in mountainous areas of Caspian zone, but Mazlum (1971) emphasized that cattle could be the most important host for adult ticks. *H. concinna* was found infected with rickettsiae of spotted-fever group (Sreter-Lancz et al., 2006), but it is considered not an important vector. Examining the *H. concinna* collected in Kazakhstan revealed *Anaplasma bovis* (Shpynov et al., 2004a) and *Rickettsia hulinii* (Shpynov et al., 2004b) in these ticks. The ability of *H. concinna* to transmit *Borrelia* was determined under laboratory conditions in China (Sun and Xu, 2003). This tick was also found to be infected with the causative agents of tularemia (Khazova and Iastrebov, 2001).

ACKNOWLEDGMENTS

We wish to express our sincere thank to Dr. Allan Walker for his kind assistance. We also appreciate the

assistance received from Iranian Center of Tick and Tick-borne Diseases.

REFERENCES

- Abbasian L (1961) Records of tick (Acarina: Ixodidae) occurring in Iran and their distributional data. *Acarologia* **3**: 546-559.
- Bouattour A, Darghouth MA, Daoud A (1999) Distribution and ecology of ticks (Acari: Ixodidae) infesting livestock in Tunisia: an overview of eighth years field collections. *Parassitologia* **41(suppl. 1)**: 5-15.
- Chen M, Fan MY, Bi DZ, Zhang JZ, Huang YP (1998) Detection of *Rickettsia sibirica* in ticks and small mammals collected in three different regions of China. *Acta Virol* **42**: 61-64.
- Delpy L (1936) Note sur les Ixodides du genre *Hyalomma* (Koch). *Ann Parasitol Hum Comp* **14**: 206-245.
- Delpy L (1938) Les especes iraniennes du genre *Haemaphysalis* Koch 1844. *Ann Parasitol Hum Comp* **16**: 1-10.
- Filopova NA, Neronov VM, Farhang-Azad A (1976) Data on ixodid tick fauna (Acarina, Ixodidae) of small mammals in Iran. *Ent Obozr* **55**: No. 2 (in Russian).
- Grebenyuk Rv (1966) Ixodid ticks of Kirgizia. *Frunze Akad Nauk Kirgiz SSR Inst Biol* **328**.
- Harwood RF, Maurice TJ (1979) Tick paralysis and toxicosis. In *Entomology in human and animal health*. pp. 460-463.
- Hoogstraal H, Wassef HY (1979) *Haemaphysalis (Allophysis) kopetdaghica*: identity and discovery of each feeding stage on the wild goat in northern Iran (Ixodoidea: Ixodidae). *J Parasitol* **65**: 783-790.
- Hoogstraal H, Valdez R (1980) Ticks (Ixodoidea) from wild sheep and goats in Iran and medical and veterinary implications. *Fieldiana Zool* **6**: 1-16.
- Khazova TG, Iastrebov VK (2001) Combined focus of tick-borne encephalitis, tick-borne rickettsiosis and tularemia in the habitat of *Haemaphysalis concinna* in south central Siberia. *Zh Mikrobiol Epidemiol Immunobiol* **1**: 78-80.
- Lewis D, Herbert I (1980) A large *Babesia* of sheep from North Wales. *Vet Rec* **107**: 352-353.
- Mazlum Z (1971) Ticks of domestic animals in Iran: Geographic distribution, host relation, and seasonal activity. *J Vet Faculty, Univ Tehran, Iran* **27**: 1-32.
- Rahbari S (1995) Studies on some ecological aspects of tick fauna of West Azarbayegan, Iran. *J Appl Anim Res* **7**: 189-194.
- Razmi GR, Naghibi A, Aslani MR, Fathivand M, Dastjerdi K (2002) An epidemiological study on ovine babesiosis in the Mashhad suburb area, province of Khorasan, Iran. *Vet Parasitol* **108**: 109-115.
- Shchelkanov MIu, Kolobukhina, LV, Moskvina TM, Aushev I.D, Kartoev AA, Kelli EI, Merkulova LN, Grenkova EP, Samokhvalov EI, Petriaev VG, Serobian AG, Klimova E.A, Galkina, IV, Malyshev NA, Aristova V.A, Slavskii AA, Luk'ianova NA, Deriabin PG, Gromashevskii VL, Efremenko VI, Onishchenko GG, L'vov DK (2005) Detection of the circulation of Crimean-Congo hemorrhagic fever virus in the piedmont steppes of the North Caucasus. *Vopr³ Virusol* **50**: 9-15.
- Shpynov SN, Rudakov NV, Iastrebov VK, Leonova GN, Khazova TG, Egorova NV, Borisova ON, Preider VP, Bezrukov GV, Fedorov EG, Fedianin AP, Sherstneva MB, Turyshev AG, Gavrillov AP, Tankibaev MA, Fournier PE, Raoult D (2004a) New evidence for the detection of *Ehrlichia* and *Anaplasma* in *Ixodes* ticks in Russia and Kazakhstan. *Med Parazitol (Mosk)* **2**: 4-10.
- Shpynov SN, Rudakov NV, Iastrebov VK, Khazova TG, Fournier PE, Raoult D (2004b) Detection of *Rickettsia hulinii* in ticks of the *Haemaphysalis concinna* species in Russia. *Zh Mikrobiol Epidemiol Immunobiol* **2**: 26-29.
- Skerman KD, Hillard GG (1966) A handbook for studies of helminth parasites of ruminants. Near East Animal Health Institutes, Iran Unit, United Nations Development Programme/Special fund Handbook 2 N.E.A.H.I 1-3.
- Sreter-Lancz Z, Szell Z, Kovacs G, Egyed L, Marialigeti K, Sreter T (2006) Rickettsiae of the spotted-fever group in ixodid ticks from Hungary: identification of a new genotype ('Candidatus *Rickettsia kotlanii*'). *Ann Trop Med Parasitol* **100**: 229-236.
- Sun Y, Xu R (2003) Ability of *Ixodes persulcatus*, *Haemaphysalis concinna* and *Dermacentor silvarum* ticks to acquire and transstadially transmit *Borrelia garrinii*. *Exp Appl Acarol* **31**: 151-160.
- Walker AR, Bouattour A, Camicas JL, Estrada- Pena A, Horak IG, Latif A, Pegram RG, Preston PM (2003) Ticks of domestic animals in Africa. In *A guide to identification of species*. Bioscience Reports, U.K. 157.
- Yin H Lu W, Luo J, Zhang Q, Lu W Dou H (1996) Experiments on the transmission of *Babesia major* and *Babesia bigemina* by *Haemaphysalis punctata*. *Vet Parasitol* **67**: 89-98.