

## COMBINATORIAL PROOF FOR $e$ -POSITIVITY OF THE POSET OF RANK 1

JAEJIN LEE

ABSTRACT. Let  $P$  be a poset and  $G = G(P)$  be the incomparability graph of  $P$ . Stanley [7] defined the chromatic symmetric function  $X_{G(P)}$  which generalizes the chromatic polynomial  $\chi_G$  of  $G$ , and showed all coefficients are nonnegative in the  $e$ -expansion of  $X_{G(P)}$  for a poset  $P$  of rank 1. In this paper, we construct a sign reversing involution on the set of special rim hook  $P$ -tableaux with some conditions. It gives a combinatorial proof for  $(\mathbf{3+1})$ -free conjecture of a poset  $P$  of rank 1.

### 1. Introduction

Let  $G$  be a simple graph with  $d$  vertices. In [7], Stanley defined a homogeneous symmetric function  $X_G$  of degree  $d$  which generalizes the chromatic polynomial  $\chi_G$  of  $G$ . Let  $P$  be a poset and  $G(P)$  be the incomparability graph of  $P$ . Then the symmetric function  $X_{G(P)}$  can be expanded in terms of various symmetric function bases. In particular, if we use the elementary symmetric function basis  $\{e_\mu\}$ , we have

$$X_{G(P)} = \sum_{\mu} c_{\mu} e_{\mu}.$$

Through their work on immanants of Jacobi-Trudi matrices, Stanley and Stembridge [9] were led to the following conjecture.

---

Received September 8, 2008. Revised September 18, 2008.

2000 Mathematics Subject Classification: 05E10.

Key words and phrases: sign-reversing involution, Kostka number, chromatic symmetric function,  $P$ -tableau.

CONJECTURE 1.1 ( $(\mathbf{3+1})$ -free conjecture). *If  $P$  is a  $(\mathbf{3+1})$ -free poset,  $X_{G(P)}$  is  $e$ -positive, i.e., if*

$$X_{G(P)} = \sum_{\mu} c_{\mu} e_{\mu},$$

then all  $c_{\mu} \geq 0$ .

Using the acyclic orientation of the incomparability graph  $G(P)$  of  $P$ , Stanley [7] proved that  $(\mathbf{3+1})$ -free conjecture is true for a poset  $P$  of rank 1.

On the other hand, Egecioglu and Remmel [2] gave a combinatorial interpretation for the entries of the inverse of Kostka matrix and Chow [1] used Egecioglu and Remmel's interpretation to get a combinatorial object for  $c_{\mu}$  appeared in Conjecture 1.1.

Using Chow's combinatorial object for  $c_{\mu}$ , we construct a sign reversing involution on the set of special rim hook  $P$ -tableaux with some conditions. It gives a combinatorial proof for  $(\mathbf{3+1})$ -free conjecture of a poset  $P$  of rank 1. In Section 2 we describe basic definitions from the theory of Young tableaux. A sign reversing involution to prove the main result with an example is given in Section 3.

## 2. Definitions and combinatorial interpretation for $K_{\mu, \lambda}^{-1}$

In this section we describe some definitions necessary for later. See [3], [6] or [8] for definitions and notations not described here.

DEFINITION 2.1. A *partition*  $\lambda$  of a positive integer  $n$  is a sequence of positive integers  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{\ell})$  such that

- (i)  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{\ell} > 0$ ,
- (ii)  $\sum_{i=1}^{\ell} \lambda_i = n$ .

We write  $\lambda \vdash n$ , or  $|\lambda| = n$ . We say each term  $\lambda_i$  is a *part* of  $\lambda$  and the number of nonzero parts is called the *length* of  $\lambda$  and is written  $\ell = \ell(\lambda)$ . In addition, we will use the notation  $\lambda = (1^{m_1}, 2^{m_2}, \dots, n^{m_n})$  which means that the integer  $j$  appears  $m_j$  times in  $\lambda$ .

DEFINITION 2.2. Let  $\lambda = (\lambda_1, \dots, \lambda_{\ell})$  be a partition. The *Ferrers diagram*  $D_{\lambda}$  of  $\lambda$  is the array of cells or boxes arranged in rows and

columns,  $\lambda_1$  in the first row,  $\lambda_2$  in the second row, etc., with each row left-justified. That is,

$$D_\lambda = \{(i, j) \in \mathbf{Z}^2 \mid 1 \leq i \leq \ell(\lambda), 1 \leq j \leq \lambda_i\},$$

where we regard the elements of  $D_\lambda$  as a collection of boxes in the plane with matrix-style coordinates.

DEFINITION 2.3. If  $\lambda, \mu$  are partitions with  $D_\lambda \supseteq D_\mu$ , the *skew shape*  $D_{\lambda/\mu}$  or just  $\lambda/\mu$  is defined as the set-theoretic difference  $D_\lambda \setminus D_\mu$ . Thus

$$D_{\lambda/\mu} = \{(i, j) \in \mathbf{Z}^2 \mid 1 \leq i \leq \ell(\lambda), \mu_i < j \leq \lambda_i\}.$$

Figure 2.1 shows the Ferrers diagram  $D_\lambda$  and skew shape  $D_{\lambda/\mu}$ , respectively, when  $\lambda = (5, 4, 2, 1) \vdash 12$  and  $\mu = (2, 2, 1) \vdash 5$ .




Figure 2.1

DEFINITION 2.4. Let  $\lambda$  be a partition. A *tableau*  $T$  of shape  $\lambda$  is an assignment  $T : D_\lambda \rightarrow \mathbf{P}$  of positive integers to the cells of  $\lambda$ . The *content* of the tableau  $T$ , denoted by  $\text{content}(T)$ , is the finite nonnegative vector whose  $i$ th component is the number of entries  $i$  in  $T$ .

A tableau  $T$  of shape  $\lambda$  is said to be *column strict* if it satisfies the following two conditions:

- (i)  $T(i, j) \leq T(i, j + 1)$ , i.e., the entries increase weakly along the rows of  $\lambda$  from left to right.
- (ii)  $T(i, j) < T(i + 1, j)$ , i.e., the entries increase strictly along the columns of  $\lambda$  from top to bottom.

In Figure 2.2,  $T$  is a tableau of shape  $(5, 4, 2, 1)$  and  $S$  is a column strict tableau of shape  $(5, 4, 2, 1)$  and of content  $(3, 3, 1, 2, 2, 1)$ .




Figure 2.2

DEFINITION 2.5. For partitions  $\lambda$  and  $\mu$  such that  $|\lambda| = |\mu|$ , the *Kostka number*  $K_{\lambda,\mu}$  is the number of column strict tableaux of shape  $\lambda$  and content  $\mu$ .

If we use the reverse lexicographic order on the set of partitions of a fixed positive integer  $n$ , the *Kostka matrix*  $K = (K_{\lambda,\mu})$  becomes upper unitriangular so that  $K$  is non-singular.

DEFINITION 2.6. A *rim hook*  $H$  is a skew shape which is connected and contains no  $2 \times 2$  square of cells. The *size of*  $H$  is the number of cells it contains. The *leg length of rim hook*  $H$ ,  $\ell(H)$ , is the number of vertical edges in  $H$  when viewed as in Figure 2.3. We define the *sign* of a rim hook  $H$  to be  $\epsilon(H) = (-1)^{\ell(H)}$ .

Figure 2.3 shows the rim hook  $H$  of size 6 with  $\ell(H) = 2$  and  $\epsilon(H) = (-1)^2 = 1$ .




Figure 2.3

DEFINITION 2.7. A *rim hook tableau*  $T$  of shape  $\lambda$  is a partition of the diagram of  $\lambda$  into rim hooks. The *type* of  $T$  is  $\text{type}(T) = (1^{m_1}, 2^{m_2}, \dots, n^{m_n})$  where  $m_k$  is the number of rim hooks in  $T$  of size  $k$ . We now define the *sign* of a rim hook tableau  $T$  as

$$\epsilon(T) = \prod_{H \in T} \epsilon(H).$$

A rim hook tableau  $S$  is called *special* if each of the rim hooks contains a cell from the first column of  $\lambda$ . We use nodes for the Ferrers diagram

and connect them if they are adjacent in the same rim hook as  $S$  in Figure 2.4.




Figure 2.4

In Figure 2.4,  $T$  is a rim hook tableau of shape  $(5, 4, 2, 1)$ ,  $\text{type}(T) = (1^2, 2, 4^2)$  and  $\epsilon(T) = (-1)^1 \cdot (-1)^1 \cdot (-1)^0 \cdot (-1)^0 \cdot (-1)^0 = 1$ , while  $S$  is a special rim hook tableau with shape  $(5, 3, 2, 1, 1)$ ,  $\text{type}(S) = (2, 4, 6)$  and  $\epsilon(S) = (-1)^0 \cdot (-1)^1 \cdot (-1)^2 = -1$ .

We can now state Egecioglu and Remmel’s interpretation for the entries of the inverse of Kostka matrix.

**THEOREM 2.8** (Egecioglu and Remmel[2]). *The entries of the inverse Kostka matrix are given by*

$$K_{\mu,\lambda}^{-1} = \sum_S \epsilon(S)$$

where the sum is over all special rim hook tableaux  $S$  with shape  $\lambda$  and type  $\mu$ . □

### 3. A sign reversing involution

We begin with Stanley’s chromatic symmetric functions in this section.

**DEFINITION 3.1.** Let  $G = G(V, E)$  be a graph with a finite set of vertices  $V$  and edges  $E$ . A *proper coloring* of  $G$  is a function  $\kappa : V \rightarrow \mathbb{P}$  such that  $uv \in E$  implies  $\kappa(u) \neq \kappa(v)$ . Now consider a countably infinite

set of variables  $\mathbf{x} = \{x_1, x_2, \dots\}$ . The *chromatic symmetric function*  $X_G$  associated with a graph  $G$  is a formal power series

$$X_G = X_G(\mathbf{x}) = \sum_{\kappa:V \rightarrow \mathbb{P}} x_{\kappa(v_1)}x_{\kappa(v_2)} \cdots x_{\kappa(v_n)}$$

where  $\kappa$  is a proper coloring.

Note that if one sets  $x_1 = x_2 = \dots = x_n = 1$  and  $x_i = 0$  for  $i > n$ , denoted  $\mathbf{x} = 1^n$ , then  $X_G$  reduces to the number of proper colorings of  $G$  from a set with  $n$  elements. So under this substitution,  $X_G(1^n) = \chi_G(n)$  where  $\chi_G(n)$  is the chromatic polynomial of Whitney [10]. Also, because permuting the colors of a proper coloring keeps the coloring proper,  $X_G(\mathbf{x})$  is a symmetric function in  $\mathbf{x}$  over the rationals. In [7], Stanley derived many interesting properties of the chromatic symmetric function  $X_G(\mathbf{x})$  some of which generalize those of the chromatic polynomial.

DEFINITION 3.2. Let  $(P, \leq)$  be a finite partially ordered set (poset). We say that  $P$  is  $(\mathbf{a}+\mathbf{b})$ -free if it contains no induced subposet isomorphic to a disjoint union of an  $a$ -element chain and a  $b$ -element chain. Also, given any poset  $P$ , *incomparability graph*  $G(P)$  of  $P$  is a graph having vertices  $V = P$  and an edge between  $u$  and  $v$  in  $G(P)$  if and only if  $u$  and  $v$  are incomparable in  $P$ .

Figure 3.1 shows a poset  $P$  and its incomparability graph  $G(P)$ .




Figure 3.1

Although  $(\mathbf{3}+\mathbf{1})$ -free conjecture introduced in Section 1 still remains open, a weak result proved by Gasharov [4]. He gave a combinatorial interpretation to the coefficients in the  $s$ -expansion of  $X_{G(P)}$  and proved that if  $P$  is  $(\mathbf{3}+\mathbf{1})$ -free then  $X_{G(P)}$  is  $s$ -positive, where  $s_\lambda$  is the Schur function corresponding to  $\lambda$ .

DEFINITION 3.3. Let  $P$  be a poset. A  $P$ -tableau  $T$  of shape  $\lambda$  is a bijection  $D_\lambda \rightarrow P$  such that for all  $(i, j) \in \lambda$ :

- (i)  $T_{i,j} < T_{i+1,j}$ , and
- (ii)  $T_{i,j} \not> T_{i,j+1}$ ,

where a condition is considered vacuously true if subscripts refer to a cell outside of  $\lambda$ . We denote the number of  $P$ -tableaux of shape  $\lambda$  by  $f_P^\lambda$ .

Note that when  $P$  is a chain, then a  $P$ -tableau is just a standard Young tableau and  $f_P^\lambda = f^\lambda$ . Figure 3.2 shows all  $P$ -tableaux of shape  $\lambda = (3, 1)$  when  $P$  is a poset given in Figure 3.1.

$a$	$b$	$d$	$b$	$a$	$d$	$b$	$d$	$a$	$b$	$a$	$c$
$c$			$c$			$c$			$d$		

Figure 3.2

Using  $P$ -tableaux, Gasharov proved the following result which immediately implies  $s$ -positivity of  $X_{G(P)}$ , where  $P$  is a  $(\mathbf{3+1})$ -free poset.

**THEOREM 3.4** (Gasharov [4]). *If  $P$  is  $(\mathbf{3+1})$ -free then*

$$(1) \quad X_{G(P)} = \sum_{\lambda} f_P^\lambda s_{\lambda'}$$

where  $\lambda'$  is the conjugate of  $\lambda$ . □

Chow [1] pointed out that (1) could be combined with Egecioglu and Remmel’s result to obtain a combinatorial interpretation of the coefficients  $c_\mu$  in Conjecture 1.1. First note that the change of basis matrix between the Schur and elementary symmetric functions is

$$s_{\lambda'} = \sum_{\mu} K_{\mu,\lambda}^{-1} e_{\mu}$$

Combining this with (1) we get

$$X_{G(P)} = \sum_{\lambda,\mu} K_{\mu,\lambda}^{-1} f_P^\lambda e_{\mu}.$$

Since the  $e_{\mu}$  are a basis, we have

$$c_{\mu} = \sum_{\lambda} K_{\mu,\lambda}^{-1} f_P^\lambda.$$

Finally we apply Theorem 2.8 to get the desired interpretation.

COROLLARY 3.5 (Chow [1]). *Let  $P$  be a finite poset and let*

$$X_{G(P)} = \sum_{\mu} c_{\mu} e_{\mu}.$$

*Then, the coefficients  $c_{\mu}$  satisfy*

$$c_{\mu} = \sum_{(S,T)} \epsilon(S)$$

*where the sum is over all pairs of a special rim hook tableau  $S$  of type  $\mu$  and a  $P$ -tableau  $T$  with the same shape as  $S$ .  $\square$*

Note that a column of a  $P$ -tableau  $T$  must be a chain in  $P$  and the number of rim hooks in  $S$  is at most the length of its first column because they are special. So the previous corollary implies that  $c_{\mu} = 0$  whenever  $\mu$  has more parts than the height of  $P$ ,  $h(P)$  (which is defined as the number of elements in the longest chain of  $P$ ).

To present pairs  $(S, T)$  described in Corollary 3.5 economically, we will combine each pair  $(S, T)$  into a single tableau  $S_T$ , called *special rim hook  $P$ -tableau*, with elements in the same places as in  $T$  and edges between pairs of elements which are adjacent in a hook of  $S$ . See Figure 3.3 for an example of special rim hook  $P$ -tableau.




Figure 3.3

Using special rim hook  $P$ -tableaux Corollary 3.5 can be rewritten as follows.

COROLLARY 3.6. *Let  $P$  be a finite poset. Then the coefficients  $c_{\mu}$  in the  $e$ -expansion of  $X_G(P)$  are*

$$c_{\mu} = \sum_S \epsilon(S)$$



where the sum is over all special rim hook  $P$ -tableaux  $S$  of type  $\mu$ .  $\square$

We can now state the main result and give a sign reversing involution to prove it.

**THEOREM 3.7.** *Let  $P$  be a poset with  $n$  elements of rank 1. Then*

$$\sum_S \epsilon(S)$$

is non-negative, where the sum is over all special rim hook  $P$ -tableaux  $S$  of type  $\mu \vdash n$ .

*Proof.* Let  $\mu$  be a fixed partition of  $n$  and  $\Gamma_\mu$  be the set of all special rim hook  $P$ -tableaux of type  $\mu$ . We divide the set  $\Gamma_\mu$  into two disjoint subsets  $\Gamma_\mu^+$  and  $\Gamma_\mu^-$  as follows.

$$\begin{aligned} \Gamma_\mu^+ &= \{ S \in \Gamma_\mu \mid \epsilon(S) = 1 \} \\ \Gamma_\mu^- &= \{ S \in \Gamma_\mu \mid \epsilon(S) = -1 \} \end{aligned}$$

Note that  $P$  cannot have a chain of three elements and a column of a  $P$ -tableau  $T$  in  $\Gamma_\mu$  must be a chain in  $P$ . This fact implies that the shape of  $T$  has at most two rows, and the number of rim hooks in  $T$  is at most two because it is special. This means that either  $\mu = (n^1)$  or  $\mu = (r^1, s^1)$  with  $r \geq s$ .

Suppose first  $\mu = (n^1)$ . Since  $T$  contains only one rim hook,  $T$  is a special rim hook  $P$ -tableaux of form

$$a_1 - a_2 - \cdots - a_{n-1} - a_n \quad \text{or} \quad \begin{array}{c} b_1 - b_2 - \cdots - b_{n-1} \\ | \\ b_n \end{array}$$

Define

$$I(a_1 - a_2 - \cdots - a_{n-1} - a_n) = \begin{cases} a_1 - a_2 - \cdots - a_{n-1} - a_n & \text{if } a_1 \not\prec a_n \\ \begin{array}{c} a_1 - a_2 - \cdots - a_{n-1} \\ | \\ a_n \end{array} & \text{otherwise} \end{cases}$$

and

$$I \left( \begin{array}{c} b_1 - b_2 - \cdots - b_{n-1} \\ | \\ b_n \end{array} \right) = b_1 - b_2 - \cdots - b_{n-1} - b_n$$

If  $b_{n-1} > b_n$  in the above,  $\{b_1, b_n, b_{n-1}\}$  forms a chain of three elements in  $P$ . Hence we have  $b_{n-1} \not> b_n$  and  $I$  is well-defined on  $\Gamma_\mu$ .

Suppose now  $\mu = (r^1, s^1)$  with  $r \geq s$ . Since there are two rim hooks in each  $P$ -tableau in  $\Gamma_\mu$ , such tableau is of form

$$T_1 = \begin{array}{cccccccc} a_1 - a_2 - \cdots - a_{s-1} - a_s - a_{s+1} - \cdots - a_r \\ b_1 - b_2 - \cdots - b_{s-1} - b_s \end{array}$$

or

$$T_2 = \begin{array}{cccccccc} c_1 - c_2 - \cdots - c_{s-1} - c_s & d_{s+2} & - \cdots - d_r \\ & | \\ d_1 - d_2 - \cdots - d_{s-1} - d_s - d_{s+1} \end{array}$$

Define

$$I(T_1) = \begin{cases} T_1 & \text{if } a_{s+1} \not> a_r \\ T_3 & \text{otherwise} \end{cases}$$

and

$$I(T_2) = T_4$$

where

$$T_3 = \begin{array}{cccccccc} a_1 - a_2 - \cdots - a_{s-1} - a_s & a_{s+1} - \cdots - a_{r-1} \\ & | \\ b_1 - b_2 - \cdots - b_{s-1} - b_s - a_r \end{array}$$

and

$$T_4 = \begin{array}{cccccccc} c_1 - c_2 - \cdots - c_s - d_{s+2} - \cdots - d_r - d_{s+1} \\ d_1 - d_2 - \cdots - d_s \end{array}$$

If  $d_r > d_{s+1}$  or  $b_s > a_r > a_{s+1}$  in the above,  $P$  contains a chain of three elements  $\{d_{s+2}, d_{s+1}, d_r\}$  or  $\{a_{s+1}, a_r, b_s\}$ . Thus we have  $d_r \not> d_{s+1}$ ,  $b_s \not> a_r$  and  $I$  is well-defined.

In either case, we can check easily that  $I$  is a sign reversing involution on  $\Gamma_\mu$ , i.e.,  $I \circ I = 1_{\Gamma_\mu}$  and

$$\epsilon(I(S)) = \begin{cases} 1 & \text{if } S \in \Gamma_\mu^-, \\ -1 & \text{if } S \in \Gamma_\mu^+ \text{ and } I(S) \neq S \\ 1 & \text{if } S \in \Gamma_\mu^+ \text{ and } I(S) = S. \end{cases}$$

Class	shape	type	sign	# of special rim hook $P$ -tableaux
I	(5)	(5)	1	42
II	(4, 1)	(5)	-1	12
III	(4, 1)	(4, 1)	1	12
IV	(3, 2)	(3, 2)	1	6
V	(3, 2)	(4, 1)	-1	6

TABLE 1

Using the above involution  $I$ , we finally have

$$\sum_S \epsilon(S) = \sum_{S \in \Gamma_\mu} \epsilon(S) = \sum_{S \in \Gamma_\mu^+, I(S)=S} \epsilon(S) \geq 0$$

which immediately implies our theorem. □

Combining Corollary 3.6 and Theorem 3.7, we get the following facts.

**COROLLARY 3.8** (Stanley[7]). *Let  $P$  be a finite poset of rank 1. For any partition  $\mu$ , the coefficient  $c_\mu$  of  $e_\mu$  in the  $e$ -expansion of  $X_{G(P)}$  is non-negative.* □

**EXAMPLE 3.9.** Consider the poset  $P$  as in Figure 3.4.




Figure 3.4

Then there are 78 special rim hook  $P$ -tableaux. Table 1 shows all possible shapes and types of special rim hook  $P$ -tableaux, and the number of special rim hook  $P$ -tableaux with given shape and type. Examples of a special rim hook  $P$ -tableaux contained in each class of Table 1 are given in Figure 3.5.

$$\begin{array}{ll}
 \text{Class I:} & a - b - c - d - e \\
 \\
 \text{Class II:} & \begin{array}{c} a - b - c - e \\ | \\ d \end{array} \\
 \\
 \text{Class III:} & \begin{array}{c} a - b - c - e \\ d \end{array} \\
 \\
 \text{Class IV:} & \begin{array}{c} a - b - c \\ d - e \end{array} \\
 \\
 \text{Class V:} & \begin{array}{c} a \quad b - c \\ d - e \quad | \end{array}
 \end{array}$$

Figure 3.5

Each special rim hook  $P$ -tableaux in Class II is matched to just one of tableaux in Class I, and each special rim hook  $P$ -tableaux in Class V is matched to one in Class III as follows;

$$\begin{array}{ccc}
 \begin{array}{c} a - b - c - e \\ | \\ d \end{array} & \iff & a - b - c - e - d \\
 \\
 \begin{array}{c} a \quad b - c \\ | \\ d - e \end{array} & \iff & \begin{array}{c} a - b - c - e \\ d \end{array}
 \end{array}$$

Figure 3.6

30 unmatched tableaux in Class I, 6 unmatched tableaux in Class III and 6 unmatched tableaux in Class IV are fixed by the involution  $I$  described in Theorem 3.7. Hence, we have

$$X_{G(P)} = 30e_{(5)} + 6e_{(4,1)} + 6e_{(3,2)}.$$

## References

- [1] T. Chow, A note on a combinatorial interpretation of the  $e$ -coefficients of the chromatic symmetric function, preprint (1997), 9 pp, arXiv math.CO/9712230.
- [2] Ö. Eğecioğlu and J. Remmel, A combinatorial interpretation of the inverse Kostka matrix, *Linear and Multilinear Algebra* **26** (1990), 59–84.
- [3] W. Fulton, “Young Tableaux,” London Mathematical Society Student Texts 35, Cambridge University Press, Cambridge, 1999.

- [4] V. Gasharov, Incomparability graphs of  $(\mathbf{3+1})$ -free posets are  $s$ -positive, *Discrete Math.* **157** (1996), 193–197.
- [5] I. G. Macdonald, “Symmetric functions and Hall polynomials,” 2nd edition, Oxford University Press, Oxford, 1995.
- [6] B. Sagan, “The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions,” 2nd edition, Springer-Verlag, New York, 2001.
- [7] R. P. Stanley, A symmetric function generalization of the chromatic polynomial of a graph, *Advances in Math.* **111** (1995), 166–194.
- [8] R. P. Stanley, “Enumerative Combinatorics, Volume 2,” Cambridge University Press, Cambridge, 1999.
- [9] R. P. Stanley and J. Stembridge, On immanants of Jacobi-Trudi matrices and permutations with restricted position, *J. Combin. Theory Ser. A* **62** (1993), 261–279.
- [10] H. Whitney, A logical expansion in mathematics, *Bull. Amer. Math. Soc.* **38** (1932), 572–579.

Department of Mathematics  
Hallym University  
Chunchon, Korea 200–702  
*E-mail*: jjlee@hallym.ac.kr