

TUBES OF WEINGARTEN TYPES IN A EUCLIDEAN 3-SPACE

JIN SUK RO* AND DAE WON YOON**

ABSTRACT. In this paper, we study a tube in a Euclidean 3-space satisfying some equation in terms of the Gaussian curvature, the mean curvature and the second Gaussian curvature.

1. Introduction

Let f and g be smooth functions on a surface M in a Euclidean 3-space \mathbb{E}^3 . The Jacobi function $\Phi(f, g)$ formed with f, g is defined by $\Phi(f, g) = \det \begin{pmatrix} f_s & f_t \\ g_s & g_t \end{pmatrix}$, where $f_s = \frac{\partial f}{\partial s}$ and $f_t = \frac{\partial f}{\partial t}$. In particular, a surface satisfying the Jacobi equation $\Phi(K, H) = 0$ with respect to the Gaussian curvature K and the mean curvature H on a surface M is called a *Weingarten surface* or a *W-surface*. Also, if a surface satisfies a linear equation with respect to K and H , that is, $aK + bH = c$ ($a, b, c \in \mathbb{R}, (a, b, c) \neq (0, 0, 0)$), then it is said to be a *linear Weingarten surface* or a *LW-surface*.

When the constant $b = 0$, a linear Weingarten surface M reduces to a surface with constant Gaussian curvature. When the constant $a = 0$, a linear Weingarten surface M reduces to a surface with constant mean curvature. In such a sense, the linear Weingarten surfaces can be regarded as a natural generalization of surfaces with constant Gaussian curvature or with constant mean curvature.

Several geometers ([3, 4, 6, 11, 12, 13]) have studied *W-surfaces* and *LW-surfaces* and obtained many interesting results.

Received April 29, 2009; Accepted August 14, 2009.

2000 Mathematics Subject Classification: Primary 53A05, 56B34; Secondary 53B25.

Key words and phrases: tube, Gaussian curvature, mean curvature, second Gaussian curvature, circular helix.

Correspondence should be addressed to Dae Won Yoon, dwyoon@gnu.ac.kr.

If the second fundamental form II of a surface M in \mathbb{E}^3 is non-degenerate, then it is regarded as a new (pseudo-) Riemannian metric. Therefore, the Gaussian curvature K_{II} of non-degenerate second fundamental form II can be defined formally on the Riemannian or pseudo-Riemannian manifold (M, II) . We call the curvature K_{II} the *second Gaussian curvature* on M .

For a pair (X, Y) , $X \neq Y$, of the curvatures K, H and K_{II} of M in \mathbb{E}^3 , if M satisfies $\Phi(X, Y) = 0$ and $aX + bY = c$, then it is said to be a (X, Y) -Weingarten surface and (X, Y) -linear Weingarten surface, respectively.

For study of these surfaces, W. Kühnel ([11]) and G. Stamou ([13]) investigate ruled (X, Y) -Weingarten surface in a Euclidean 3-space \mathbb{E}^3 . Also, C. Baikoussis and Th. Koufogiorgos ([1]) studied helicoidal (H, K_{II}) -Weingarten surfaces. F. Dillen and W. Kühnel ([3]) and F. Dillen and W. Sodsiri ([4, 5]) gave a classification of ruled (X, Y) -Weingarten surface in a Minkowski 3-space \mathbb{E}_1^3 , where $X, Y \in \{K, H, K_{II}\}$. D. Koutroufiotis ([10]) and Th. Koufogiorgos and T. Hasanis ([9]) investigate closed ovaloid (X, Y) -linear Weingarten surface in \mathbb{E}^3 . D. W. Yoon ([14]) and D. E. Blair and Th. Koufogiorgos ([2]) classified ruled (X, Y) -linear Weingarten surface in \mathbb{E}^3 .

Following the Jacobi equation and the linear equation with respect to the Gaussian curvature K , the mean curvature H and the second Gaussian curvature K_{II} , an interesting geometric question is raised:

“Classify all surfaces in a Euclidean 3-space and a Minkowski 3-space satisfying the conditions

$$(1.1) \quad \Phi(X, Y) = 0,$$

$$(1.2) \quad aX + bY = c,$$

where $X, Y \in \{K, H, K_{II}\}$, $X \neq Y$ and $(a, b, c) \neq (0, 0, 0)$.”

In this paper, we would like to contribute the solution of the above question, by studying this question for a tube in a Euclidean 3-space \mathbb{E}^3 .

Throughout this paper, we assume that all objects are smooth and all surfaces are connected unless stated otherwise.

2. Preliminaries

We denote a surface M in \mathbb{E}^3 by

$$x(s, t) = (x_1(s, t), x_2(s, t), x_3(s, t)).$$

Let U be the standard unit normal vector field on a surface M defined by $U = \frac{x_s \times x_t}{\|x_s \times x_t\|}$, where $x_s = \frac{\partial x(s,t)}{\partial s}$. Then the first fundamental form I and the second fundamental form II of a surface M are defined by, respectively

$$I = E ds^2 + 2F ds dt + G dt^2,$$

$$II = e ds^2 + 2f ds dt + g dt^2,$$

where

$$E = \langle x_s, x_s \rangle, \quad F = \langle x_s, x_t \rangle, \quad G = \langle x_t, x_t \rangle,$$

$$e = \langle x_{ss}, U \rangle, \quad f = \langle x_{st}, U \rangle, \quad g = \langle x_{tt}, U \rangle.$$

On the other hand, the Gaussian curvature K and the mean curvature H are given by, respectively

$$K = \frac{eg - f^2}{EG - F^2},$$

$$H = \frac{1}{2} \frac{Eg - 2Ff + Ge}{EG - F^2}.$$

From Brioschi's formula in a Euclidean 3-space, we are able to compute K_{II} of a surface by replacing the components of the first fundamental form E, F, G by the components of the second fundamental form e, f, g respectively in Brioschi's formula. Consequently, the second Gaussian curvature K_{II} of a surface is defined by (cf. [2].)

$$(2.1) \quad K_{II} = \frac{1}{(eg - f^2)^2} \left\{ \begin{array}{c} \left| \begin{array}{cc} -\frac{1}{2}e_{tt} + f_{st} - \frac{1}{2}g_{ss} & \frac{1}{2}e_s \quad f_s - \frac{1}{2}e_t \\ f_t - \frac{1}{2}g_s & e \quad f \end{array} \right| \\ \frac{1}{2}g_t & f \quad g \end{array} \right| \\ - \left| \begin{array}{ccc} 0 & \frac{1}{2}e_t & \frac{1}{2}g_s \\ \frac{1}{2}e_t & e & f \\ \frac{1}{2}g_s & f & g \end{array} \right| \end{array} \right\}.$$

REMARK 2.1. (cf. [1]) It is well known that a minimal surface has vanishing second Gaussian curvature but that a surface with vanishing second Gaussian curvature need not be minimal.

3. Tubes of Weingarten types

Let $\gamma : (a, b) \rightarrow \mathbb{E}^3$ be a smooth unit speed curve of finite length which is topologically imbedded in \mathbb{E}^3 . The total space N_γ of the normal bundle of $\gamma((a, b))$ in \mathbb{E}^3 is naturally diffeomorphic to the direct product $(a, b) \times \mathbb{E}^2$ via the translation along γ with respect to the induced normal

connection. For a sufficiently small $r > 0$, the tube of radius r about the curve γ is the set :

$$T_r(\gamma) = \{\exp_{\gamma(t)} v \mid v \in N_{\gamma(t)}, \|v\| = r, a < t < b\}.$$

For a sufficiently small $r > 0$, the tube $T_r(\gamma)$ is a smooth surface in \mathbb{E}^3 . Let $\mathbf{t}, \mathbf{n}, \mathbf{b}$ denote the Frenet frame of the unit speed curve $\gamma = \gamma(t)$. Then the position vector of $T_r(\gamma)$ can be expressed as

$$(3.1) \quad x = x(t, \theta) = \gamma(t) + r(\cos \theta \mathbf{n}(t) + \sin \theta \mathbf{b}(t)).$$

We denote by κ, τ the curvature and the torsion of the curve γ . Then Frenet formula of $\gamma(t)$ is defined by

$$(3.2) \quad \begin{cases} \mathbf{t}' &= \kappa \mathbf{n}, \\ \mathbf{n}' &= -\kappa \mathbf{t} + \tau \mathbf{b}, \\ \mathbf{b}' &= -\tau \mathbf{n}, \end{cases}$$

where the prime denotes the differentiation with respect to t . Furthermore, we have the natural frame $\{x_s, x_t\}$ given by

$$\begin{aligned} x_t &= (1 - r\kappa \cos \theta) \mathbf{t} - r\tau \sin \theta \mathbf{n} + r\tau \cos \theta \mathbf{b} \\ &= \alpha \mathbf{t} + r\tau \mathbf{v}, \\ x_\theta &= -r \sin \theta \mathbf{n} + r \cos \theta \mathbf{b} = r\mathbf{v}, \end{aligned}$$

where we put $\alpha = 1 - r\kappa(t) \cos \theta$ and $\mathbf{v} = -\sin \theta \mathbf{n}(t) + \cos \theta \mathbf{b}(t)$. From which the components of the first fundamental form are

$$(3.3) \quad E = \alpha^2 + r^2 \tau^2, \quad F = r^2 \tau, \quad G = r^2.$$

On the other hand, the unit normal vector field U is obtained by

$$U = \frac{1}{\|x_t \times x_\theta\|} (x_t \times x_\theta) = -\cos \theta \mathbf{n} - \sin \theta \mathbf{b},$$

from this, the components of the second fundamental form of x are given by

$$e = r\tau^2 - \kappa\alpha \cos \theta, \quad f = r\tau, \quad g = r.$$

If the second fundamental form is non-degenerate, $eg - f^2 \neq 0$, that is, κ, α and $\cos \theta$ are nowhere vanishing. In this case, we can define formally the second Gaussian curvature K_{II} on $T_r(\gamma)$.

On the other hand, the Gauss curvature K , the mean curvature H and the second Gaussian curvature K_{II} are given by, respectively

$$(3.4) \quad K = -\frac{1}{r\alpha} \kappa \cos \theta,$$

$$(3.5) \quad H = \frac{1}{2r\alpha}(1 - 2r\kappa \cos \theta),$$

$$(3.6) \quad K_{II} = \frac{1}{4r\alpha^2 \cos^2 \theta}(4r^2\kappa^2 \cos^4 \theta - 6r\kappa \cos^3 \theta + \cos^2 \theta + 1).$$

Differentiating K, H and K_{II} with respect to t and θ , we get

$$(3.7) \quad K_t = -\frac{\kappa' \cos \theta}{r\alpha^2}, \quad K_\theta = \frac{\kappa \sin \theta}{r\alpha^2},$$

$$(3.8) \quad H_t = -\frac{\kappa' \cos \theta}{2r\alpha^2}, \quad H_\theta = \frac{\kappa \sin \theta}{2\alpha^2},$$

$$(3.9) \quad \begin{aligned} (K_{II})_t &= \frac{1}{4r\alpha^4 \cos \theta}(-2r^3\kappa^2\kappa' \cos^4 \theta + 6r^2\kappa\kappa' \cos^3 \theta \\ &\quad - 4r\kappa' \cos^2 \theta - 2r^2\kappa\kappa' \cos \theta + 2r\kappa'), \\ (K_{II})_\theta &= \frac{1}{4r\alpha^4 \cos^4 \theta}(2r^3\kappa^3 \cos^6 \theta \sin \theta - 6r^2\kappa^2 \cos^5 \theta \sin \theta \\ &\quad + 4r\kappa \cos^4 \theta \sin \theta + 4r^2\kappa^2 \cos^3 \theta \sin \theta \\ &\quad - 6r\kappa \cos^2 \theta \sin \theta + 2 \cos \theta \sin \theta). \end{aligned}$$

Now, we investigate a tube $T_r(\gamma)$ in \mathbb{E}^3 satisfying the Jacobi equation $\Phi(X, Y) = 0$.

By using (3.7) and (3.8), $T_r(\gamma)$ satisfies identically the Jacobi equation $\Phi(K, H) = K_t H_\theta - K_\theta H_t = 0$. Therefore, $T_r(\gamma)$ is a Weingarten surface.

We consider a tube $T_r(\gamma)$ with non-degenerate second fundamental form in \mathbb{E}^3 satisfying the Jacobi equation

$$(3.10) \quad \Phi(K, K_{II}) = K_t(K_{II})_\theta - K_\theta(K_{II})_t = 0$$

with respect to the Gaussian curvature K and the second Gaussian curvature K_{II} . Then, by (3.7) and (3.9) equation (3.10) becomes

$$r^2\kappa^2\kappa' \cos^2 \theta + 4r\kappa\kappa' \cos \theta \sin \theta + \kappa' \sin \theta = 0.$$

Since this polynomial is equal to zero for every θ , all its coefficients must be zero. Therefore, we conclude that $\kappa' = 0$.

We suppose that a tube $T_r(\gamma)$ with non-degenerate second fundamental form in \mathbb{E}^3 is (H, K_{II}) -Weingarten surface. Then it satisfies the equation

$$(3.11) \quad H_t(K_{II})_\theta - H_\theta(K_{II})_t = 0,$$

which implies

$$(3.12) \quad r^2\kappa^2\kappa' \cos^2 \theta \sin \theta - 2r\kappa\kappa' \cos \theta \sin \theta + \kappa' \sin \theta = 0.$$

From (3.12) we can obtain $\kappa' = 0$.

Consequently, we have the following theorems:

THEOREM 3.1. *A tube in a Euclidean 3-space is a Weingarten surface.*

THEOREM 3.2. *Let $(X, Y) \in \{(K, K_{II}), (H, K_{II})\}$ and let $T_r(\gamma)$ be a tube defined by (3.1) in a Euclidean 3-space with non-degenerate second fundamental form. If $T_r(\gamma)$ is a (X, Y) -Weingarten surface, then the curvature of γ is a non-zero constant.*

THEOREM 3.3. *Under the same hypothesis and notation as in Theorem 3.2, if γ has non-zero constant torsion, then $T_r(\gamma)$ is generated by a circular helix γ .*

Finally, we study a tube $T_r(\gamma)$ in \mathbb{E}^3 satisfying a linear equation $aX + bY = c$.

First of all, we suppose that a tube $T_r(\gamma)$ in \mathbb{E}^3 is a linear Weingarten surface, that is, it satisfies the equation

$$(3.13) \quad aK + bH = c.$$

Then, by (3.4) and (3.5) we have

$$(2a\kappa + 2br\kappa - 2cr^2\kappa) \cos \theta - b + 2rc = 0.$$

Since $\cos \theta$ and 1 are linearly independent, we get

$$2a\kappa + 2br\kappa - 2cr^2\kappa = 0, \quad b = 2rc,$$

which imply

$$\kappa(a + cr^2) = 0.$$

If $a + cr^2 \neq 0$, then $\kappa = 0$. Thus, $T_r(\gamma)$ is an open part of a circular cylinder.

Next, suppose that a tube $T_r(\gamma)$ with non-degenerate second fundamental form in \mathbb{E}^3 satisfies the equation

$$(3.14) \quad aK + bK_{II} = c.$$

By (3.4) and (3.6), equation (3.14) becomes

$$(4ar\kappa^2 + 4br^2\kappa^2 - 4cr^3\kappa^2) \cos^4 \theta - (4a\kappa + 6br\kappa - 8cr^2\kappa) \cos^3 \theta + (b - 4cr) \cos^2 \theta + b = 0.$$

Since the identity holds for every θ , all the coefficients must be zero. Therefore, we have

$$\begin{cases} 4ar\kappa^2 + 4br^2\kappa^2 - 4cr^3\kappa^2 = 0, \\ 4a\kappa + 6br\kappa - 8cr^2\kappa, \\ b - 4cr = 0, \\ b = 0. \end{cases}$$

Thus, we get $b = 0$, $c = 0$ and $\kappa = 0$. In this case, the second fundamental form of $T_r(\gamma)$ is degenerate.

Suppose that a tube $T_r(\gamma)$ with non-degenerate second fundamental form in \mathbb{E}^3 satisfies the equation

$$(3.15) \quad aH + bK_{II} = c.$$

By (3.5), (3.6) and (3.15), we have

$$\begin{aligned} & (4ar^2\kappa^2 + 4br^2\kappa^2 - 4cr^3\kappa^2) \cos^4 \theta - (6ar\kappa + 6br\kappa - 8cr^2\kappa) \cos^3 \theta \\ & + (2a + b - 4cr) \cos^2 \theta + b = 0, \end{aligned}$$

from which we can obtain $b = 0$ and $\kappa = 0$.

Consequently, we have the following theorems:

THEOREM 3.4. *Let $T_r(\gamma)$ be a tube satisfying the linear equation $aK + bH = c$. If $a + br \neq 0$, then it is an open part of a circular cylinder.*

THEOREM 3.5. *Let $(X, Y) \in \{(K, K_{II}), (H, K_{II})\}$. Then there are no (X, Y) -linear Weingarten tubes in a Euclidean 3-space.*

Acknowledgments. The authors would like to thank the referees for the helpful suggestions.

References

- [1] C. Baikoussis and Th. Koufogiorgos, *On the inner curvature of the second fundamental form of helicoidal surfaces*, Arch. Math. **68** (1997), 169-176.
- [2] D. E. Blair and Th. Koufogiorgos, *Ruled surfaces with vanishing second Gaussian curvature*, Monatsh. Math. **113** (1992), 177-181.
- [3] F. Dillen and W. Kühnel, *Ruled Weingarten surfaces in Minkowski 3-space*, Manuscripta Math. **98** (1999), 307-320.
- [4] F. Dillen and W. Sodsiri, *Ruled surfaces of Weingarten type in Minkowski 3-space*, J. Geom. **83** (2005), 10-21.
- [5] F. Dillen and W. Sodsiri, *Ruled surfaces of Weingarten type in Minkowski 3-space II*, J. Geom. **84** (2005), 37-44.
- [6] Y. H. Kim and D. W. Yoon, *Classification of ruled surfaces in Minkowski 3-spaces*, J. Geom. Phys. **49** (2004), 89-100.

- [7] Y. H. Kim and D. W. Yoon, *On non-developable ruled surfaces in Lorentz-Minkowski 3-spaces*, Taiwanese J. Math. **11** (2007), 197-214.
- [8] N. G. Kim and D. W. Yoon, *Mean curvature of non-degenerate second fundamental form of ruled surfaces*, Honam Math. J. **28** (2006), 549-558.
- [9] Th. Koufogiorgos and T. Hasanis, *A characteristic property of the sphere*, Proc. Amer. Math. Soc. **67** (1977), 303-305.
- [10] D. Koutroufiotis, *Two characteristic properties of the sphere*, Proc. Amer. Math. Soc. **44** (1974), 176-178.
- [11] W. Kühnel, *Ruled W-surfaces*, Arch. Math. **62** (1994), 475-480.
- [12] R. López, *Special Weingarten surfaces foliated by circles*, to appear in Monatsh. Math.
- [13] G. Stamou, *Regelflächen vom Weingarten-type*, Colloq. Math. **79** (1999), 77-84.
- [14] D. W. Yoon, *Some properties of the helicoid as ruled surfaces*, JP Jour. Geom. Topology **2** (2002), 141-147.
- [15] D. W. Yoon, *On non-developable ruled surfaces in Euclidean 3-spaces*, Indian J. pure appl. Math. **38** (2007), 281-289.

*

Seonggwang High School
Daegu 702-817, Republic of Korea
E-mail: blackr1@naver.com

**

Department of Mathematics Education and RINS
Gyeongsang National University
Jinju 660-701, Republic of Korea
E-mail: dwyoon@gnu.ac.kr