

Role of Autophagy in the Control of Cell Death and Inflammation

Myung-Shik Lee*

Department of Medicine, Samsung Medical Center, Sungkyunkwan University, School of Medicine, Seoul 135-710, Korea

There is mounting evidence that autophagy is involved in diverse physiological and pathological processes that have immense relevance in human development, diseases and aging. Immunity and inflammation are not exceptions. Here, the role of autophagy in the control of immune processes particularly that related to cell death and inflammation is discussed.

[Immune Network 2009;9(1):8-11]

INTRODUCTION

Macroautophagy (here referred to as autophagy) is a process of sequestration and removal of damaged organelles/proteins for recycling of their constituents and supply of nutrients. In addition, autophagy provides signals for the removal of apoptotic cells and genomic stability (1). Hence, autophagy can be generally considered as a protective measure for cells against various types of injuries or from continuous cellular wear-and-tear. The physiological role of autophagy in the mammalian system was demonstrated in several *in vivo* animal models. Disruption of autophagic processes leads to failure of cavitation during embryogenesis (2) or accumulation of abnormal mitochondria in adult tissues (3). In addition to the physiological roles, dysregulated autophagy has been suggested to play pathogenic roles in a variety of disease processes including neurodegeneration (4,5) and cardiomyopathy (6) particularly when cellular stress is increased, which is likely to be due to accumulation of damaged molecules and organelles (6). Other studies revealed important roles for autophagy in tumor suppression (1) and ER stress responses (7).

AUTOPHAGY AND CELL DEATH

As discussed above, autophagy plays a protective role in diverse types of cellular stress. Paradoxically, autophagy can also lead to a form of non-apoptotic cell death known as "type 2 programmed cell death" (8-10) particularly when autophagy is excessive (11). Thus, the dual nature of autophagy can either promote cell death or protect cells from diverse types of injuries depending on the cellular and environmental context. In mice models that have a deficiency of autophagy specifically in neurons, increased apoptosis of neurons were observed (4,5). We and others have shown in mice with a pancreatic β -cell-specific deletion of an essential autophagy gene (*Atg7*) that the pancreatic β -cell mass was decreased due to increased pancreatic β -cell death and decreased β -cell proliferation leading to a reduction of insulin production and hyperglycemia (12,13). Large intracellular aggregates containing ubiquitin and p62 were observed in the pancreatic islets of β -cell-specific *Atg7*-null mice (13,14). However, inflammation was not observed but in the case of hepatocyte-specific *Atg7*-null mice it was observed (data not shown). Increased pancreatic β -cell death in β -cell-specific *Atg7*-null mice is consistent with previous data that suggests autophagy plays a protective role against cell injury and autophagy-deficient cells would be susceptible to diverse types of cell death. Several potential mechanisms of increased susceptibility to cell death were proposed such as a shortage of bioenergetic sources like NADPH₂ and ATP. The shortage of these sources may trigger apoptosis presumably through a direct effect on the mitochondria or de-inhibit the activation of caspases or permeabilization of the mitochondrial outer mem-

Received on January 9, 2009. Revised on January 16, 2009. Accepted on January 20, 2009.

*Corresponding Author. Tel: 82-2-3410-3436; Fax: 82-2-3410-6491; E-mail: mslee0923@skku.edu

Keywords: autophagy, inflammation, immunity, apoptosis, necrosis

brane (15) or through activation of p27 by metabolic stress and energy deficiency that may also lead to cell cycle arrest and decreased proliferation (16). Modulation of mitochondrial mass or cytochrome *c* content by autophagy may be related to the changes in the susceptibility to apoptotic inducers (17). However, detailed mechanism of increased pancreatic β -cell death in β -cell-specific *Atg7*-null mice is unknown. Furthermore, it was suggested that the real process inhibited by autophagy is necrosis rather than apoptosis which may be of relevance to the control of cell death-associated inflammation and tumorigenesis (18).

AUTOPHAGY, MACROPHAGE AND INFLAMMATION

Autophagy may also play a crucial role in the regulation of inflammation and immunity. For instance, autophagy has been reported to be critically involved in the eradication of microbes such as *Mycobacterium tuberculosis* (19), antigen presentation by professional antigen-presenting cells (APC) (20) and tolerance induction in T cells (21). Autophagy may also regulate inflammation or immunity by modulating survival or death of lymphocytes (22) and APC such as macrophages. We have reported the role of ROS and STAT1 in the necrotic death of macrophage by LPS in the presence of zVAD, a pan-caspase inhibitor (23). Then, a role for autophagy in this type of macrophage death was reported. Xu et al. showed the death of macrophages by LPS in combination with zVAD had microscopic features of autophagy such as punctuate LC3 pattern and formation of autophagosomes surrounded by double membranes (24). Furthermore, Yu et al. reported that autophagy selectively degraded catalase leading to the accumulation of ROS and non-apoptotic death of macrophages (25). Contrary to those reports, a protective role of autophagy in macrophage death by zVAD was also reported (26). If zVAD is able to inhibit cathepsin and lysosomal function as reported, then the apparent increased conversion of LC3-I to LC3-II and the accumulation of autophagosomes may be due to the blockade of autophagy at the lysosomal step rather than to an increased autophagic activity. Such a theory may provide a clue to the long-sought question regarding the additional target molecules of zVAD in addition to caspases. Thus, it is still unclear whether autophagy protects macrophages from death or induces macrophage death in the presence of zVAD.

Autophagy may also modulate functions of macrophages

in addition to their survival or death. LC3 recruitment was observed when macrophages ingested latex beads associated with TLR ligands, but not without TLR ligands (27). However, classical autophagosomes with double membranes were not observed in that case. Furthermore, microbes engulfed by autophagy-deficient macrophages showed a marked survival advantage suggesting that TLR signaling may usurp the autophagic machinery to deliver phagocytic material to the lysosomal pathway (27). TLR agonists themselves have been reported to activate the autophagic pathway (28); however, a recent paper reported contradictory results (29).

AUTOPHAGY AND INFLAMMATORY/IMMUNE DISEASES

Besides *in vitro* or animal models, direct involvement of the autophagy-inflammation relationship in human diseases was recently reported. Crohn's disease is characterized by chronic mucosal or transmural inflammation in the small intestines particularly in the distal ileum. Several theories have been proposed related to the pathogenesis of Crohn's disease such as mutations of NOD₂, an important intracellular innate immune receptor (30). Also several animal models are available for the study of Crohn's disease. However, the detailed molecular pathogenic mechanism is still far from clear. Large-scale genetic analysis revealed more than 30 loci showing linkage to the disease development process (31). Among them, *Atg16L1* locus is of particular interest because *Atg16L1* is a gene homolog to yeast *Atg16*, an essential gene for autophagy in yeast. *Atg16L1* is an integral component of the large molecular weight complex comprising *Atg12-Atg5* conjugate and has been suggested to play an important role in the conjugation of LC3 (*Atg8*) to phosphatidylethanolamine. Thus, *Atg16L1* has been regarded as a key gene that may provide a clue to the relationship between autophagy and inflammation in human diseases. Indeed, a recent paper showed a role of *Atg16L1* in the control of inflammatory cytokines from macrophages (29). Release of cytokines such as IL-1 β and IL-18 after LPS or infection with non-invasive bacteria such as *E. coli* was significantly increased in *Atg16L1*-null macrophages. Intriguingly, IL-1 β release after infection with invasive bacteria such as *Salmonella typhimurium* was not increased in *Atg16L1*-null macrophages. Increased IL-1 β or IL-18 secretion from *Atg16L1*-null macrophages appears to be due to increased ROS production inducing activation of caspase-1. However, the molecular details

leading to the increase ROS production in autophagy-deficient macrophages are still unknown. Mice lacking Atg16L1 in hematopoietic cells did not show spontaneous colitis; however, they showed much severer colitis after dextran sulphate sodium (DSS) challenge. Another recent paper showed an abnormality in Paneth cells that are specialized cells in the intestinal epithelium secreting antimicrobial peptides or other intestinal peptides (32) in Atg16L1-deficient mice (33). Furthermore, production of adiponectin and leptin that modify the intestinal environment was also altered in autophagy-deficient Paneth cells.

CONCLUSION

Autophagy appears to be critically involved in inflammation and immunity. So far, several pioneering studies have been done. Some of the results are not consistent with each other, which are more than likely due to an incomplete knowledge or misconception regarding the process of autophagy. For example, it is becoming clear that autophagy level and autophagic activity are not always the same, and the blockade of the autophagic process may be confused with increased autophagy due to the possible accumulation of autophagosomes. Thus, more thorough evaluation will be required for proper interpretation of future experimental findings and also for that of previously reported results. More investigations are being done and will be necessary to reveal the relationship between autophagy and the diverse types of inflammatory and (auto) immune disorders which will lead to the elucidation of hitherto unknown and unsuspected aspects of both adaptive and innate immunity.

ACKNOWLEDGEMENTS

This work was supported by a Nano/Bio Science Program Grant (2004-00716) and by the 21C Frontier Functional Proteomics Project of the Korean Ministry of Science & Technology (FPR05C1-160). Lee M-S is the recipient of The BioScience Grant (2008-057-6001) from the Korea Science & Engineering Foundation.

CONFLICTS OF INTEREST

Disclosures: The authors have no financial conflict of interest.

REFERENCES

1. Levine B, Kroemer G: Autophagy in the pathogenesis of disease. *Cell* 132;27-42, 2008
2. Qu X, Zou Z, Sun Q, Luby-Phelps K, Cheng P, Hogan RN, Gilpin C, Levine B: Autophagy gene-dependent clearance of apoptotic cells during embryonic development. *Cell* 128;931-946, 2007
3. Komatsu M, Waguri S, Ueno T, Iwata J, Murata S, Tanida I, Ezaki J, Mizushima N, Ohsumi Y, Uchiyama Y, Kominami E, Tanaka K, Chiba T: Impairment of starvation-induced and constitutive autophagy in Atg7-deficient mice. *J Cell Biol* 169;425-434, 2005
4. Hara T, Nakamura K, Matsui M, Yamamoto A, Nakahara Y, Suzuki-Migishima R, Yokoyama M, Mishima K, Saito I, Okano H, Mizushima N: Suppression of basal autophagy in neural cells causes neurodegenerative disease in mice. *Nature* 441;885-889, 2006
5. Komatsu M, Waguri S, Chiba T, Murata S, Iwata J, Tanida I, Ueno T, Koike M, Uchiyama Y, Kominami E, Tanaka K: Loss of autophagy in the central nervous system causes neurodegeneration in mice. *Nature* 441;880-884, 2006
6. Nakai A, Yamaguchi O, Takeda T, Higuchi Y, Hikoso S, Taniike M, Omiya S, Mizote I, Matsumura Y, Asahi M, Nishida K, Hori M, Mizushima N, Otsu K: The role of autophagy in cardiomyocytes in the basal state and in response to hemodynamic stress. *Nat Med* 13;619-624, 2007
7. Ogata M, Hino SI, Saito A, Morikawa K, Kondo S, Kanemoto S, Murakami T, Taniguchi M, Tani I, Yoshinaga K, Shiosaka S, Hammarback JA, Urano F, Imaizumi K: Autophagy is activated for cell survival after endoplasmic reticulum stress. *Mol Cell Biol* 26;9220-9231, 2006
8. Kim EH, Sohn S, Kwon HJ, Kim SU, Kim MJ, Lee SJ, Choi KS: Sodium selenite induces superoxide-mediated mitochondrial damage and subsequent autophagic cell death in malignant glioma cells. *Cancer Res* 67;6314-6324, 2007
9. Park KJ, Lee SH, Kim TI, Lee HW, Lee CH, Kim EH, Jang JY, Choi KS, Kwon MH, Kim YS: A human scFv antibody against TRAIL receptor 2 induces autophagic cell death in both TRAIL-sensitive and TRAIL-resistant cancer cells. *Cancer Res* 67;7327-7334, 2008
10. Pyo JO, Jang MH, Kwon YK, Lee HJ, Jun JI, Woo HN, Cho DH, Choi B, Lee H, Kim JH, Mizushima N, Ohsumi Y, Jung YK: Essential roles of Atg5 and FADD in autophagic cell death: dissection of autophagic cell death into vacuole formation and cell death. *J Biol Chem* 280;20722-20729, 2005
11. Boya P, González-Polo RA, Casares N, Perfettini JL, Dessen P, Larochette N, Métivier D, Meley D, Souquere S, Yoshimori T, Piéron G, Codogno P, Kroemer G: Inhibition of macroautophagy triggers apoptosis. *Mol Cell Biol* 25;1025-1040, 2005
12. Ebato C, Uchida T, Arakawa M, Komatsu M, Ueno T, Korniya K, Azuma K, Hirose T, Tanaka K, Kominami E, Kawamori R, Fujitani Y, Watada H: Autophagy is important in islet homeostasis and compensatory increase of beta cell mass in response to high-fat diet. *Cell Metab* 8;325-332, 2008
13. Jung HS, Chung KW, Won KJ, Kim J, Komatsu M, Tanaka

- K, Nguyen YH, Kang TM, Yoon KH, Kim JW, Jeong YT, Han MS, Lee MK, Kim KW, Shin J, Lee MS: Loss of autophagy diminishes pancreatic beta cell mass and function with resultant hyperglycemia. *Cell Metab* 8:318-324, 2008
14. Jung HS, Lee MS: Macroautophagy in homeostasis of pancreatic b-cells. *Autophagy* 5:1-3, 2009
 15. Maiuri MC, Zalckvar E, Kimchi A, Kroemer G: Self-eating and self-killing: crosstalk between autophagy and apoptosis. *Nat Rev Mol Cell Biol* 8:741-752, 2007
 16. Liang J, Shao SH, Xu ZX, Hennessy B, Ding Z, Larrea M, Kondo S, Dumont DJ, Guterman JU, Walker CL, Slingerland JM, Mills GB: The energy sensing LKB1-AMPK pathway regulates p27(kip1) phosphorylation mediating the decision to enter autophagy or apoptosis. *Nat Cell Biol* 9:218-224, 2007
 17. Ravikumar B, Berger Z, Vacher C, O'Kane CJ, Rubinsztein DC: Rapamycin pre-treatment protects against apoptosis. *Hum Mol Genet* 15:1209-1216, 2006
 18. Degenhardt K, Mathew R, Beaudoin B, Bray K, Anderson D, Chen G, Mukherjee C, Shi Y, Gélinas C, Fan Y, Nelson DA, Jin S, White E: Autophagy promotes tumor cell survival and restricts necrosis, inflammation, and tumorigenesis. *Cancer Cell* 10:51-64, 2006
 19. Gutierrez MG, Master SS, Singh SB, Taylor GA, Colombo MI, Deretic V: Autophagy is a defense mechanism inhibiting BCG and Mycobacterium tuberculosis survival in infected macrophages. *Cell* 119:753-766, 2004
 20. Schmid D, Pypaert M, Münz C: Antigen-loading compartments for major histocompatibility complex class II molecules continuously receive input from autophagosomes. *Immunity* 26:79-92, 2007
 21. Nedjic J, Aichinger M, Emmerich J, Mizushima N, Klein L: Autophagy in thymic epithelium shapes the T-cell repertoire and is essential for tolerance. *Nature* 455:396-400, 2008
 22. Feng CG, Zheng L, Jankovic D, Báfica A, Cannons JL, Watford WT, Chaussabel D, Hieny S, Caspar P, Schwartzberg PL, Lenardo MJ, Sher A: The immunity-related GTPase Irgm1 promotes the expansion of activated CD4+ T cell populations by preventing interferon-gamma-induced cell death. *Nat Immunol* 9:1279-1287, 2008
 23. Kim HS, Lee MS: Essential role of STAT1 in caspase-independent cell death of activated macrophages through the p38 mitogen activated protein kinase/STAT1/reactive oxygen species pathway. *Mol Cell Biol* 25:6821-6833, 2005
 24. Xu Y, Kim SO, Li Y, Han J: Autophagy contributes to caspase-independent macrophage cell death. *J Biol Chem* 281:19179-19187, 2006
 25. Yu L, Wan F, Dutta S, Welsh S, Liu ZH, Freundt E, Baehrecke EH, Lenardo MJ: Autophagic programmed cell death by selective catalase degradation. *Proc Natl Acad Sci U S A* 103:4952-4957, 2006
 26. Wu YT, Tan HL, Huang Q, Kim YS, Pan N, Ong WY, Liu ZG, Ong CN, Shen HM: Autophagy plays a protective role during zVAD-induced necrotic cell death. *Autophagy* 4:457-466, 2008
 27. Sanjuan MA, Dillon CP, Tait SWG, Moshiah S, Dorsey F, Connell S, Komatsu M, Tanaka K, Cleveland JC, Withoff S, Green DR: Toll-like receptor signalling in macrophages links the autophagy pathway to phagocytosis. *Nature* 450:1253-1257, 2007
 28. Xu Y, Jagannath C, Liu XD, Sharafkhaneh A, Kolodziejka KE, Eissa NT: Toll-like receptor 4 is a sensor for autophagy associated with innate immunity. *Immunity* 27:135-144, 2007
 29. Saitoh T, Fujita N, Jang MH, Uematsu S, Yang BG, Satoh T, Omori H, Noda T, Yamamoto N, Komatsu M, Tanaka K, Kawai T, Tsujimura T, Takeuchi O, Yoshimori T, Akira S: Loss of the autophagy protein Atg16L1 enhances endotoxin-induced IL-1beta production. *Nature* 456:264-269, 2008
 30. Maeda S, Hsu LC, Liu H, Bankston LA, Iimura M, Kagnoff MF, Eckmann L, Karin M: Nod2 mutation in Crohn's disease potentiates NF-kappaB activity and IL-1beta processing. *Science* 307:734-738, 2005
 31. Massey DC, Parkes M: Genome-wide association scanning highlights two autophagy genes, ATG16L1 and IRGM, as being significantly associated with Crohn's disease. *Autophagy* 3:649-651, 2007
 32. Vaishnava S, Behrandt CL, Ismail AS, Eckmann L, Hooper LV: Paneth cells directly sense gut commensals and maintain homeostasis at the intestinal host-microbial interface. *Proc Natl Acad Sci U S A* 105:20858-20863, 2008
 33. Cadwell K, Liu JY, Brown SL, Miyoshi H, Loh J, Lennerz JK, Kishi C, Kc W, Carrero JA, Hunt S, Stone CD, Brunt EM, Xavier RJ, Sleckman BP, Li E, Mizushima N, Stappenbeck TS, Virgin HW 4th: A key role for autophagy and the autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature* 456:259-264, 2008