

# Involvement of Hepatic Innate Immunity in Alcoholic Liver Disease

Jin-Seok Byun and Won-Il Jeong\*

Graduate School of Medical Science and Engineering, KAIST Institute for the BioCentury, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Korea

Excessive alcohol consumption is one of the critical causative factors leading to alcoholic liver disease (ALD). ALD is characterized by a wide spectrum of liver damage, ranging from simple uncomplicated liver steatosis (fatty liver) to steatohepatitis and liver fibrosis/cirrhosis. It has been believed that the obvious underlying cause for ALD is due to hepatocyte death induced by alcohol itself. However, recent sparkling studies have shown that diverse immune responses contribute to ALD because liver is enriched with numerous immune cells. Especially, a line of evidence has suggested that innate immune cells such as Kupffer cells and natural killer (NK)/NKT cells are significantly involved in the pathogenesis of ALD via production of pro-inflammatory cytokines and other mediators. Indeed, more interestingly, hepatic stellate cells (HSCs), known as a major cell inducing liver steatosis and fibrosis, can be killed by liver NK cells, which could be suppressed by chronic alcohol consumption. In this review, with the view of liver as predominant innate immune organ, we describe the pathogenesis of ALD in which what roles of innate immune cells are and how they are interacting with HSCs.

[Immune Network 2010;10(6):181-187]

## INTRODUCTION

Liver disease is one of major causes of morbidity and mortality worldwide and alcohol is one of the most prominent factors in liver disease (1,2). Alcoholic liver disease (ALD) encompasses a broad spectrum of diseases ranging from steatosis (fatty liver), steatohepatitis, fibrosis, cirrhosis to hepatocellular carcinoma (3,4). Of the ALD, steatosis is consid-

ered as mild condition, whereas steatohepatitis is pathogenic condition, which has a potential of progression to liver fibrosis, cirrhosis or hepatocellular carcinoma (5). Generally, the liver is composed of parenchymal cells (e.g. hepatocyte) and non-parenchymal cells such as sinusoidal endothelial cells, Kupffer cells, hepatic stellate cells (HSCs), dendritic cells and other lymphocytes. Interestingly, liver comprises enrichment of innate immune cells such as resident macrophages (Kupffer cells), natural killer (NK), NKT and  $\gamma \delta$  T cells. (6-8). For example, mouse liver lymphocytes contain 10% NK cells, whereas rat and human liver lymphocytes contain about 30% to 50% NK cells, NKT cells constitute up to 30% and 10% of the intrahepatic lymphocyte population in mice and human respectively (6,7,9). Interestingly, ALD has been considered as complex consequences of interaction among these cells, especially between hepatocytes and hepatic stellate cells or innate immune cells (3,4). Previous our studies demonstrated that alcoholic liver steatosis was induced by HSC-mediated endocannabinoid and its hepatic CB1 receptor (10) and alcoholic liver fibrosis was accelerated due to abrogation of antifibrotic effects of NK cells/interferon- $\gamma$  (IFN- $\gamma$ ) against HSCs (11). Other studies suggest that Kupffer cells are mainly involved in alcohol-mediated inflammation via lipopolysaccharide (LPS)/toll like receptor 4 (TLR4) signaling-dependant mechanisms (4,12). Based on these evidences, researchers recognized the importance of innate immune response in ALD. Therefore, in this review, we will focus and discuss about diverse involvements of HSCs and innate immunity (Kupffer cells/macrophages and NK cells) in ALD

Received on December 13, 2010. Accepted on December 13, 2010.

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

\*Corresponding Author. Tel: 82-42-350-4239; Fax: 82-42-350-4241; E-mail: [wijeong@kaist.ac.kr](mailto:wijeong@kaist.ac.kr)

Keywords: Natural killer cell, Interferon-gamma, Hepatic stellate cell, Steatosis, Steatohepatitis, Fibrosis, Alcohol

such as alcoholic steatosis, steatohepatitis and liver fibrosis.

## ALCOHOLIC LIVER STEATOSIS (FATTY LIVER)

Alcoholic liver steatosis has been considered as a weak condition for a long time. However, increasing evidence suggests that it seems to be a potentially pathologic state and it will progress more severe state in the presence of other co-factors such as sustained consumption of alcohol, hepatic virus infection, diabetes or drugs (13,14). Fat accumulation in hepatocytes is the result of imbalanced fat metabolism such as decrease mitochondrial lipid oxidation but enhanced synthesis of triglycerides. Several underlying mechanism of this have been suggested that it might be related with increased NADH/NAD<sup>+</sup> ratio (15,16), increased sterol regulatory element-binding protein-1 (SREBP-1) activity (17,18), decreased peroxisome proliferator-activated receptor- $\alpha$  (PPAR- $\alpha$ ) activity (19,20) and decreased AMP-activated protein kinase (AMPK) activity (10,18).

In addition to pathological mechanisms for this, there are several interesting evidences that innate immune cells of liver are related in alcoholic liver steatosis (3,4,12,21,22). Kupffer cells are one of the main innate immune cells involved in steatotic liver. First, alcohol increased gut permeabilization, which allows more uptake of endotoxin/LPS in portal circulation (21) and then Kupffer cells are activated in response to LPS via TLR4 signaling cascade, leading to produce several kinds of pro-inflammatory cytokine such as tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin (IL)-1, IL-6 and reactive oxygen species (ROS) (3,4,22). Of these cytokines, contribution of TNF- $\alpha$  in developing alcoholic liver steatosis has been less characterized compared with TNF- $\alpha$ -mediated hepatic inflammation. However, increased expression of TNF- $\alpha$  has been observed in alcoholic liver steatosis of mouse (23,24) and absence of its receptor (TNF- $\alpha$  R1) activity inhibits the development of alcoholic liver steatosis (25,26). In addition, it has been reported that TNF- $\alpha$  has a potential to increase mRNA expression of SREBP-1c, a potent transcription factor of fat synthesis, in the liver of mice and to stimulate the maturation of SREBP-1 in human hepatocytes, respectively (27,28). Furthermore, recent report showed that alcohol-mediated infiltration of macrophages decreased amounts of adiponectin (known as anti-steatosis peptide hormone) production of adipocytes, leading to alcoholic liver steatosis (29). Therefore, Kupffer cells/macrophages may contribute to the development of alcoholic liver steatosis by upregulating the SREBP1

activity in hepatocytes and downregulating the production of adiponectin in adipocytes. In contrast, IL-6 produced by Kupffer cells may play a crucial role in protecting against alcoholic liver steatosis via activation of signal transducer and activator of transcription 3 (STAT3), consequently inhibiting of SREBP1 gene expression in hepatocytes (30-32).

More interestingly, a recent our study reported that alcoholic liver steatosis was mediated mainly through activated HSCs-derived endocannabinoid and its receptor (CB1R) of hepatocytes (10). This study suggested that chronic alcohol drinking activates HSCs to produce 2-arachidonoylglycerol (2-AG), one of endocannabinoid, which then increased the expression of SREBP1c and fatty acid synthase (FAS) but decreased AMPK activation, consequently leading to accumulation of fat in hepatocyte. Another studies reported that activated HSC can be killed by liver NK cells through IFN- $\gamma$  and tumor necrosis factor-related apoptosis inducing ligand (TRAIL)-dependent manners (11,33-35). However, chronic alcohol consumption inhibited NK cytotoxicity against activated HSC, leading to prolonged survival of activated HSC (11).

## ALCOHOLIC STEATOHEPATITIS (ASH)

Alcoholic steatohepatitis means united condition with fat accumulation and inflammation in the liver and more progressed pathologic state compared with alcoholic liver steatosis. In response to alcohol intake, innate immune cells initiate and maintain hepatic inflammation via pattern recognition receptors, especially TLRs (36-38).

Among activation of innate immune cells in liver, Kupffer cells have been identified as one of the significant elements in the pathogenesis of alcoholic steatohepatitis (22). Considering their specific location at the interface between the portal and the systemic circulation, Kupffer cells play a key role in orchestrating the immune response against endotoxin/LPS. LPS, a component of gram-negative bacteria wall, has been considered to be a key molecule of activation in Kupffer cells, in which signals are transmitted through TLR4 (3,38-40). TLR4 is a major component of the LPS recognition receptor complex, which also involves the co-receptors CD14 and MD-2, and LPS binding protein (LBP) (41,42). LBP is a soluble shuttle protein that directly binds LPS and facilitates the association between LPS and CD14 (43). Studies in knockout mouse models have shown that chronic alcohol feeding in mice deficient of CD14, TLR4 and LPS-binding protein (LBP) results in alleviation of alcohol-induced liver injury indicating an im-

portant role for the TLR4 pathway (40,44,45). In addition, the importance of gut-derived endotoxin/LPS in ALD was suggested by experiments where treating the animals either with antibiotics or with lactobacilli to remove or reduce the gut microflora provided protection from the features of ALD (46). TLR4 initiates two major pathways. After binding LPS with TLR4, recruited TIR domain-containing adaptor protein (TIRAP) and myeloid differentiation factor 88 (MyD88) lead to early-phase activation of nuclear factor- $\kappa$  B (NF- $\kappa$  B), producing pro-inflammatory cytokines including TNF- $\alpha$ , IL-6 and monocyte chemoattractant protein-1 (MCP-1). Meanwhile TIR-domain containing adaptor inducing IFN- $\beta$  (TRIF) and TRIF-related adaptor molecule (TRAM) activate interferon regulatory factor 3 (IRF3) leading to the production of type I IFN and late activation of NF- $\kappa$  B (38,47). However, there is a recent report that alcohol-mediated liver injury and inflammation are mainly induced by TLR4-dependent but MyD88-independent manners (48). Among pro-inflammatory cytokines, particularly TNF- $\alpha$  mainly contributes to the development of ALD and its level is increased in patients with ASH (22) and in the liver of alcohol-fed animals (23,24). In addition, Kupffer cells secrete other important cytokines, including IL-8, IL-12, and interferons, that contribute to the intrahepatic recruitment and activation of granulocytes that are characteristically found in severe ALD, influence immune system polarization (49). Interestingly, TLR4 is expressed not only on innate immune cells such as Kupffer cells and recruited macrophages, but also on hepatocytes, sinusoidal endothelial cells and stellate cells in the liver (39). The role of TLR4 in these cells has to be investigated.


In addition to LPS, oxidative stress-mediated cellular responses also play an important role in innate immune cell activation. Kupffer cells are also a major source of ROS in response to chronic alcohol exposure (50,51). One important ROS is the superoxide ion, which in activated phagocytes is mainly generated by the enzyme complex NADPH oxidase. Underscoring the important role of ROS in mediating ethanol damage, treatment with antioxidants and deletion of the p47phox subunit of NADPH oxidase in ethanol-fed animals could reduce oxidative stress, activation of NF- $\kappa$  B, and TNF- $\alpha$  release in Kupffer cells, thus, preventing liver injury (50,52). In addition, NADPH oxidase induces TLR2 and TLR4 expression in human monocytic cells (53). Furthermore, evidence shows that direct interaction of NADPH oxidase isozyme 4 with TLR4 is involved in LPS-mediated ROS generation and NF- $\kappa$  B activation in neutrophils (54).

## ALCOHOLIC LIVER FIBROSIS

HSCs are known as a major cell type inducing liver fibrosis (55). Generally, HSCs are located in the Disse's space as a quiescent status in normal healthy liver, but by liver injuries they become activated and differentiated into myofibroblastic cells that are characterized by a loss of vitamin A (retinol) and enhanced extracellular matrix, especially collagen fibers, leading to liver fibrosis.

Interestingly, recent reports suggest that innate immune cells (Kupffer cells and NK/NKT cells) are closely involved in the liver fibrogenesis. First, Kupffer cells are generally considered as a positive regulator in liver fibrosis. Actually, increasing evidences suggest that chronic alcohol consumption leads to liver fibrosis via the activation of the profibrogenic effects of Kupffer cells due to enhanced transport of endotoxin/LPS in the gut, inducing production of pro-inflammatory cytokines such as TNF- $\alpha$ , IL-6 and transforming growth factor- $\beta$  (TGF- $\beta$ ) (3,14,23,24). Among these cytokines, TGF- $\beta$  is known as a key regulator of HSC activation and differentiation. Thus, it is believed that Kupffer cells positively regulate alcoholic liver fibrosis via TGF- $\beta$ -mediated HSC activation and subsequently promote collagen synthesis of HSCs. In contrast, recent findings interestingly suggest the novel role of NK cells having anti-fibrotic effects via multiple mechanisms (33-35,56,57). First, NK cells directly kill early activated HSCs in NKG2D- and TRAIL-dependent manners but not quiescent HSCs (35,56). This is because early activated HSCs express NK cell-activating ligand retinoic acid early inducible gene 1 (RAE-1) via retinol metabolism and TRAIL receptors but express decreased MHC-I, NK cell-inhibitory ligand (33,57). Second, NK cells can suppress liver fibrosis via production of IFN- $\gamma$ , which induces HSC cell cycle arrest and apoptosis in STAT1-dependant manner (34,58). Analogous to NK cells, NKT cells (invariant NKT cells) can also suppress activation of HSCs via direct killing and IFN- $\gamma$  production (59). However, anti-fibrotic effects of NKT cells are beneficial only at early stage of liver fibrosis because of their quick depletion.

Although the functions of hepatic NK/NKT cells are known a little in ALD, it is generally accepted now that chronic alcohol consumption accelerates liver fibrosis due to the suppressed activity of NK cell in patients and mice and enhanced alcoholic liver injury as well (11,60-62). The activation status of NK cells has been reported to be decreased by alcohol, possibly because of increased IL-10 and TGF- $\beta$  release by



**Figure 1.** Involvement of innate immune cells and HSCs in alcoholic liver disease (ALD).

monocytes and activated HSCs (11,63) In case of NKT cells, they seem to contribute to alcoholic liver injury since activation of NKT cell accelerates whereas NKT deficiency delays alcoholic liver injury (64,65). Nevertheless reports of alcohol effects on NK/NKT cell functions are still controversial. Therefore, further investigation is needed in animal and human.

## CONCLUSIONS

The innate immune cells of liver play important roles in dangerous stress such as alcoholic liver injuries and infection of microorganism. Recently a line of evidences has demonstrated that innate immune responses are closely connected with the development of ALD and we integrate these findings in Fig. 1. Especially, activation of Kupffer cells by alcohol appears to be required for the development of alcoholic steatohepatitis via LPS-TLR4 signaling pathways. However, endocannabinoid production (2-AG) of activated HSCs might be another major factor for the induction of alcoholic steatosis. Therefore, we should simultaneously consider both types of

cells when developing therapeutics for alcoholic steatohepatitis. For example, although we successfully inhibit activation of Kupffer cells by drugs activated HSCs still induce accumulation of fat in the liver, leading to lipotoxicity and in turn to generative of oxidative stress and inflammation, subsequently restoring steatohepatitis. More interestingly, although the role of NKT in ALD is obscure, NK cells are responsible for anti-steatotic and anti-fibrotic functions via killing activated HSCs that may reduce production of 2-AG and collagen. However, functions of NK cells are abrogated or suppressed by chronic alcohol consumption and high level of TGF- $\beta$  in the liver. Therefore, further understanding of the roles of innate immunity and HSCs help us to develop novel therapeutic targets to treat ALD.

## ACKNOWLEDGEMENTS

This work was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2009-0072049).

## CONFLICTS OF INTEREST

The authors have no financial conflict of interest.

## REFERENCES

1. Williams R: Global challenges in liver disease. *Hepatology* 44:521-526, 2006
2. O'Shea RS, Dasarthy S, McCullough AJ: Practice Guideline Committee of the American Association for the Study of Liver Diseases; Practice Parameters Committee of the American College of Gastroenterology: Alcoholic liver disease. *Hepatology* 51:307-328, 2010
3. Jeong WI, Gao B: Innate immunity and alcoholic liver fibrosis. *J Gastroenterol Hepatol* 23(Suppl 1):S112-S118, 2008
4. Purohit V, Gao B, Song BJ: Molecular mechanisms of alcoholic fatty liver. *Alcohol Clin Exp Res* 33:191-205, 2009
5. Purohit V, Brenner DA: Mechanisms of alcohol-induced hepatic fibrosis: a summary of the Ron Thurman Symposium. *Hepatology* 43:872-878, 2006
6. Gao B, Jeong WI, Tian Z: Liver: An organ with predominant innate immunity. *Hepatology* 47:729-736, 2008
7. Racanelli V, Rehermann B: The liver as an immunological organ. *Hepatology* 43(2 Suppl 1):S54-S62, 2006
8. Crispe IN: The liver as a lymphoid organ. *Annu Rev Immunol* 27:147-163, 2009
9. Doherty DG, O'Farrelly C: Innate and adaptive lymphoid cells in the human liver. *Immunol Rev* 174:5-20, 2000
10. Jeong WI, Osei-Hyiaman D, Park O, Liu J, Bátkai S, Mukhopadhyay P, Horiguchi N, Harvey-White J, Marsicano G, Lutz B, Gao B, Kunos G: Paracrine activation of hepatic CB1 receptors by stellate cell-derived endocannabinoids mediates alcoholic fatty liver. *Cell Metab* 7:227-235, 2008
11. Jeong WI, Park O, Gao B: Abrogation of the antifibrotic effects of natural killer cells/interferon-gamma contributes to alcohol acceleration of liver fibrosis. *Gastroenterology* 134:248-258, 2008
12. Mandrekar P, Szabo G: Signalling pathways in alcohol-induced liver inflammation. *J Hepatol* 50:1258-1266, 2009
13. Otani K, Korenaga M, Beard MR, Li K, Qian T, Showalter LA, Singh AK, Wang T, Weinman SA: Hepatitis C virus core protein, cytochrome P450 2E1, and alcohol produce combined mitochondrial injury and cytotoxicity in hepatoma cells. *Gastroenterology* 128:96-107, 2005
14. Lieber CS: Alcoholic fatty liver: its pathogenesis and mechanism of progression to inflammation and fibrosis. *Alcohol* 34:9-19, 2004
15. Crabb DW: Recent developments in alcoholism: the liver. *Recent Dev Alcohol* 11:207-230, 1993
16. Fromenty B, Berson A, Pessayre D: Microvesicular steatosis and steatohepatitis: role of mitochondrial dysfunction and lipid peroxidation. *J Hepatol* 26(Suppl 1):13-22, 1997
17. Yahagi N, Shimano H, Hasty AH, Matsuzaka T, Ide T, Yoshikawa T, Amemiya-Kudo M, Tomita S, Okazaki H, Tamura Y, Iizuka Y, Ohashi K, Osuga J, Harada K, Gotoda T, Nagai R, Ishibashi S, Yamada N: Absence of sterol regulatory element-binding protein-1 (SREBP-1) ameliorates fatty livers but not obesity or insulin resistance in *Lep(ob)/Lep(ob)* mice. *J Biol Chem* 277:19353-19357, 2002
18. You M, Matsumoto M, Pacold CM, Cho WK, Crabb DW: The role of AMP-activated protein kinase in the action of ethanol in the liver. *Gastroenterology* 127:1798-1808, 2004
19. Costet P, Legendre C, Moré J, Edgar A, Galtier P, Pineau T: Peroxisome proliferator-activated receptor alpha-isoform deficiency leads to progressive dyslipidemia with sexually dimorphic obesity and steatosis. *J Biol Chem* 273:29577-29585, 1998
20. Ip E, Farrell GC, Robertson G, Hall P, Kirsch R, Leclercq I: Central role of PPARalpha-dependent hepatic lipid turnover in dietary steatohepatitis in mice. *Hepatology* 38:123-132, 2003
21. Nagy LE: Recent insights into the role of the innate immune system in the development of alcoholic liver disease. *Exp Biol Med* (Maywood) 228:882-890, 2003
22. Bird GL, Sheron N, Goka AK, Alexander GJ, Williams RS: Increased plasma tumor necrosis factor in severe alcoholic hepatitis. *Ann Intern Med* 112:917-920, 1990
23. Lin HZ, Yang SQ, Zeldin G, Diehl AM: Chronic ethanol consumption induces the production of tumor necrosis factor-alpha and related cytokines in liver and adipose tissue. *Alcohol Clin Exp Res* 22(5 Suppl):231S-237S, 1998
24. Pritchard MT, McMullen MR, Stavitsky AB, Cohen JI, Lin F, Medof ME, Nagy LE: Differential contributions of C3, C5, and decay-accelerating factor to ethanol-induced fatty liver in mice. *Gastroenterology* 132:1117-1126, 2007
25. Ji C, Deng Q, Kaplowitz N: Role of TNF-alpha in ethanol-induced hyperhomocysteinemia and murine alcoholic liver injury. *Hepatology* 40:442-451, 2004
26. Yin M, Wheeler MD, Kono H, Bradford BU, Gallucci RM, Luster MI, Thurman RG: Essential role of tumor necrosis factor alpha in alcohol-induced liver injury in mice. *Gastroenterology* 117:942-952, 1999
27. Lawler JF Jr, Yin M, Diehl AM, Roberts E, Chatterjee S: Tumor necrosis factor-alpha stimulates the maturation of sterol regulatory element binding protein-1 in human hepatocytes through the action of neutral sphingomyelinase. *J Biol Chem* 273:5053-5059, 1998
28. Endo M, Masaki T, Seike M, Yoshimatsu H: TNF-alpha induces hepatic steatosis in mice by enhancing gene expression of sterol regulatory element binding protein-1c (SREBP-1c). *Exp Biol Med* (Maywood) 232:614-621, 2007
29. Kang L, Sebastian BM, Pritchard MT, Pratt BT, Previs SF, Nagy LE: Chronic ethanol-induced insulin resistance is associated with macrophage infiltration into adipose tissue and altered expression of adipocytokines. *Alcohol Clin Exp Res* 31:1581-1588, 2007
30. El-Assal O, Hong F, Kim WH, Radaeva S, Gao B: IL-6-deficient mice are susceptible to ethanol-induced hepatic steatosis: IL-6 protects against ethanol-induced oxidative stress and mitochondrial permeability transition in the liver. *Cell Mol Immunol* 1:205-211, 2004
31. Hong F, Radaeva S, Pan HN, Tian Z, Veech R, Gao B: Interleukin 6 alleviates hepatic steatosis and ischemia/re-

- perfusion injury in mice with fatty liver disease. *Hepatology* 40;933-941, 2004
32. Horiguchi N, Wang L, Mukhopadhyay P, Park O, Jeong WI, Lafdil F, Osei-Hyiaman D, Moh A, Fu XY, Pacher P, Kunos G, Gao B: Cell type-dependent pro- and anti-inflammatory role of signal transducer and activator of transcription 3 in alcoholic liver injury. *Gastroenterology* 134;1148-1158, 2008
  33. Radaeva S, Wang L, Radaev S, Jeong WI, Park O, Gao B: Retinoic acid signaling sensitizes hepatic stellate cells to NK cell killing via upregulation of NK cell activating ligand RAE1. *Am J Physiol Gastrointest Liver Physiol* 293;G809-G816, 2007
  34. Jeong WI, Park O, Radaeva S, Gao B: STAT1 inhibits liver fibrosis in mice by inhibiting stellate cell proliferation and stimulating NK cell cytotoxicity. *Hepatology* 44;1441-1451, 2006
  35. Radaeva S, Sun R, Jaruga B, Nguyen VT, Tian Z, Gao B: Natural killer cells ameliorate liver fibrosis by killing activated stellate cells in NKG2D-dependent and tumor necrosis factor-related apoptosis-inducing ligand-dependent manners. *Gastroenterology* 130;435-452, 2006
  36. McClain CJ, Barve S, Deaciuc I, Kugelmas M, Hill D: Cytokines in alcoholic liver disease. *Semin Liver Dis* 19; 205-219, 1999
  37. Szabo G: Consequences of alcohol consumption on host defence. *Alcohol Alcohol* 34;830-841, 1999
  38. Akira S, Uematsu S, Takeuchi O: Pathogen recognition and innate immunity. *Cell* 124;783-801, 2006
  39. Seki E, Brenner DA: Toll-like receptors and adaptor molecules in liver disease: update. *Hepatology* 48;322-335, 2008
  40. Uesugi T, Froh M, Arteeel GE, Bradford BU, Thurman RG: Toll-like receptor 4 is involved in the mechanism of early alcohol-induced liver injury in mice. *Hepatology* 34;101-108, 2001
  41. Chow JC, Young DW, Golenbock DT, Christ WJ, Gusovsky F: Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction. *J Biol Chem* 274;10689-10692, 1999
  42. Visintin A, Mazzoni A, Spitzer JA, Segal DM: Secreted MD-2 is a large polymeric protein that efficiently confers lipopolysaccharide sensitivity to Toll-like receptor 4. *Proc Natl Acad Sci U S A* 98;12156-12161, 2001
  43. Wright SD, Tobias PS, Ulevitch RJ, Ramos RA: Lipopolysaccharide (LPS) binding protein opsonizes LPS-bearing particles for recognition by a novel receptor on macrophages. *J Exp Med* 170;1231-1241, 1989
  44. Uesugi T, Froh M, Arteeel GE, Bradford BU, Wheeler MD, Gäbele E, Isayama F, Thurman RG: Role of lipopolysaccharide-binding protein in early alcohol-induced liver injury in mice. *J Immunol* 168;2963-2969, 2002
  45. Yin M, Bradford BU, Wheeler MD, Uesugi T, Froh M, Goyert SM, Thurman RG: Reduced early alcohol-induced liver injury in CD14-deficient mice. *J Immunol* 166;4737-4742, 2001
  46. Nanji AA, Khettry U, Sadrzadeh SM: Lactobacillus feeding reduces endotoxemia and severity of experimental alcoholic liver (disease). *Proc Soc Exp Biol Med* 205;243-247, 1994
  47. Kawai T, Akira S: The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 11;373-384, 2010
  48. Hritz I, Mandrekar P, Velayudham A, Catalano D, Dolganiuc A, Kodys K, Kurt-Jones E, Szabo G: The critical role of toll-like receptor (TLR) 4 in alcoholic liver disease is independent of the common TLR adapter MyD88. *Hepatology* 48;1224-1231, 2008
  49. Bautista AP: Neutrophilic infiltration in alcoholic hepatitis. *Alcohol* 27;17-21, 2002
  50. Kono H, Rusyn I, Yin M, Gäbele E, Yamashina S, Dikalova A, Kadiiska MB, Connor HD, Mason RP, Segal BH, Bradford BU, Holland SM, Thurman RG: NADPH oxidase-derived free radicals are key oxidants in alcohol-induced liver disease. *J Clin Invest* 106;867-872, 2000
  51. Thakur V, Pritchard MT, McMullen MR, Wang Q, Nagy LE: Chronic ethanol feeding increases activation of NADPH oxidase by lipopolysaccharide in rat Kupffer cells: role of increased reactive oxygen in LPS-stimulated ERK1/2 activation and TNF-alpha production. *J Leukoc Biol* 79;1348-1356, 2006
  52. Wheeler MD, Kono H, Yin M, Rusyn I, Froh M, Connor HD, Mason RP, Samulski RJ, Thurman RG: Delivery of the Cu/Zn-superoxide dismutase gene with adenovirus reduces early alcohol-induced liver injury in rats. *Gastroenterology* 120;1241-1250, 2001
  53. Dasu MR, Devaraj S, Zhao L, Hwang DH, Jialal I: High glucose induces toll-like receptor expression in human monocytes: mechanism of activation. *Diabetes* 57;3090-3098, 2008
  54. Park HS, Jung HY, Park EY, Kim J, Lee WJ, Bae YS: Cutting edge: direct interaction of TLR4 with NAD(P)H oxidase 4 isozyme is essential for lipopolysaccharide-induced production of reactive oxygen species and activation of NF-kappa B. *J Immunol* 173;3589-3593, 2004
  55. Friedman SL: Hepatic stellate cells: protean, multifunctional, and enigmatic cells of the liver. *Physiol Rev* 88;125-172, 2008
  56. Melhem A, Muhanna N, Bishara A, Alvarez CE, Ilan Y, Bishara T, Horani A, Nassar M, Friedman SL, Safadi R: Anti-fibrotic activity of NK cells in experimental liver injury through killing of activated HSC. *J Hepatol* 45;60-71, 2006
  57. Taimr P, Higuruchi H, Kocova E, Rippe RA, Friedman S, Gores GJ: Activated stellate cells express the TRAIL receptor-2/death receptor-5 and undergo TRAIL-mediated apoptosis. *Hepatology* 37;87-95, 2003
  58. Baroni GS, D'Ambrosio L, Curto P, Casini A, Mancini R, Jezequel AM, Benedetti A: Interferon gamma decreases hepatic stellate cell activation and extracellular matrix deposition in rat liver fibrosis. *Hepatology* 23;1189-1199, 1996
  59. Park O, Jeong WI, Wang L, Wang H, Lian ZX, Gershwin ME, Gao B: Diverse roles of invariant natural killer T cells in liver injury and fibrosis induced by carbon tetrachloride. *Hepatology* 49;1683-1694, 2009
  60. Laso FJ, Madruga JI, Girón JA, López A, Ciudad J, San Miguel JF, Alvarez-Mon M, Orfao A: Decreased natural killer cytotoxic activity in chronic alcoholism is associated with alcohol liver disease but not active ethanol consumption. *Hepatology* 25;1096-1100, 1997

61. Pan HN, Sun R, Jaruga B, Hong F, Kim WH, Gao B: Chronic ethanol consumption inhibits hepatic natural killer cell activity and accelerates murine cytomegalovirus-induced hepatitis. *Alcohol Clin Exp Res* 30;1615-1623, 2006
  62. Szabo G, Mandrekar P: A recent perspective on alcohol, immunity, and host defense. *Alcohol Clin Exp Res* 33;220-232, 2009
  63. Szabo G, Mandrekar P, Girouard L, Catalano D: Regulation of human monocyte functions by acute ethanol treatment: decreased tumor necrosis factor-alpha, interleukin-1 beta and elevated interleukin-10, and transforming growth factor-beta production. *Alcohol Clin Exp Res* 20;900-907, 1996
  64. Jaruga B, Hong F, Kim WH, Sun R, Fan S, Gao B: Chronic alcohol consumption accelerates liver injury in T cell-mediated hepatitis: alcohol dysregulation of NF-kappaB and STAT3 signaling pathways. *Am J Physiol Gastrointest Liver Physiol* 287;G471-G479, 2004
  65. Minagawa M, Deng Q, Liu ZX, Tsukamoto H, Dennert G: Activated natural killer T cells induce liver injury by Fas and tumor necrosis factor-alpha during alcohol consumption. *Gastroenterology* 126;1387-1399, 2004
-