

Analysis of causes of injuries among children in Daegu, Korea

Geol Ha, M.D.¹, Man Joong Jeon, M.D.², and Joon Sakong, M.D.²

Departments of Medicine¹, Preventive Medicine and Public Health², College of Medicine, Yeungnam University, Daegu, Korea

Received: 4 June 2010, Revised: 14 July 2010

Accepted: 4 August 2010

Corresponding author: Joon Sakong, M.D.

Department of Preventive Medicine and Public Health, College of Medicine, Yeungnam University, 317-1 Daemyung-dong, Nam-gu, Daegu 705-717, Korea

Tel: +82.53-620-4614, Fax: +82.53-653-2061

E-mail: jsakong@med.yu.ac.kr

Copyright © 2010 by The Korean Pediatric Society

Purpose: The purpose of this study was to determine the mechanisms and exact causes of injuries to children by analyzing the medical records of children who visited emergency medical centers.

Methods: We reviewed the medical records of 7,114 pediatric patients (aged below 15 years) who visited the emergency care center (ECC) of a Yeungnam university hospital in Daegu between January 2007 and December 2009.

Results: Among the pediatric patients who visited the ECC, 1,526 (27.2%) were admitted for injuries (boys, 979 [64.2%] and girls, 547 [35.8%]); the boy-to-girl ratio in this study population was 1.8:1. The proportion of boys was higher than that of girls over all age ranges, except for children aged less than 1 year. The highest incidence of injury due to slipping (20.5%) was noted in the 1-year-old group, and nearly half the total number of cases (49.3%) involved children who were under 4 years of age. Of the sites of injury, 999 involved the head and face (73.7%) and 134 involved the hand and wrist (9.9%). Seventy-one injuries (5.7%) occurred on the bed, 70 (5.6%) at the door, and 67 (5.3%) on the stairs. Of the approximately 250 types of injuries classified by cause, 17 accounted for 50.4% of all the injuries.

Conclusion: The cause of injury could be attributed to approximately 20 causes in more than half the cases of pediatric injuries. A classification of the types of injuries and the development of appropriate preventive strategies can help avoid such injuries.

Key words: Wounds and Injuries, Child, Emergency medical centers

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Injuries in children are a medically important subject because injuries account for the largest proportion of the causes of children's deaths. According to data from the Statistics Korea (Korean national statistical office), of the 10 major causes of children's deaths,

3 to 4 causes related to unexpected accidents were responsible for deaths of children¹. For children 1-9 years of age, the number of deaths due to injuries is comparable with the number of deaths due to all other causes. Moreover, for adolescents 10-19 years of age, unexpected accidents are still the leading causes of deaths, except for suicide². For global deaths of children (950,000 children) under

18 years of age per annum, injury is the leading cause of death, and death due to unintentional injuries accounts for approximately 90% of all causes of deaths³. In United States, deaths in childhood from injuries exceed childhood deaths from all other causes combined⁴. As injuries and children's deaths are closely related, reduction of the number of unintentional injuries in children can considerably decrease overall child mortality.

Although there have been many studies on child accidents in Korea, only a few studies have dealt with the 'causes' of child accidents. Thus, more studies on the causes of child accidents are required. Also, many studies have focused on the mechanisms of accidents, but few have addressed the detailed circumstances of accidents, except for a study that was performed with the use of a survey questionnaire. In this context, the detailed causes of accidents were also investigated, in addition to the general mechanisms of the accidents.

Because there are few training materials for children with respect to the prevention of accidents, practical guidelines for the prevention of accidents are required. This study was performed to establish an appropriate preventative measure by identifying the exact circumstances under which childhood injuries occur based on the medical records of the ECC.

Materials and methods

Of the 7,114 pediatric patients under 15 years of age who visited the ECC of Yeungnam university medical center between January 1, 2007 and December 31, 2008, 1,934 cases of children who visited the ECC due to injuries or burns were reviewed. Among the 1,934 patients, 408 who were not eligible for inclusion in the study based on the predetermined criteria were excluded from the study and the emergency room outpatient records of the 1,526 children who were eligible were analyzed. The cases that were excluded were: ECC visits due to traffic accidents, inadequate records of the cause of the accident, accidents due to basal diseases (e.g., gait disturbance or Moyamoya disease), ECC visits due to assaults, revisits within 2 days, and absence of a medical chart, although the patient's name was listed on the computerized database received from the patient records department.

The computer-based data on the subject included the visit date, arrival time, discharge date, registration number, gender, age at the time of the visit, and other necessary information that were not available in the computer-based data and were obtained from the relevant medical charts. The causes of the accidents that were determined from the charts were classified according to the mechanism of injury (full version) of ICECI (international

classification of external causes of injury) in 2003 (Table 1)⁵. The mechanism of an accident that was listed on the ICECI classification, but was not among the subjects of this study, was excluded from the results. The classification of the injured body parts followed S00-T98 of KCD-5 (XIX. injury, poisoning, and certain other consequences of external causes). Also, the detailed causes of the accidents were organized using the authors' classification chart.

The frequency of the accidents by age and gender, the associations between gender and the mechanism of the accident, age and the mechanism of the accident, time or season and the mechanism of the accident, the ECC visit time and the mechanism of the accident, the injured body part and the mechanism of the accident, and the detailed causes of the accidents were examined and organized for use as basic data for establishing proper preventative measures.

The data collected was assessed with a frequency analysis for which SPSS Statistics 17.0 was used.

Results

The proportion of pediatric patients who visited the ECC for injuries due to causes other than traffic accidents among the total number of pediatric patients who visited the ECC was 27.2%.

The number of male and female pediatric patients were 979 (64.2%) and 547 (35.8%), respectively, with a male-to-female ratio

Table 1. Mechanism of Injury and ICECI

Mechanism of injury	ICECI
Blunt force	1
Contact with object or animal	1.2
Contact with person	1.3
Crushing	1.4
Falling, stumbling, jumping, pushed	1.5
Slipping down	1.5.1-1.5.3
Falling down	1.5.4-1.5.6
<1 m	1.5.4
≥1 m	1.5.5
Unspecified	1.5.6
On stairs and steps	1.5.7
Piercing, penetrating force, abrasion	2
Scratching, cutting, tearing, severing	2.1
Puncturing, stabbing	2.2
Biting, stinging, in venomating	2.3
Thermal mechanism	4
Heating	4.1
Cooling	4.2
Poisoning, chemical injury	6
Other mechanism	98
Foreign body	98.1
Electricity	98.2

Abbreviations: ICECI, international classification of external causes of injury cited from reference 6

of 1.8:1. The greatest number of accidents occurred among children 1 year of age [313 accidents (20.5%)]. Nearly one-half of all of the accidents occurred in children under 4 years of age [753 accidents (49.3%)]. Before 15 years of age, more accidents occurred in boys than in girls, except under 1 year of age, for which the ratio of boys to girls was 1:1.1 (Table 2).

Except for crushing and poisoning, the accidents occurred more often in boys than in girls, regardless of the mechanism of the accident and the age. More blunt injuries, falls (including down the stairs), penetrations, lacerations, or animal or insect bites occurred in boys than in girls, whereas more slips, crushing, burns, or swallowing of foreign bodies occurred in girls than in boys. The number of children who visited the ECC due to poisoning or electrical burns was very small, and thus insignificant. Slips were the leading causes of accidents, regardless of the child's gender, with 448 cases (29.4%). No frostbite occurred (Table 3).

Regardless of the cause of the accident, a large number of accidents occurred at about 1 year of age. There were 9 collisions with humans over 11 years of age (90%). Of all the falls, the proportion of falls from a height over 1 meter in children 0-6 years of age was relatively low. For falls down the stairs, the largest number occurred in children 1 year of age (20 cases [29.9%]). Thirty-five burns (46.1%) occurred in children 1 year of age, which is a high incidence rate considering that 20.5% of all the accidents occurred in children 1 year of age. Electrical burns occurred in 3 children 1 year of age, and poisoning occurred in 3 children 3 years of age. For children under 1 year of age, falls accounted for the largest proportion (31.7%), followed by burns (18.3%), and blunt injuries (16.3%). For children 1 year of age, the proportion

Table 2. Distribution of the Injured Children Visiting the Emergency Care Center (ECC), January 1, 2007~December 31, 2008, by Age and Sex

Age (yrs.)	Male (%)	Female (%)	Total (%)
0	50 (5.1)	54 (9.9)	104 (6.8)
1	189 (19.3)	124 (22.7)	313 (20.5)
2	117 (12.0)	79 (14.4)	196 (12.8)
3	79 (8.1)	61 (11.2)	140 (9.2)
4	80 (8.2)	49 (9.0)	129 (8.5)
5	62 (6.3)	37 (6.8)	99 (6.5)
6	59 (6.0)	34 (6.2)	93 (6.1)
7	48 (4.9)	22 (4.0)	70 (4.6)
8	54 (5.5)	24 (4.4)	78 (5.1)
9	35 (3.6)	11 (2.0)	46 (3.0)
10	34 (3.5)	13 (2.4)	47 (3.1)
11	25 (2.6)	12 (2.2)	37 (2.4)
12	28 (2.9)	5 (0.9)	33 (2.2)
13	52 (5.3)	2 (0.4)	54 (3.5)
14	38 (3.9)	7 (1.3)	45 (2.9)
15	29 (3.0)	13 (2.4)	42 (2.8)
Total (%)	979 (64.2)	547 (35.8)	1526 (100.0)

of slips was the largest (32.3%), which indicates that there was a considerable difference in the mechanisms of the accidents that occurred between children 0 and 1 year of age (Table 4). The proportion of children inflicted with injuries was highest (8.2-10.4%) from March to September, and under 8% from January to February and October to December, regardless of the type of accident. Most of the accidents occurred from March to September, except that crushing occurred most often in January (19.2%), respectively (Table 5).

There was no considerable difference in the seasonal distribution, regardless of mechanisms of injury (Table 6).

The largest number of visits occurred between 18:00 and 20:00 (25.9%) (Table 7). Nine hundred ninety-nine cases (73.7%) involved the head and face, followed by the hand and wrist (9.9%) (Table 8).

Regarding the detailed causes of accidents, the largest number of accidents occurred in bed [71 cases (5.7%)]. Of the more than 250 detailed causes of accidents, 17 types accounted for 50.4% of all accidents (Table 9).

Regarding the relationship between the mechanism of the accident and the detailed cause of the accident, blunt injuries caused by collision occurred mostly due to collisions with structures or furniture in the house, and blunt injuries caused by collision with other people occurred mostly due to sports activities. For crushing injuries, 31 cases (59.6%) occurred at the door. Although slips

Table 3. Distribution of Injured Children by Sex and Mechanism of Injury

Mechanism of injury	Male (%)	Female (%)	Total (%)
Blunt force	244 (24.9)	124 (22.7)	368 (24.1)
Contact with object or animal	212 (21.7)	94 (17.2)	306 (20.1)
Contact with person	9 (0.9)	1 (0.2)	10 (0.7)
Crushing	23 (2.3)	29 (5.3)	52 (3.4)
Falling, stumbling, jumping, pushed	519 (53.0)	270 (49.4)	789 (51.7)
Slipping down	275 (28.1)	173 (31.6)	448 (29.4)
Falling down	196 (20.0)	78 (14.3)	274 (18.0)
<1 m	89 (9.1)	47 (8.6)	136 (8.9)
≥1 m	51 (5.2)	11 (2.0)	62 (4.1)
Unspecified	56 (5.7)	20 (3.7)	76 (5.0)
On stairs and steps	48 (4.9)	19 (3.5)	67 (4.4)
Piercing, penetrating force, abrasion	81 (8.3)	38 (6.9)	119 (7.8)
Scratching, cutting, tearing, severing	51 (5.2)	25 (4.6)	76 (5.0)
Puncturing, stabbing	20 (2.0)	9 (1.6)	29 (1.9)
Biting, stinging, invenomating	10 (1.0)	4 (0.7)	14 (0.9)
Thermal mechanism	40 (4.1)	36 (6.6)	76 (5.0)
Heating	40 (4.1)	36 (6.6)	76 (5.0)
Cooling	0 (0)	0 (0)	0 (0)
Poisoning, chemical injury	1 (0.1)	2 (0.4)	3 (0.2)
Other mechanism	94 (9.6)	77 (14.1)	171 (11.2)
Foreign body	91 (9.3)	77 (14.1)	168 (11.0)
Electricity	3 (0.3)	0 (0)	3 (0.2)
Total (%)	979 (64.2)	547 (35.8)	1,526 (100.0)

Table 4. Distribution of Injured Children by Age and Mechanism of Injury

Mechanism of injury	Age (yrs.)															Total (%)
	0 (%)	1 (%)	2 (%)	3 (%)	4 (%)	5 (%)	6 (%)	7 (%)	8 (%)	9 (%)	10 (%)	11 (%)	12 (%)	13 (%)	14 (%)	
Burnt force	19 (16.3)	74 (23.6)	47 (24.0)	34 (24.3)	24 (16.6)	28 (28.3)	22 (23.7)	14 (20.0)	20 (25.6)	11 (23.9)	13 (27.7)	7 (18.9)	9 (27.3)	18 (33.3)	13 (28.9)	15 (35.7)
Contact with object or animal	17 (15.3)	64 (20.4)	43 (21.9)	28 (20.0)	20 (15.5)	19 (19.2)	18 (19.4)	9 (12.9)	16 (20.5)	9 (19.6)	13 (27.7)	3 (8.1)	8 (24.2)	15 (27.8)	10 (22.2)	15 (33.3)
Contact with person	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (5.4)	1 (3.0)	3 (5.6)	2 (4.4)	10 (20.1)
Cushing	2 (1.9)	10 (3.2)	4 (2.0)	6 (4.3)	4 (3.1)	9 (9.1)	3 (3.2)	5 (7.1)	4 (5.1)	2 (4.3)	0 (0.0)	2 (5.4)	0 (0.0)	0 (0.0)	1 (2.2)	5 (10.7)
Falling, stumbling, jumping, pushed	45 (43.3)	166 (52.7)	107 (54.6)	80 (57.1)	73 (55.6)	45 (45.5)	53 (57.0)	34 (48.6)	42 (53.8)	25 (56.5)	20 (42.6)	18 (48.6)	19 (57.6)	23 (42.6)	21 (46.7)	18 (42.9)
Slipping down	11 (10.6)	101 (32.3)	72 (36.7)	44 (31.4)	41 (31.8)	26 (26.3)	24 (25.8)	22 (31.4)	16 (34.8)	16 (34.8)	11 (23.4)	13 (35.1)	11 (33.3)	15 (27.8)	7 (15.6)	44 (81.2)
Falling down	33 (31.7)	44 (14.1)	27 (13.8)	28 (20.0)	25 (19.4)	16 (16.2)	26 (28.0)	10 (14.3)	13 (16.7)	9 (19.6)	8 (17.0)	5 (13.5)	7 (21.2)	5 (9.3)	12 (26.7)	6 (14.3)
< 1 m	22 (21.2)	29 (9.3)	17 (8.7)	15 (10.7)	18 (14.0)	7 (7.1)	3 (4.3)	4 (5.4)	0 (0.0)	1 (2.7)	1 (2.1)	1 (2.7)	0 (0.0)	1 (1.9)	4 (8.9)	13 (28.9)
> 1 m	5 (4.8)	8 (2.6)	5 (2.6)	6 (4.3)	4 (3.1)	1 (1.0)	2 (2.2)	4 (5.7)	5 (6.4)	3 (6.5)	1 (2.1)	1 (2.7)	2 (6.1)	3 (5.6)	6 (13.3)	6 (14.3)
Unspecified	6 (5.8)	7 (2.2)	5 (2.6)	7 (5.0)	3 (2.3)	8 (8.1)	3 (3.2)	3 (4.3)	4 (5.1)	6 (13.0)	6 (12.8)	3 (7.7)	5 (15.2)	1 (1.9)	2 (4.4)	0 (0.0)
On stairs and steps	1 (1.0)	20 (6.4)	8 (4.1)	8 (5.7)	7 (5.4)	3 (3.0)	3 (3.2)	2 (2.9)	4 (5.1)	1 (2.2)	1 (2.1)	0 (0.0)	1 (3.0)	3 (5.6)	2 (4.4)	6 (13.3)
Piercing, penetrating force, abrasion	6 (5.8)	15 (4.8)	7 (3.6)	3 (2.1)	9 (7.0)	8 (8.1)	10 (10.8)	14 (20.0)	9 (11.5)	4 (8.7)	5 (10.6)	9 (24.3)	3 (9.1)	7 (13.0)	8 (17.8)	2 (4.8)
Scratching, cutting, tearing, severing	4 (3.8)	11 (3.5)	3 (1.5)	0 (0.0)	7 (5.4)	3 (3.0)	5 (5.4)	10 (14.3)	5 (6.4)	3 (6.5)	4 (8.5)	5 (13.5)	2 (6.1)	6 (11.1)	2 (4.8)	7 (15.0)
Puncturing, stabbing	1 (1.0)	4 (1.3)	2 (1.0)	3 (2.1)	1 (0.8)	2 (2.0)	3 (4.3)	1 (1.4)	1 (1.3)	1 (2.2)	1 (2.1)	4 (10.8)	1 (3.0)	2 (4.4)	1 (2.2)	29 (61.9)
Biting, stinging, insectbaiting	1 (1.0)	0 (0.0)	2 (1.0)	0 (0.0)	1 (0.8)	3 (3.0)	2 (2.2)	1 (1.4)	3 (3.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.2)	14 (30.0)
Thermal mechanism	19 (18.3)	35 (11.2)	5 (2.6)	2 (1.4)	4 (3.1)	1 (1.0)	1 (1.1)	0 (0.0)	0 (0.0)	3 (6.5)	2 (4.3)	0 (0.0)	1 (3.0)	2 (3.7)	0 (0.0)	7 (15.0)
Heating	19 (18.3)	35 (11.2)	5 (2.6)	2 (1.4)	4 (3.1)	1 (1.0)	1 (1.1)	0 (0.0)	0 (0.0)	3 (6.5)	2 (4.3)	0 (0.0)	1 (3.0)	2 (3.7)	0 (0.0)	7 (15.0)
Cooling	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Poisoning, chemical injury	0 (0.0)	0 (0.0)	0 (0.0)	3 (2.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (6.5)
Other mechanism	15 (14.4)	24 (7.7)	30 (15.3)	18 (12.9)	19 (14.7)	17 (17.2)	8 (11.4)	8 (11.4)	7 (9.0)	2 (4.3)	7 (14.9)	3 (8.1)	1 (3.0)	4 (7.4)	3 (6.7)	6 (14.3)
Foreign body	15 (14.4)	24 (7.7)	30 (15.3)	18 (12.9)	19 (14.7)	17 (17.2)	8 (11.4)	8 (11.4)	7 (9.0)	2 (4.3)	7 (14.9)	3 (8.1)	1 (3.0)	4 (7.4)	3 (6.7)	6 (14.3)
Electricity	0 (0.0)	3 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (6.5)
Total (%)	104 (6.8)	313 (20.5)	196 (12.8)	140 (9.2)	129 (8.5)	99 (6.5)	98 (6.1)	70 (4.6)	78 (5.1)	46 (3.0)	47 (3.1)	37 (2.4)	38 (2.2)	38 (2.2)	54 (3.5)	45 (2.9)

Table 5. Distribution of Injured Children by Month and Mechanism of Injury

Mechanism of injury	Month												Total (%)
	Jan (%)	Feb (%)	Mar (%)	Apr (%)	May (%)	Jun (%)	Jul (%)	Aug (%)	Sep (%)	Oct (%)	Nov (%)	Dec (%)	
Burnt force	35 (31.5)	21 (18.4)	39 (31.2)	28 (20.7)	35 (22.0)	29 (23.2)	36 (24.8)	38 (25.2)	40 (27.0)	20 (17.1)	19 (22.1)	28 (25.5)	368 (24.1)
Contact with object or animal	23 (20.7)	17 (14.9)	32 (25.6)	21 (15.6)	29 (18.2)	24 (19.2)	30 (20.7)	34 (22.5)	36 (24.3)	19 (16.2)	14 (16.3)	27 (24.5)	306 (20.1)
Contact with person	2 (1.8)	0 (0.0)	4 (3.5)	1 (0.7)	0 (0.0)	1 (0.8)	1 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	10 (0.7)
Cushing	10 (9.0)	4 (3.5)	3 (2.4)	6 (4.4)	6 (3.8)	4 (3.2)	5 (3.4)	4 (2.6)	4 (2.7)	1 (0.9)	4 (4.7)	1 (0.9)	52 (3.4)
Falling, stumbling, jumping, pushed	48 (43.2)	60 (52.6)	57 (45.6)	73 (54.1)	87 (54.7)	70 (56.0)	81 (55.9)	75 (49.7)	75 (50.7)	67 (57.3)	40 (46.5)	56 (50.9)	789 (51.7)
Slipping down	28 (25.2)	41 (36.0)	32 (25.6)	39 (28.9)	42 (26.4)	39 (31.2)	44 (30.3)	44 (29.1)	43 (29.1)	36 (30.8)	18 (20.8)	42 (38.2)	448 (29.4)
Falling down	15 (13.5)	14 (12.3)	20 (16.0)	28 (20.7)	40 (25.2)	23 (18.4)	32 (22.1)	24 (15.9)	24 (16.2)	24 (20.5)	19 (22.1)	11 (10.0)	274 (18.0)
< 1 m	9 (8.1)	10 (8.8)	15 (12.0)	15 (11.1)	19 (11.9)	11 (8.8)	14 (9.7)	11 (7.3)	8 (5.4)	10 (8.5)	7 (8.1)	7 (6.4)	136 (8.9)
> 1 m	3 (2.7)	0 (0.0)	3 (2.4)	8 (5.9)	7 (4.4)	5 (4.0)	10 (6.9)	7 (4.6)	3 (2.0)	7 (6.0)	6 (7.0)	3 (2.7)	62 (4.1)
Unspecified	3 (2.7)	4 (3.5)	2 (1.6)	5 (3.7)	8 (5.1)	7 (5.6)	8 (5.5)	6 (4.0)	13 (8.8)	7 (6.0)	7 (8.1)	1 (0.9)	76 (5.0)
On stairs and steps	5 (4.5)	5 (4.4)	5 (4.0)	6 (4.4)	5 (3.1)	8 (6.4)	5 (3.4)	7 (4.6)	8 (5.4)	7 (6.0)	3 (3.5)	3 (2.7)	67 (4.4)
Piercing, penetrating force, abrasion	8 (7.2)	12 (10.5)	12 (9.6)	12 (8.9)	10 (6.3)	9 (7.2)	9 (6.2)	11 (7.9)	14 (9.5)	6 (5.1)	7 (8.1)	8 (7.3)	119 (7.8)
Scratching, cutting, tearing, severing	6 (5.4)	7 (6.1)	7 (5.6)	9 (6.7)	7 (4.4)	5 (4.0)	5 (3.4)	5 (3.4)	9 (6.1)	5 (4.3)	2 (2.3)	5 (4.3)	76 (5.0)
Puncturing, stabbing	1 (0.9)	3 (2.6)	4 (3.2)	1 (0.7)	3 (1.9)	4 (3.2)	2 (1.4)	0 (0.0)	1 (0.7)	1 (0.9)	5 (5.8)	4 (3.6)	29 (1.9)
Biting, stinging, insectbaiting	1 (0.9)	2 (1.8)	1 (0.8)	2 (1.5)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.7)	4 (2.7)	0 (0.0)	0 (0.0)	0 (0.0)	14 (0.9)
Thermal mechanism	5 (4.5)	4 (3.5)	8 (6.4)	6 (4.4)	8 (5.0)	2 (1.6)	5 (3.4)	10 (6.6)	7 (4.7)	9 (7.7)	6 (7.0)	6 (5.5)	76 (5.0)
Heating	5 (4.5)	4 (3.5)	8 (6.4)	6 (4.4)	8 (5.0)	2 (1.6)	5 (3.4)	10 (6.6)	7 (4.7)	9 (7.7)	6 (7.0)	6 (5.5)	76 (5.0)
Cooling	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.4)	0 (0.0)	0 (0.0)	0 (0.0)	3 (0.2)
Poisoning, chemical injury	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other mechanism	15 (13.5)	17 (14.9)	9 (7.2)	16 (11.9)	19 (11.9)	15 (12.0)	14 (9.7)	15 (9.9)	10 (6.8)	15 (12.8)	14 (16.3)	12 (10.9)	171 (11.2)
Foreign body	15 (13.5)	17 (14.9)	9 (7.2)	16 (11.9)	19 (11.9)	15 (12.0)	13 (9.0)	15 (9.9)	9 (6.1)	15 (12.8)	13 (15.1)	12 (10.9)	168 (11.0)
Electricity	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	3 (0.2)
Total (%)	111 (7.3)	114 (7.5)	125 (8.2)	135 (8.9)	159 (10.4)	125 (8.2)	145 (9.5)	151 (9.9)	148 (9.7)	117 (7.7)	86 (5.6)	110 (7.2)	1526 (100.0)

%, n/monthx100, %; n/totalx100

Table 6. Distribution of Injured Children by Four Seasons and Mechanism of Injury

Mechanism of injury	Season				Total (%)
	Spring*	Summer†	Autumn‡	Winter§	
Blunt force	102 (24.3)	103 (24.5)	79 (22.5)	84 (25.1)	368 (24.1)
Contact with object or animal	82 (19.6)	88 (20.9)	69 (19.7)	67 (20.0)	306 (20.1)
Contact with person	5 (1.2)	2 (0.5)	1 (0.3)	2 (0.6)	10 (0.7)
Crushing	15 (3.6)	13 (3.1)	9 (2.6)	15 (4.5)	52 (3.4)
Falling, stumbling, jumping, pushed	217 (51.8)	226 (53.7)	182 (51.9)	164 (49.0)	789 (51.7)
Slipping down	113 (27.0)	127 (30.2)	97 (27.6)	111 (33.1)	448 (29.4)
Falling down	88 (21.0)	79 (20.9)	67 (25.1)	40 (26.3)	274 (5.8)
<1 m	49 (11.7)	36 (8.6)	25 (7.1)	26 (7.8)	136 (8.9)
≥1 m	18 (4.3)	22 (5.2)	16 (4.6)	6 (1.8)	62 (4.1)
Unspecified	21 (5.0)	21 (5.0)	26 (7.4)	8 (2.4)	76 (5.0)
On stairs and steps	16 (3.8)	20 (4.8)	18 (5.1)	13 (3.9)	67 (4.4)
Piercing, penetrating force, abrasion	34 (8.1)	30 (7.1)	27 (7.7)	28 (8.4)	119 (7.8)
Scratching, cutting, tearing, severing	23 (5.5)	21 (5.0)	16 (4.6)	16 (4.8)	76 (5.0)
Puncturing, stabbing	8 (1.9)	6 (1.4)	7 (2.0)	8 (2.4)	29 (1.9)
Biting, stinging, inenoming	3 (0.7)	3 (0.7)	4 (1.1)	4 (1.2)	14 (0.9)
Thermal mechanism	22 (5.3)	17 (4.0)	22 (6.3)	15 (4.5)	76 (5.0)
Heating	22 (5.3)	17 (4.0)	22 (6.3)	15 (4.5)	76 (5.0)
Cooling	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Poisoning, chemical injury	0 (0)	1 (0.2)	2 (0.6)	0 (0)	3 (0.2)
Other mechanism	44 (10.5)	44 (10.5)	39 (11.1)	44 (13.1)	171 (11.2)
Foreign body	44 (10.5)	43 (10.2)	37 (10.5)	44 (13.1)	168 (11.0)
Electricity	0 (0)	1 (0.2)	2 (0.6)	0 (0)	3 (0.2)
Total (%)	419 (100.0)	421 (100.0)	351 (100.0)	335 (100.0)	1526 (100.0)

*Spring: March, April, May

†Summer: June, July, August

‡Autumn: September, October, November

§Winter: December, January, February

Table 7. Distribution of Injured Children by Time and Mechanism of Injury

Mechanism of injury	Time (hour)								Total (%)
	0-2 (%*)	3-5 (%*)	6-8 (%*)	9-11 (%*)	12-14 (%*)	15-17 (%*)	18-20 (%*)	21-23 (%*)	
Blunt force	16 (24.2)	2 (16.7)	7 (35.0)	28 (18.9)	51 (22.7)	68 (23.6)	110 (27.8)	86 (23.1)	368 (24.1)
Contact with object or animal	15 (22.7)	2 (16.7)	6 (30.0)	25 (16.9)	43 (19.1)	59 (20.5)	87 (22.0)	69 (18.5)	306 (20.1)
Contact with person	1 (1.5)	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.9)	0 (0.0)	4 (1.0)	3 (0.8)	10 (0.7)
Crushing	0 (0.0)	0 (0.0)	1 (5.0)	3 (2.0)	6 (2.7)	9 (3.1)	19 (4.8)	14 (3.8)	52 (3.4)
Falling, stumbling, jumping, pushed	34 (51.5)	7 (58.3)	6 (30.0)	86 (58.1)	128 (56.9)	167 (58.0)	188 (47.6)	173 (46.5)	789 (51.7)
Slipping down	23 (34.8)	3 (25.0)	2 (10.0)	49 (33.1)	59 (26.2)	94 (32.6)	107 (27.1)	111 (29.8)	448 (29.4)
Falling down	11 (16.7)	4 (33.3)	4 (20.0)	27 (18.2)	53 (23.6)	53 (18.4)	71 (18.0)	51 (13.7)	274 (18.0)
<1 m	6 (9.1)	1 (8.3)	4 (20.0)	17 (11.5)	27 (12.0)	20 (6.9)	30 (7.6)	31 (8.3)	136 (8.9)
≥1 m	3 (4.5)	2 (16.7)	0 (0.0)	5 (3.4)	16 (7.1)	11 (3.8)	17 (4.3)	8 (2.2)	62 (4.1)
Unspecified	2 (3.0)	1 (8.3)	0 (0.0)	5 (3.4)	10 (4.4)	22 (7.6)	24 (6.1)	12 (3.2)	76 (5.0)
On stairs and steps	0 (0.0)	0 (0.0)	0 (0.0)	10 (6.8)	16 (7.1)	20 (6.9)	10 (2.5)	11 (3.0)	67 (4.4)
Piercing, penetrating force, abrasion	4 (6.1)	1 (8.3)	1 (5.0)	12 (8.1)	20 (8.9)	21 (7.3)	27 (6.8)	33 (8.9)	119 (7.8)
Scratching, cutting, tearing, severing	3 (4.5)	1 (8.3)	1 (5.0)	6 (4.1)	15 (6.7)	13 (4.5)	19 (4.8)	18 (4.8)	76 (5.0)
Puncturing, stabbing	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.4)	5 (2.2)	5 (1.7)	7 (1.8)	10 (2.7)	29 (1.9)
Biting, stinging, inenoming	1 (1.5)	0 (0.0)	0 (0.0)	4 (2.7)	0 (0.0)	3 (1.0)	1 (0.3)	5 (1.3)	14 (0.9)
Thermal mechanism	4 (6.1)	0 (0.0)	4 (20.0)	5 (3.4)	11 (4.9)	7 (2.4)	28 (7.1)	17 (4.6)	76 (5.0)
Heating	4 (6.1)	0 (0.0)	4 (20.0)	5 (3.4)	11 (4.9)	7 (2.4)	28 (7.1)	17 (4.6)	76 (5.0)
Cooling	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Poisoning, chemical injury	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	1 (0.3)	1 (0.3)	3 (0.2)
Other mechanism	8 (12.1)	2 (16.7)	2 (10.0)	17 (11.5)	15 (6.7)	24 (8.3)	41 (10.4)	62 (16.7)	171 (11.2)
Foreign body	8 (12.1)	2 (16.7)	2 (10.0)	17 (11.5)	15 (6.7)	22 (7.6)	41 (10.4)	61 (16.4)	168 (11.0)
Electricity	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.7)	0 (0.0)	1 (0.3)	3 (0.2)
Total (%)	66 (4.3)	12 (0.8)	20 (1.3)	148 (9.7)	225 (14.7)	288 (18.9)	395 (25.9)	372 (24.4)	1526 (100.0)

*%: n/time×100

†%: n/total×100

Table 9. Distribution of Injured Children by Specific Cause of Injury

Cause of injury	N (%)
Bed	71 (5.7)
Door	70 (5.6)
Stairs	67 (5.3)
Hot water	56 (4.5)
Bicycle	48 (3.8)
Bathroom	48 (3.8)
Chair	46 (3.7)
Desk	40 (3.2)
Fish bone	39 (3.1)
Table	33 (2.6)
Dining table (short legged)	23 (1.8)
Soccer	19 (1.5)
Sofa	18 (1.4)
Slide	17 (1.4)
Piece of broken glass	17 (1.4)
Playground	16 (1.4)
Dining table	16 (1.3)
Total (%)	1255* (100.0)

*Cases of unknown specific causes: excluded

accounted for 29.4% of all accidents (448 cases), the exact causes were not identified in most cases (Table 10).

Discussion

The total number of children who visited the ECCs was 7,114 during the 2-year period, 1,526 (27.2%) of whom visited the ECC due to accidents other than traffic accidents. A previous study conducted at the same ECC from January 1, 2001 to December 31, 2002 showed, however, that the proportion of children who visited the ECC due to accidents other than traffic accidents was 23.0%⁶. Based on the increased proportion of children who visited the ECC for accidents other than traffic accidents, overcrowding of the emergency room and childhood accidents were associated.

Similar to the results of other studies, the ratio of boys to girls who visited the ECC was 1.8:1, which can be attributed to the characteristics of boys who have strong curiosity and are very active⁶⁻²⁰. Notably, under 15 years of age, the proportion of boys to girls was higher except under 1 year of age, at which age the ratio of boys to girls was 1:1.1. This seems to be related to the developmental process of children. There is little difference in the amount of activity between boys and girls under 1 year of age, when they can start to walk due to improved motor ability. Although the frequency of slips, which is considered most closely related to physical activity, was the fourth highest in children under 1 year of age, it was the first and second highest in children over 1 year of age. Based on this finding, it is suggested that accidents due to physical activity can occur more in boys than in girls.

Although for children under 1 year of age, falls accounted for the largest proportion of injuries (31.7%), followed by burns (18.3%) and blunt injuries (16.3%), for children 1 year of age, slips accounted for the largest proportion of injuries (32.3%), which indicates that there is a big difference in the mechanisms of accidents between boys and girls. This also suggests that the ability to ambulate considerably affects the mechanism of accidents.

The age at which the largest number of accidents occurred was 1 year, which accounted for 20.5% (313 cases) of all accidents. Similar to one report, nearly one-half of all the accidents occurred under 4 years of age [753 cases (49.3%)]. Under 1 year of age, accidents mostly occur with parental supervision, and at 1 year of age accidents may occur without parental supervision because children can already move on their own. Thus, the incidence of accidents is higher at 1 year of age than under 1 year. In addition, at 12-18 months, children tend to explore the things around them¹, which may cause accidents.

Except for crushing and poisoning, more accidents occurred in boys than in girls in all the classifications, regardless of the mechanism of the accident. The proportions of blunt injuries, falls, falls down the stairs, penetrations, lacerations, and animal or insect bites was higher among the boys who visited the ECCs than among girls, whereas the proportions of slips, crushing, burns, and swallowing of foreign bodies were higher than among the boys. The number of children who visited the ECCs due to poisoning or electrical burns was small, and thus insignificant for this study.

Slips were the leading causes of accidents in boys and girls, with 448 cases (29.4%). Previous studies did not classify slips as among the separate causes of accidents⁹⁻¹¹ or classified them in the same category as other causes⁷, which made it difficult to compare the results of previous studies with those of this study. Two studies that were conducted relatively recently, however, showed that slips were the leading causes of accidents^{8,21}.

Regardless of the cause, accidents occurred at approximately 1 year of age, and 9 cases of collisions with other people (90%) occurred over 11 years of age. Those collisions with other people were related to sports activities.

The proportion of falls from a height over 1 m at age 0-6 years was low. This might have been because children at 0-6 years of age cannot climb to a height over 1 m in most cases.

The largest number of falls down the stairs 20 cases (29.9%) occurred at 1 year of age. This was because children at 18 months of age can go up the stairs by holding other people's hands¹. Before 18 months of age, children cannot go up the stairs on their own. It appears that the number of falls increases as they start to go up the stairs on their own. Incomplete motor function, unstable body

Table 8. Distribution of Injured Children by Part of Body and Mechanism of Injury

Mechanism of injury	Part of body										Total (%)				
	Head (%)	Neck (%)	Thorax (%)	Upper back (%)	Abdomen, pelvis (%)	Lower back (%)	Shoulder upper arm (%)	Elbow, forearm (%)	Wrist, hand (%)	Hip, thigh (%)		Knee, lower leg (%)	Ankle, foot (%)	Unknown (%)	Whole body (%)
Blunt force	293 (29.3)	1 (11.1)	0 (0.0)	0 (0.0)	4 (23.5)	0 (0.0)	1 (1.9)	0 (0.0)	45 (33.6)	0 (0.0)	7 (21.9)	17 (43.6)	0 (0.0)	0 (0.0)	368 (27.1)
Contact with object or animal	284 (28.4)	1 (11.1)	0 (0.0)	0 (0.0)	3 (17.6)	0 (0.0)	1 (1.9)	0 (0.0)	8 (6.0)	0 (0.0)	3 (9.4)	6 (15.4)	0 (0.0)	0 (0.0)	306 (22.6)
Contact with person	7 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.7)	0 (0.0)	1 (3.1)	1 (2.6)	0 (0.0)	0 (0.0)	10 (0.7)
Crushing	2 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	1 (5.9)	0 (0.0)	0 (0.0)	0 (0.0)	36 (26.9)	0 (0.0)	3 (9.4)	10 (25.6)	0 (0.0)	0 (0.0)	52 (3.8)
Falling, stumbling, jumping, pushed	635 (63.6)	4 (4.4)	2 (28.6)	1 (100)	6 (35.3)	2 (100)	44 (83.0)	36 (87.8)	23 (17.2)	6 (42.9)	19 (59.4)	8 (20.5)	3 (75.0)	0 (0.0)	789 (58.2)
Slipping down	377 (37.7)	3 (33.3)	0 (0.0)	1 (100)	4 (23.5)	0 (0.0)	20 (37.7)	15 (36.6)	10 (7.5)	3 (21.4)	10 (31.3)	5 (12.8)	0 (0.0)	0 (0.0)	448 (33.0)
Falling down	200 (20.0)	0 (0.0)	1 (14.3)	0 (0.0)	2 (11.8)	0 (0.0)	24 (45.3)	19 (46.3)	11 (8.2)	3 (21.4)	7 (21.9)	2 (5.1)	3 (75.0)	0 (0.0)	274 (20.2)
<1 m	109 (10.9)	0 (0.0)	0 (0.0)	0 (0.0)	1 (5.9)	1 (50.0)	8 (15.1)	8 (19.5)	3 (2.2)	1 (7.1)	3 (9.4)	0 (0.0)	2 (50.0)	0 (0.0)	136 (10.0)
≥1 m	42 (4.2)	0 (0.0)	1 (14.3)	0 (0.0)	1 (5.9)	1 (50.0)	8 (15.1)	8 (19.5)	2 (1.5)	1 (7.1)	1 (3.1)	0 (0.0)	1 (25.0)	0 (0.0)	62 (4.6)
Unspecified	49 (4.9)	0 (0.0)	0 (0.0)	0 (0.0)	1 (5.9)	0 (0.0)	8 (15.1)	8 (19.5)	6 (4.5)	1 (7.1)	3 (9.4)	0 (0.0)	0 (0.0)	0 (0.0)	76 (5.6)
On stairs and steps	58 (5.8)	1 (11.1)	1 (14.3)	0 (0.0)	1 (5.9)	0 (0.0)	8 (15.1)	2 (4.9)	2 (1.5)	1 (7.1)	2 (6.3)	1 (2.6)	0 (0.0)	0 (0.0)	67 (4.9)
Piercing, penetrating force, abrasion	57 (5.7)	1 (11.1)	1 (14.3)	0 (0.0)	1 (5.9)	0 (0.0)	1 (1.9)	2 (4.9)	43 (32.1)	3 (21.4)	3 (9.4)	6 (15.4)	0 (0.0)	0 (0.0)	119 (8.8)
Scratching, cutting, tearing, severing	25 (2.5)	1 (11.1)	1 (14.3)	0 (0.0)	1 (5.9)	0 (0.0)	0 (0.0)	2 (4.9)	38 (28.4)	1 (7.1)	1 (3.1)	6 (15.4)	0 (0.0)	0 (0.0)	76 (5.6)
Puncturing, stabbing	28 (2.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	29 (2.1)
Biting, stinging, insectbiting	4 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.9)	0 (0.0)	4 (3.0)	2 (14.3)	2 (6.3)	0 (0.0)	0 (0.0)	0 (0.0)	14 (1.0)
Thermal mechanism	13 (1.3)	3 (33.3)	4 (57.1)	0 (0.0)	6 (35.3)	0 (0.0)	7 (13.2)	3 (7.3)	20 (14.9)	5 (35.7)	3 (9.4)	8 (20.5)	1 (25.0)	3 (75.0)	76 (5.6)
Heating	13 (1.3)	3 (33.3)	4 (57.1)	0 (0.0)	6 (35.3)	0 (0.0)	7 (13.2)	3 (7.3)	20 (14.9)	5 (35.7)	3 (9.4)	8 (20.5)	1 (25.0)	3 (75.0)	76 (5.6)
Cooling	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Poisoning, chemical injury	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.1)
Other mechanism	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (2.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (0.2)
Foreign body	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Electricity	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (2.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (0.2)
Total (%)	999 (73.7)	9 (0.7)	7 (0.5)	1 (0.1)	17 (1.3)	2 (0.1)	53 (3.9)	41 (3.0)	134 (9.9)	14 (1.0)	32 (2.4)	39 (2.9)	4 (0.3)	4 (0.3)	1,356 (100.0)

*Foreign body; excluded, %: n/total×100, %: n/part of body×100

Table 10. Distribution of Injured Children by Frequency of Cause and Mechanism of Injury

Mechanism of injury	Frequency										Total (%)		
	1 (N, %)	2 (N, %)	3 (N, %)	4 (N, %)	5 (N, %)	6 (N, %)	7 (N, %)	8 (N, %)	9 (N, %)	10 (N, %)			
Blunt force													
Contact with object or animal	Table 22 (7.2)	Door 19 (6.2)	Dining table 17 (5.6)	Desk 16 (5.2)	Bed 14 (4.6)	Dining table 14 (4.6)	Baseball 12 (3.9)	Chair 11 (3.6)	Wall 10 (3.3)	Furniture 9 (2.9)	306 (20.1)		
Contact with person	Soccer 5 (50.0)	Taekwondo 2 (20.0)	Baseball 1 (10.0)	Kendo 1 (10.0)	Head 1 (10.0)	Plyground 1 (1.9)	Metal rod 1 (1.9)	Window 1 (1.9)	Mirabelle 1 (1.9)	Escalator 1 (1.9)	10 (0.7)		
Crushing	Door 31 (59.6)	Bicycle 2 (3.8)	Closet 2 (3.8)	Automatic door 2 (3.8)	Car 2 (3.8)	Plyground 3 (3.9)	Plyground 3 (3.9)	Plyground 1 (1.9)	Escalator 1 (1.9)	Escalator 1 (1.9)	52 (3.4)		
Falling, stumbling, jumping, pushed	Bathroom 47 (10.5)	Desk 19 (4.2)	Door 17 (3.8)	Bed 15 (3.3)	Slide 11 (2.5)	Table 11 (2.5)	Plyground 7 (1.6)	Window 6 (1.3)	Mirabelle 1 (1.9)	Escalator 1 (1.9)	448 (28.4)		
Slipping down													
Falling down	<1 m	Bed 36 (26.5)	Chair 29 (21.3)	Bicycle 19 (14.0)	Sofa 12 (8.8)	Toy car 2 (1.5)	Quick board 2 (1.5)	Toilet 2 (1.5)	Desk 2 (1.5)	Car 2 (1.5)	62 (4.1)		
	≥1 m	Bicycle 6 (9.7)	Cloud ladder 4 (6.5)	Second floor 4 (6.5)	Bed 4 (6.5)	Trolley 4 (6.5)	Plyground 3 (4.8)	Fences 3 (4.8)	Banister 2 (3.2)	Desk 2 (3.2)	76 (5.0)		
	Unspecified	Bicycle 20 (26.3)	Swing 4 (5.3)	Horizontal bar 3 (3.9)	Slide 3 (3.9)	Trampoline 3 (3.9)	Trolley 3 (3.9)	Plyground 2 (2.6)	Cloud ladder 2 (2.6)	Window 2 (2.6)	67 (4.4)		
	On stairs and steps	Stars 67 (100)									67 (4.4)		
Piercing, penetrating force, abrasion	Knife 17 (22.4)	Piece of broken glass 13 (17.1)	Glass window 5 (6.6)	Toy 4 (5.3)	Dish 3 (3.9)	Sessor 3 (3.9)	Door 2 (2.6)	Cup 2 (2.6)	Bathtub 1 (1.3)	Eyeglasses 1 (1.3)	76 (5.0)		
Scratching, cutting, tearing, severing	Ball bullet 6 (20.7)	Sessor 2 (6.9)	Tree branch 2 (6.9)	Rod 1 (3.4)	Wood toy 1 (3.4)	Door 1 (3.4)	Broomstick 1 (3.4)	Umbrella 1 (3.4)	Thermometer 1 (3.4)	Cotton swab 1 (3.4)	29 (1.9)		
Biting, stinging, insectbiting	Dog 10 (7.4)	Bug 2 (14.3)	Bee 1 (7.1)	Rabbit 1 (7.1)	-	-	-	-	-	-	14 (0.9)		
Thermal mechanism	How water 59 (77.6)	Steam cleaner 2 (2.6)	Electric rice cooker 2 (2.6)	Hot grill 2 (2.6)	Freezer 1 (1.3)	Faucet 1 (1.3)	Electric iron 1 (1.3)	Oven 1 (1.3)	Cooking oil 1 (1.3)	Tukdoagi 1 (1.3)	76 (5.0)		
Heating													
Cooling													
Poisoning, chemical injury	Sodium hypochlorite 1 (33.3)	Thinner 1 (33.3)	Mercury thermometer 1 (33.3)	-	-	-	-	-	-	-	3 (0.2)		
Other mechanism													
Foreign body	Fish bone 36 (21.4)	Con 13 (7.7)	Marble 12 (7.1)	Bean 10 (6.0)	Chicken bone 9 (5.4)	Baduk-piece 7 (4.2)	Slicker 3 (1.8)	Ball bullet 3 (1.8)	Piece of broken glass 3 (1.8)	Magnat 3 (1.8)	188 (11.0)		
Electricity	Electrical outlet 1 (33.3)	Chopstick 1 (33.3)	Mobile phone charger 1 (33.3)	-	-	-	-	-	-	-	3 (0.2)		
Total (%)											1,326 (100.0)		

%, n:mechanism ×100, %: n/total×100

posture while going up the stairs, and higher stairs than children's height can cause accidents.

For burns, 46.1% (35 cases) occurred at 1 year of age, which is a considerably high incidence rate, considering that the proportion of accidents that occurred at 1 year of age was 20.5% of all accidents. The largest number of burns occurred due to hot water. In this study, the meaning of hot water includes hot water in a cup, a pot, and a bowl. Considering that many burn accidents occurred due to parents' mistakes or children spilling hot water by mistake, parents should be cautious and hot water should be placed away from children to prevent burns. Two cases of burn accidents due to a steam cleaner also occurred.

Most of the mechanisms of the accidents occurred from March to September, except for crushing and pricking, with the largest number of crushing and pricking cases occurring in January (19.2%) and November (17.2%), respectively. Generally, the proportion of accidents related to physical activity was high. It is postulated that this was because there are more hours of outdoor activity when the weather is suitable and because days are longer from March to September. For the seasonal distribution of the accidents, more accidents occurred in spring and summer than in fall and winter. This appears to be related to the amount and length of physical activity, as in the case of the monthly distribution of the accidents. The number of slips slightly increased from 97 cases (21.7%) in the fall to 111 cases (4.8%) in the winter. Most of the crushing was indoors, which was unrelated to outdoor activity, and the largest number of crushing incidents occurred in winter, when there are more hours of indoor activity.

The time when the largest number of accidents (25.9%) occurred was from 18:00 to 20:00 hours. This was based on the ECC visit time, so that the actual accident time could be slightly earlier than 18:00 to 20:00 hours.

For the injured body parts, 999 cases (73.7%) occurred in the head and face, followed by the hand and the wrist (134 cases [9.9%]). The hands are generally highly used body parts and are used to protect other body parts in cases of slips or falls. The high incidence rate of accidents in the hands and wrists could be related to this. In the cases of falls, many face injuries occurred because the head and face bump strongly against the floor.

Seventy-one cases (5.7%) occurred in bed. As the bed is related to falling down, slips or blunt injuries, it should be checked first to prevent accidents in the home. Of the more than 250 detailed causes of accidents, 17 were related to 50.4% of the accidents. As childhood accidents occur under certain circumstances, the identification of the common causes of such accidents can be very useful for their prevention.

Blunt injuries due to collisions with things appeared to have been caused by structures or furniture in the house, and blunt injuries due to collisions with other people appeared to have been related to sports activities. As for crushing injuries, 31 cases (59.6%) were caused by doors. Slips accounted for the largest proportion of mechanisms of the accidents [448 cases (29.4%)]. In most of the cases, however, the relevant medical charts did not contain information on the place where the slip occurred. For the accidents with identified causes, the number of slips in the bathroom was the largest [47 cases (10.5%)]. Notably, no frostbite occurred, which could have been because this study was conducted in a downtown hospital where the living environment has improved from the past and where parental supervision of children is good. As for swallowing of foreign bodies, fish bones were the major causes. Thus, parents should avoid giving fish to young children or should give them fish after removing all of the fish bones. In addition, bicycle accidents accounted for a large proportion of the accidents. Thus, children must wear protective gear when riding a bicycle.

This study was conducted on pediatric patients who visited ECCs using the outpatient medical chart. As the emergency room outpatient medical chart was not prepared for the purpose of the study, it had limitations with respect to the use of its contents as basic data, unlike had data been prepared precisely for the purpose of this study. The main limitation of this study was that some emergency room medical charts did not contain the detailed causes or mechanisms of the accidents. However, the selection of relatively more serious injuries can be the strength of this study.

It appeared that during the preparation of the ER chart, more focus was given to the outcome of the accidents than to the mechanisms of the accidents. In particular, in most of the cases of slips, the place where the slip occurred, and whether the injury was due to the slip itself or was caused secondarily by bumping into things after the slip, were not mentioned in the chart. Had the causes of the accidents been classified as primary, secondary, or tertiary when the chart was prepared, the data in the chart could have been used more systematically.

Even when the cause of an accident was described in the medical chart, its classification was not easy because the exact height of the falls was not mentioned and some injuries occurred due to various reasons, through a complex process. In such cases, one cause that was considered the direct cause of the accident was referred to as the primary cause of the accident. Although every effort was made to perform the analysis based on the given data, avoiding subjective judgment, there may have been some subjective opinions in the course of this study. To overcome these limitations, the classification criteria to be used for future studies should be used in

the preparation of the outpatient medical chart.

If a system for monitoring injuries will be established, many studies on subjects similar to the subject of this study are expected to be conducted. Previous studies had limitations in terms of the inconsistency of the classification system, which limited comparison between the studies. Also, the existing classification of the mechanisms of accidents is too broad to be used to establish practical preventative measures. In this context, a novel classification system and criteria are required. A new classification system that covers the mechanisms and detailed causes of accidents and that can be used for future studies is necessary. As ICECI is too complex to be used and unsuitable for the local situation, a classification system based on it will need amendment and complementation. Because many studies have focused on the mechanisms of accidents, future studies must focus on the relationship between the mechanisms and the causes of accidents to be useful in the prevention of accidents.

Acknowledgement

This work was supported by research grant for medical student from Yeungnam University college of medicine.

References

- 1) Hong CE. Textbook of pediatrics, 9th ed. Seoul: Daehan Printing & Publishing Co, 2008:16-36.
- 2) Statistics Korea. Ranking of causes of death by age [Internet]. Statistics Korea; 2008 [Updated 2008]. Available from: http://kosis.kr/nsportal/ups/ups_02List01.jsp?kor_id=6&pubcode=YD&ctype=F.
- 3) Sminkey L. World report on child injury prevention. *Inj Prev* 2008;14:69.
- 4) Mace SE, Gerardi MJ, Dietrich AM, Knazik SR, Mulligan-Smith D, Sweeney RL, et al. Injury prevention and control in children. *Ann Emerg Med* 2001;38:405-14.
- 5) WHO. International Classification of External Causes of Injury (ICECI) [Internet]. WHO; 1998 [Updated 2003]. Available from: <http://www.who.int/classifications/icd/adaptations/iceci/en/index.html>.
- 6) Lee HJ, Park SY, Lee YH, Do BS, Lee SB. Clinical analysis of the pediatric patients seen in the emergency medical center. *Korean J Pediatr* 2005;48:1061-7.
- 7) Park JH, Bae YS. Causes of childhood injuries observed at the emergency rooms of five hospitals in Daegu. *Korean J Prev Med* 1988;21:224-37.
- 8) Yoon OB, Kang HS. Characteristics of children admitted to a pediatric emergency department following an accident. *J Korean Clin Nurs Res* 2009;15:79-91.
- 9) Ahn BH, Chung YH, Kim YW, Kim KB. Statistical study on pediatric emergency room patients. *J Korean Pediatr Soc* 1992;35:621-9.
- 10) Shin KS. Clinical analysis on accident of children. *J Korean Pediatr Soc* 1982;25:829-38.
- 11) Park TJ. Clinical study of childhood accident. *J Korean Pediatr Soc* 1986;29:60-70.
- 12) Kim CM. Clinical study of childhood accidents from a hospital over ten years with regard to foreign body aspiration. *J Korea Pediatr Soc* 2002;45:1134-40.
- 13) Choi HJ. Statistical study on childhood accidents. *J Korean Pediatr Soc* 1981;24:837-43.
- 14) Lee MS, Eum KD, Lee EH, Lee KM, Min JY, Ha TK. A study on childhood injury in general population. *Korean J Public Health* 2001;38:41-50.
- 15) Langley JD, McLoughlin E. Injury mortality and morbidity in New Zealand. *Accid Anal Prev* 1989;21:243-54.
- 16) Kim SJ, Lee JS, Cha HD, Baek TW, Kwo TC, Kang JM. Statistical study on pediatric emergency room patients. *J Korean Pediatr Soc* 1988;31:93-9.
- 17) Park JH, Wi DH. Clinical analysis of the pediatric patients visiting emergency centre. *J Korean Soc Emerg Med* 1998;9:445-51.
- 18) Seo SS, Lee JW, Kim CY, Shin JW. Clinical and statistical studies on the emergency room patients. *J Korean Pediatr Soc* 1979;22:46-53.
- 19) Ryu JM, Yoon CH, Choi SH, Cho KG. The statistical studies on the emergency room patients under the 15 years of age. *J Korean Pediatr Soc* 1985;28:372-7.
- 20) Crawley-Coha T. Childhood injury: a status report. *J Pediatr Nurs* 2001;16:371-4.
- 21) Burt CW, Fingerhut LA. Injury visits to hospital emergency departments: United States, 1992-95. *Vital Health Stat* 13 1998;1-76.