

Review of the Shell-bearing Gastropods in the Russian Waters of the East Sea.

II. Caenogastropoda: Sorbeoconcha, Hypsogastropoda

Vladimir V. Gulbin

A.V. Zhirmunsky Institute of Marine Biology, Far East Branch of the Russian Academy of Sciences, Vladivostok 690041, Russia

ABSTRACT

Based on investigation of collections at the A.V.Zhirmunsky Institute of Marine Biology of the Far-Eastern Branch of the Russian Academy of Sciences (Vladivostok) and Zoological Institute of the Russian Academy of Sciences (Saint-Petersburg), as well as on the analysis of published data, the species composition of shell-bearing gastropod mollusks of the Russian waters of the East Sea, consisting of 331 species and subspecies, has been identified. Each species is assigned to a biogeographic grouping in the study area. Its type locality and habitat (depth and substrate) are also documented. The second part of this review includes 101 species from Sorbeoconcha and Hypsogastropoda (Caenogastropoda).

Keywords: Gastropoda, East Sea, faunal

Introduction

The fauna of shell-bearing gastropod mollusks of the Russian waters of the East Sea includes 331 species and subspecies: 306 prosobranchs, 24 opisthobranchs, and 1 pulmonate. The second part of this review includes 101 species from Sorbeoconcha and Hypsogastropoda (Caenogastropoda).

For each species a brief synonymy is given, including the original description, and reference to the most complete faunal reports on the mollusks of Korea (Min *et al.*, 2004), Japan (Higo *et al.*, 1999), and Russia (Kantor and Sysoev, 2006). Its type locality is also listed, as well as its distribution. The distribution includes its biogeographic grouping, the specific faunal zones (Fig. 1) where it was obtained in the Russian waters of the East Sea (Roman numerals), and a general statement of its abundance.

Its habitat is also listed, and includes data on depth and substrate which refer only to its distribution in




Fig. 1. The faunal zones in the Russian Waters of the East Sea.

Received April 7, 2010; Accepted June 14, 2010

Corresponding author: V.V. Gulbin

e-mail: vgulbin@imb.dvo.ru

1225-3480/24344

the Russian waters of the East Sea.

The classification of gastropod mollusks developed by Bouchet and Rocroi (2005) is used in the paper up to a superfamily rank. Also, all shell-bearing gastropod mollusks have been divided into the groups of Patellogastropoda, Vetigastropoda, Cocculiniformia, Caenogastropoda and Heterobranchia, without assigning them a taxonomic rank. Subordinate taxa within a higher rank taxon are listed, as a rule, in alphabetical order. Subgenera were not distinguished.

SYSTEMATICS

Sorbeoconcha

Superfamily Cerithioidea Fleming, 1822

Family Batillariidae Thiele, 1929

Genus **Batillaria** Benson, 1842

Type species: *Cerithium zonalis* Bruguière, 1792.

56. **Batillaria cumingii** (Crosse, 1862).

[*Lampania cumingii* Crosse, 1862: 54, pl. 1, fig. 10, 11; *Batillaria cumingii*: Higo et al., 1999: 88 Min et al., 2004: 131, fig. 225; Kantor and Sysoev, 2006: 47-48, pl. 22 K].

Type locality: Northern China.

Distribution: Asian subtropical-low-boreal species; I; common.

Habitat: Intertidal -9 m; muddy sand, muddy gravel

Family Litiopidae Gray, 1847

Genus **Diffalaba** Iredale, 1936

Type species: *Diffalaba opioniosa* Iredale, 1936.

57. **Diffalaba picta** (A. Adams, 1861).

[*Diala picta* A. Adams, 1861b: 243; *Diffalaba picta*: Higo et al., 1999: 85 Min et al., 2004: 125, fig. 203; Kantor and Sysoev, 2006: 48, pl. 22 E].

Type locality: Laoshang (Shangong Province, China).

Distribution: Asian subtropical-low-boreal species; I; common.

Habitat: Intertidal -2 m; rock and stones

Family Turritellidae Lovén, 1847

Genus **Neohaustator** Ida, 1952

Type species: *Turritella nipponica* Yokoyama, 1920.

58. **Neohaustator fortيلرatus** Sowerby 1914

[*Turritella fortيلرata* Sowerby, 1914: 36, pl. 2, fig. 12; *Neohaustator fortيلرata fortيلرata*: Higo et al., 1999: 88 Min et al., 2004: 131, fig. 227; *Neohaustator fortيلرata*: Kantor and Sysoev, 2006: 48, pl. 23 A].

Type locality: Nemuro (Hokkaido).

Distribution: Asian low-boreal species; I, II, III, IV, V, VII, VIII (empty shells only); common.

Habitat: 30-200 m; mainly sand and muddy sand

Genus **Tachyrhynchus** Mörch, 1868

Type species: *Turritella reticulata* Mighels and Adams, 1842.

59. **Tachyrhynchus erosus erosus** (Couthouy, 1838)

[*Turritella erosa* Couthouy, 1838: 103, pl. III, fig. 1; *Tachyrhynchus erosus erosus*: Higo et al., 1999: 89 Kantor and Sysoev, 2006: 49, pl. 23 C].

Type locality: Massachusetts Bay.

Distribution: Boreal-arctic species; I-VII; common.

Habitat: 10-220 m; mainly sand, muddy sand, and muddy sand with gravel

60. **Tachyrhynchus reticulatus reticulatus** (Mighels and Adams, 1842)

[*Turritella reticulata* Mighels and Adams, 1842: 50, pl. 4, fig. 19; *Tachyrhynchus reticulatus*: Higo et al., 1999: 89; *Tachyrhynchus reticulatus reticulatus*: Kantor and Sysoev, 2006: 49, Table 23 B].

Type locality: Chaleur Bay (Gulf of St. Lawrence, eastern Canada).

Distribution: Boreal-arctic subspecies; I, II, III, IV, V, VI common.

Habitat: 65-373 m; sand, muddy sand and muddy sand with gravel

Clade Littorinimorpha

Superfamily Calyptraeoidea Lamarck 1809

Family Calyptraeidae Lamarck 1809

Genus **Crepidula** Lamarck, 1799

Type species: *Patella fornicata* Linnaeus, 1758.

61. *Crepidula derjugini* Golikov and Kussakin, 1962.
[Golikov and Kussakin, 1962: 291, pl. 1, fig. 4; Kantor and Sysoev, 2006: 50-51, pl. 23 G].

Type locality: Sovetskaya Gavan Bay (Tatar Strait).
Distribution: Asian low-boreal species; I, II, III, IV, V, VI, VII; common
Habitat: Intertidal -62 m; rock and stones, sand; often on Neptunea shells.

62. *Crepidula grandis* Middendorff, 1849
[Middendorff, 1849: 101-103, pl. XI, figs. 8-10 Higo et al., 1999: 116; Kantor and Sysoev, 2006: 51 pl. 23 F].
Type locality: St. Paul Island, Bering Sea.
Distribution: Pacific boreal species I (storm drift), IV, VI, VII; single records only
Habitat: 8-500 m; rock and stones, sand, muddy sand with pebbles often on Neptunea shells.

Genus *Crepipatella* Lesso, 1830
Type species: *Patella dilatata* Linnaeus, 1803.

63. *Crepipatella lingulata* Gould, 1846
[*Crepidula lingulata* Gould, 1846: 160; *Crepipatella lingulata*: Higo et al., 1999: 116; Kantor and Sysoev, 2006: 51, pl. 23 F].
Type locality: Puget Sound, Oregon.
Distribution: Pacific boreal species II, III, IV, V; single records only
Habitat: 40-130 m; sand, muddy sand with pebbles

Superfamily Capuloidea Fleming, 1822

Family Capulidae Fleming, 1822

Genus *Ariadnaria* Habe, 1961

Type species: *Trichotropis borealis* Broderip and Sowerby, 1829.

64. *Ariadnaria borealis* (Broderip and Sowerby, 1829).
[*Trichotropis borealis* Broderip and Sowerby, 1829: 375; Higo et al., 1999: 117; *Ariadnaria borealis*: Kantor and Sysoev, 2006: 51-52, pl. 24 H].
Type locality: Melville Island, northern Canada.
Distribution: Boreal-arctic species; I, II, III, VI; rare.
Habitat: 31-246 m; sand and sand with shells

65. *Ariadnaria insignis* (Middendorff 1849)
[*Trichotropis insignis* Middendorff, 1848: 107, pl. X, figs. 7-9 Higo et al., 1999: 117; *Ariadnaria insignis*: Kantor and Sysoev, 2006: 52, pl. 24 G].

Type locality: Sitka (Bering Sea).
Distribution: Pacific boreal species I, II, III, IV, V, VI, VII, VIII; common.
Habitat: 5-100 m; stone, sand with shells

Genus *Ciliatotropis* Golikov, 1986
Type species: *Ciliatotropis striata* Golikov, 1986.

66. *Ciliatotropis japonica* Golikov, 1986.
[*Ciliatotropis japonica* Golikov, 1986a: 14, figs. 1, 7 Min et al., 2004: 153, fig. 314; Kantor and Sysoev, 2006: 53, pl. 25 B].
Type locality: south-western part of Tatar Strait.
Distribution: Asian low-boreal species; II, III; common.
Habitat: 50-130 m; stone, muddy sand with pebbles

Genus *Neophinoe* Habe, 1978
Type species: *Trichotrpis unicarinata* Sowerby, 1834.

67. *Neophinoe coronata* (Gould, 1860).
[*Trichotropis coronata* Gould, 1860: 324; *Iphinoe coronata*: Higo et al., 1999: 117; *Neophinoe coronata*: Kantor and Sysoev, 2006: 54, pl. 26 C].
Type locality: Strait of Seniavine, Arctic Ocean.
Distribution: Boreal-arctic species; III; single record only
Habitat: 205 m; muddy sand

68. *Neophinoe echinata* (Egorov and Alexeyev 1998)
[*Iphinoe echinata* Egorov and Alexeyev 1998: 15, fig. 12; *Neophinoe echinata*: Kantor and Sysoev, 2006: 54; pl. 25 F].

Type locality: Tatar Strait.
Distribution: Asian low-boreal species; VI very rare.
Habitat: 200 m.

69. *Neophinoe kroyeri* (Philippi, 1849)
[*Trichotropis kroyeri* Philippi, 1849: 175; *Iphinopsis kroyeri*: Higo et al., 1999: 118; Kantor and Sysoev, 2006: 54-55, pl. 25 F].

Review of the Shell-bearing Gastropods in the Russian Waters of the East Sea. II. Caenogastropoda:
Sorbeoconcha, Hypsogastropoda

Type locality: Spitzbergen.

Distribution: Boreal-arctic species; VIII single record only

Habitat: 110 m; sand.

70. ***Neophinoe triseriata*** Golikov 1986

[*Iphinoe triseriata* Golikov, 1986a: 18-19, figs. 4, 10;
Neophinoe triseriata: Kantor and Sysoev, 2006:
554-55, pl. 25 E].

Type locality: 4th Kurile Strait.

Distribution: Asian boreal species; VI; single record only

Habitat: 221 m; sand, mud.

Genus ***Torellia*** Loven in Jeffreys, 1867

Type species: *Torellia vestita* Jeffreys, 1867.

71. ***Torellia*** Golikov and Gulbin, 1978.

[*Torellia acuta* Golikov and Gulbin, 1978: 205-206,
figs. 16, 24, 4; Kantor and Sysoev, 2006: 55, pl. 25 F].

Type locality: Urup Island (Middle Kurile Islands).

Distribution: Asian boreal species; VII; single record only

Habitat: 31 m; stones and sand

Genus ***Trichamathina*** Habe, 1962

Type species: *Amathina nobilis* A. Adams, 1867.

72. ***Trichamathina buccinula*** Golikov in Golikov and Scarlato, 1967

[*Capulus buccinulum* Golikov and Scarlato, 1967: 43,
pl. 1, fig. 17; *Trichamatina buccinula*: Higo et al.,
1999: 118, Kantor and Sysoev, 2006: 55, pl. 26 G].

Type locality: Posjet Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I, II, IV, V;
single records only

Habitat: 35-667 m; pebbles with sandy mud

73. ***Trichamathina nobilis*** (A. Adams, 1867)

[*Amathina nobilis* A. Adams, 1867: 312, pl. 19, fig. 27;
Trichamatina nobilis: Higo et al., 1999: 118; Min et
al., 2004: 153, fig. 317; Kantor and Sysoev, 2006: 55,
pl. 26 H].

Type locality: Cape Ostry (= Notoro), South-Eastern
Sakhalin Island (Sea of Okhotsk).

Distribution: Asian boreal species; II, VII, VIII rare.

Habitat: 5-120 m; stone, pebble and muddy sand

Superfamily Littorioidea Children, 1834

Family Littorinidae Children, 1834

Subfamily Lacuninae Gray, 1857

Genus ***Epheria*** Leach in Gray, 1847

Type species: *Turbo vinctus* Montagu, 1803.

74. ***Epheria decorata*** (Adams, 1861)

[*Lacuna decorata* A. Adams, 1861b: 304; *Epheria
decorata*: Higo et al., 1999: 93; Min et al., 2004: 135,
fig. 243; Kantor and Sysoev, 2006: 60, pl. 29 D].

Type locality: Rishiri Island, Hokkaido (East Sea).

Distribution: Asian low-boreal species; I, II, IV, VI,
VII, VIII; common.

Habitat: intertidal -23 m; rock and stones

75. ***Epheria turrita*** (Adams, 1861)

[*Lacuna turrita* A. Adams, 1861b: 305; *Tamanella
turrita*: Higo et al., 1999: 93 *Temanella turrita* Min et
al., 2004: 135, fig. 242; *Epheria turrita* Kantor and
Sysoev, 2006: 61, pl. 29 G].

Type locality: Rishiri Island, Hokkaido (East Sea).

Distribution: Asian low-boreal species; I-VIII;
abundant

Habitat: intertidal -25 m; mainly rock and stones
usually in algae.

Genus ***Haloconcha*** Dall, 1886

Type species: *Lacunella refleza* Dall, 1884.

76. ***Haloconcha minor*** (Dall, 1919).

[*Haloconcha minor* Dall, 1919: 350 Higo et al., 1999:
94; Kantor and Sysoev, 2006: 61, pl. 29 I-I'].

Type locality: St. Paul Island, Pribilof Group (Bering
Sea).

Distribution: Pacific boreal species I, II-VII; common.

Habitat: intertidal -20 m; mainly rock and stones
usually in algae.

77. ***Haloconcha refleza*** (Dall, 1884)

[*Lacunella refleza* Dall, 1884: 344-345, pl. 2, figs. 1-3; *Haloconcha refleza*: Higo et al., 1999: 93-94; Kantor and Sysoev, 2006: 61-62, pl. 29 H].

Type locality: St. Paul Island, Pribilof Group (Bering Sea).

Distribution: Pacific boreal species II, VIII; single records only

Habitat: intertidal -30 m; rock.

Genus ***Stenotis*** A.Adams, 1863

Type species: *Stenotis laxata* A. Adams, 1863.

78. ***Stenotis uchidai*** (Habe, 1953)

[*Stenotis uchidai* Habe, 1953: 136, fig. 19; Higo et al., 1999: 93; Min et al., 2004: 137, fig. 247; Kantor and Sysoev, 2006: 62, pl. 30 C].

Type locality: Akkeshi Bay, Hokkaido.

Distribution: Asian low-boreal species; II, V, VII, VIII; single records only

Habitat: intertidal -25 m; rock; usually in algae and eelgrass

Subfamily Littorininae Children, 1834

Genus ***Littorina*** Ferrussac, 1821

Type species: *Turbo littoreus* Linnaeus, 1758.

79. ***Littorina brevicula*** (Philippi, 1844)

[*Turbo brevicula* Philippi, 1844: 166; *Littorina brevicula*: Higo et al., 1999: 90; Min et al., 2004: 133, fig. 233; Kantor and Sysoev, 2006: 57, pl. 27 E].

Type locality: mouth of the Yangtze River, China.

Distribution: Asian subtropical-low-boreal species; I-VIII; abundant.

Habitat: intertidal -3 m; various, mainly rock and stones

80. ***Littorina mandshurica*** Schrenck, 1867

[*Littorina mandshurica* Schrenck, 1867: 333-339, pl. XIV, figs. 14-20; Higo et al., 1999: 90 Kantor and Sysoev, 2006: 58, pl. 27 H].

Type locality: Bay of Chihachev, Tatar Strait.

Distribution: Asian low-boreal species; I-VIII; abundant.

Habitat: intertidal -1 m; various, mainly rock and stones

81. ***Littorina sitkana*** Philippi, 1846

[*Littorina sitkana* Philippi, 1846: 140; *Littorina sitkana*: Higo et al., 1999: 90-91; Min et al., 2004: 133, fig. 234; Kantor and Sysoev, 2006: 59, pl. 28 A-B].

Type locality: Sitka, Alaska.

Distribution: Pacific boreal species I-VIII; abundant.

Habitat: intertidal -1.5 m; various, mainly rock and stones

82. ***Littorina squalida*** Broderip and Sowerby, 1829

[*Littorina squalida* Broderip and Sowerby, 1829: 370; Higo et al., 1999: 90; Kantor and Sysoev, 2006: 59, pl. 28 I].

Type locality: Icy Cape (northwestern Alaska).

Distribution: Pacific boreal species I-VIII; abundant.

Habitat: intertidal -16 m; various, mainly rock and stones

Superfamily Naticoidea Guilding, 1834

Family Naticidae Guilding, 1834

Genus ***Bulbus*** Brown in Smith, 1839

Type species: *Bulbus smithii* Brown in Smith, 1839.

83. ***Bulbus normalis*** (Middendorff, 1851)

[*Natica aperta* forma *normalis* Middendorff, 1851: 206-208, pl. 11, figs. 1-3; *Bulbus flavus elongates*: Higo et al., 1999: 143; *Bulbus flavus flavus*: Min et al., 2004: 171, fig. 385; *Bulbus normalis*: Kantor and Sysoev, 2006: 63, pl. 30 J].

Type locality: Shantar Islands (Sea of Okhotsk).

Distribution: Asian boreal species; I, II, VI, VIII; common.

Habitat: 5-60 m; muddy sand with pebbles

84. ***Bulbus tenuiculus*** (Sowerby, 1915)

[*Natica (Polynices) tenuiculus* Sowerby, 1915: 166, pl. 10, fig. 3; *Bulbus flavus flavus*: Higo et al., 1999: 142; *Bulbus tenuiculus*: Kantor and Sysoev, 2006: 64, pl. 30 I].

Type locality: Nemuro Peninsula (Sea of Okhotsk, Japan).

Distribution: Asian low-boreal species; I, IV, VI; VIII; rare.

Habitat: 40-130 m; muddy sand with pebbles

Genus ***Cryptonatica*** Dall, 1892

Type species: *Natica (Cryptonatica) floridiana* Dall, 1892.

85. ***Cryptonatica affinis*** (Gmelin, 1791)
[*Nerita affinis* Gmelin, 1791: 3675; *Cryptonatica clausa*: Higo et al., 1999: 148; Min et al., 2004: 177, fig. 411; *Cryptonatica affinis*: Kantor and Sysoev, 2006: 64, pl. 31 A].

Type locality: Iceland.

Distribution: Boreal-arctic species; I-VIII; common.

Habitat: 47-900 m; mainly sand and muddy sand

86. ***Cryptonatica aleutica*** (Dall, 1919)
[*Natica aleutica* Dall, 1919: 352; *Cryptonatica russa*: Higo et al., 1999: 149; Min et al., 2004: 177, fig. 413; *Cryptonatica aleutica*: Kantor and Sysoev, 2006: 64, pl. 31 B].

Type locality: Unalaska, Aleutian Islands.

Distribution: Pacific boreal species I-VII; common.

Habitat: 10-400 m; mainly sand and muddy sand

87. ***Cryptonatica hirasei*** (Pilsbry, 1905)
[*Natica hirasei* Pilsbry, 1905: 105; *Cryptonatica hirasei*: Higo et al., 1999: 149; Min et al., 2004: 179, fig. 419; Kantor and Sysoev, 2006: 64-65, pl. 31 D].
Type locality: Akkeshi and Kushiro, Hokkaido.
Distribution: Asian low-boreal species; I-VIII; common.
Habitat: intertidal -70 m; mainly sand and muddy sand, rock (rare)

88. ***Cryptonatica janthostoma*** (Deshayes, 1839)
[*Natica janthostoma* Deshayes, 1839: 361 *Cryptonatica janthostoma*: Higo et al., 1999: 149; Min et al., 2004: 179, fig. 417; Kantor and Sysoev, 2006: 65, pl. 31 F; *Cryptonatica janthostomoides*: Higo et al., 1999: 149; Min et al., 2004: 179, fig. 418].
Type locality: Kamchatka Peninsula.
Distribution: Asian boreal species; I-VIII; common.
Habitat: intertidal -120 m; mainly sand and muddy sand

89. ***Cryptonatica wakkanaiensis*** Habe and Ito, 1976

[*Cryptonatica wakkanaiensis* Habe and Ito, 1976: 8, fig. 3; Higo et al., 1999: 148; Min et al., 2004: 177, fig. 412; Kantor and Sysoev, 2006: 65, pl. 31 G].

Type locality: Wakkanai Bay, northernmost Hokkaido.

Distribution: Asian low-boreal species; I, II, IV, V, VIII; common.

Habitat: intertidal -550 m; mainly muddy sand with gravel

90. ***Cryptonatica zenryumaruae*** Habe and Ito, 1976

[*Cryptonatica zenryumaruae* Habe and Ito, 1976, pp. 79-82, fig. 4; Higo et al., 1999: 149 Min et al., 2004: 177-178, fig. 416; Kantor and Sysoev, 2006: 65, pl. 31 C].

Type locality: Cape Soya, northernmost Hokkaido.

Distribution: Asian low-boreal species; VIII; rare.

Habitat: 18-115 m; shell-sand-gravel-shells with sand and gravel

Genus ***Glossaulax*** Pilsbry, 1929

Type species: *Neverita reclusiana* Deshayes, 1841.

91. ***Glossaulax didyma didyma*** (Röding, 1789)

[*Albula didyma* Röding, 1789: 20; *Glossaulax didyma didyma*: Higo et al., 1999: 144; Min et al., 2004: 173, fig. 392; *Glossaulax didyma*: Kantor and Sysoev, 2006: 66, pl. 32 F].

Type locality: none designated.

Distribution: Asian subtropical species; I; one record only (empty shell)

Habitat: beach drift only

Genus ***Lunatia*** Gray, 1847

Type species: *Natica ampullarius* Lamarck, 1822.

92. ***Lunatia pallida*** (Broderip and Sowerby, 1829)

[*Natica pallida* Broderip and Sowerby, 1829: 372; *Lunatia pallida*: Higo et al., 1999: 143; Kantor and Sysoev, 2006: 66, pl. 32 E].

Type locality: Icy Cape (northwestern Alaska).

Distribution: Boreal-arctic species; I-VIII; common.

Habitat: 5-400 m mainly muddy sand

93. **Lunatia pila** (Pilsbry, 1911)

[*Polinices pila* Pilsbry, 1911: 32; *Lunatia pila pila*: Higo et al., 1999: 143; *Lunatia pila*: Kantor and Sysoev, 2006: 66, pl. 32 F].

Type locality: Kushiro and Akkeshi, Hokkaido.

Distribution: Asian low-boreal species; I, II, IV, V, VIII; common.

Habitat: 8–80 m; mainly muddy sand

Genus **Pseudopolinices** Golikov et Sirenko, 1983

Type species: *Natica nana* Møller, 1842.

94. **Pseudopolinices nanus** (Møller, 1842)

[*Natica nana* Møller, 1842: 80 *Pseudopolinices nana*: Higo et al., 1999: 143; Kantor and Sysoev, 2006: 67, pl. 32 C-C'].

Type locality: Greenland.

Distribution: Boreal-arctic species; I, VIII uncommon (dead shells only)

Habitat: 59–25 m sand

Superfamily Rissooidea Gray, 1847

Family Assimineidae H. et A. Adams, 1856 in
1853–1858

Genus **Assiminea** Fleming, 1828

Type species: *Assiminea grayana* Fleming, 1828.

95. **Assiminea lutea** A. Adams, 1861

[*Assiminea lutea* A. Adams, 1861b: 307; Min et al., 2004: 139, fig. 259; Kantor and Sysoev, 2006: 78, pl. 38 G; *Assiminea septentrionalis* Habe, 1942: 39, pl. 1 fig. 2, pl. 4 fig. 13; Higo et al., 1999: 97; Min et al., 2004: 139, fig. 257;].

Type locality: Baihe, Hebei, northern China.

Distribution: Asian subtropical-low-boreal species; I, VI; estuaries and lagoons common.

Habitat: intertidal ~2 m; muddy sand, sand, mud, or stones, usually in sea grass

96. **Assiminea possietica** Golikov and Kussakin in Golikov et Scarlato, 1967

[*Assiminea possietica* Golikov and Scarlato, 1967: 36–37, fig. 28; Kantor and Sysoev, 2006: 78–79, pl. 38 H].

Type locality: Posjet Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I; estuaries and lagoons common.

Habitat: intertidal; sand, muddy sand

Family Barleidae Gray, 1853

Genus **Barleeia** Clark, 1855

Type species: *Turbo ruber* Adams, 1797.

97. **Barleeia angustata** (Pilsbry, 1901)

[*Assiminea angustata* Pilsbry, 1901: 396; *Barleeia angustata*: Higo et al., 1999: 105; Min et al., 2004: 145, fig. 281; *Ansola angustata*: Kantor and Sysoev, 2006: 79, pl. 38 J].

Type locality: Rishiri Island, northwestern Hokkaido.

Distribution: Asian subtropical-low-boreal species; I, II, VII, VIII; common.

Habitat: intertidal ~40 m; rock and stones mainly on algal fronds

98. **Barleeia angulosa** (Golikov and Kussakin, 1978)

[*Ansola angulosa* Golikov and Kussakin, 1978: 127, fig. 85; Kantor and Sysoev, 2006: 79, pl. 38 K].

Type locality: Putjatin Island (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I; common.

Habitat: intertidal ~1 m rock and stones

99. **Barleeia labiosa** (Golikov and Kussakin, 1978)

[*Ansola labiosa* Golikov and Kussakin, 1978: 127, fig. 85; Kantor and Sysoev, 2006: 79, pl. 38 I].

Type locality: Peter the Great Bay, East Sea.

Distribution: Asian low-boreal species; I; common.

Habitat: intertidal; rock and stones on *Sargassum* sp.

Family Caecidae Gray, 1847

Genus **Brochina** Gray, 1857

Type species: *Dentalium glaber* Montagu, 1803.

100. **Brochina derjugini** Golikov in Golikov and Scarlato, 1967

[*Brochina derjugini* Golikov and Scarlato, 1967: 38–39, fig. 29; Kantor and Sysoev, 2006: 80, pl. 39 A].

Type locality: Patrokl Bay (Peter the Great Bay, East Sea).

Review of the Shell-bearing Gastropods in the Russian Waters of the East Sea. II. Caenogastropoda:
Sorbeoconcha, Hypsogastropoda

Distribution: Asian low-boreal species; I, II, VIII;
common.

Habitat: intertidal -70 m; rock and stones, sand (rare)

Genus **Fartulum** Carpenter, 1858

Type species: *Caecum leave* C. B. Adams, 1852.

101. ***Fartulum bucerium*** Golikov in Golikov and Scarlato, 1967

[*Fartulum bucerium* Golikov and Scarlato, 1967: 39 fig. 30 a, b; Kantor and Sysoev, 2006: 80, pl. 39 B].
Type locality: Patrokl Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I; common.

Habitat: Intertidal -9 m; muddy sand, rock and stones (rare) usually in sea grass

Family Falsicingulidae Slavoschevskaya, 1975

Genus **Falsicingula** Habe, 1958

Type species: *Cingula kurilensis* Pilsbry, 1905.

102. ***Falsicingula athera*** Bartsch in Golikov and Scarlato, 1967

[*Falsicingula athera* Golikov and Scarlato, 1967: 28-29, fig. 19; Kantor and Sysoev, 2006: 82, pl. 39 H-H'].

Type locality: Yuzhno-Morskoy (= Tafuin) Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I, II, IV, VII, VIII; common.

Habitat: intertidal -45 m; rock and stones, muddy sand (rare)

103. ***Falsicingula kurilensis*** (Pilsbry, 1905)

[*Cingula kurilensis* Pilsbry, 1905: 106, pl. 4, fig. 31; *Falsicingula kurilensis*: Higo et al., 1999: 96 Kantor and Sysoev, 2006: 82, pl. 39 M].

Type locality: Kunashir Island (South Kurile Islands).

Distribution: Asian low-boreal species; II, V, VII, VIII; common.

Habitat: intertidal -5 m; rock and stones usually in algae

104. ***Falsicingula mundana*** (Yokoyama, 1926)

[*Rissoa (Cingula) mundana* Yokoyama, 1926: 276, pl.

33, fig. 11; *Cingula mundana*: Higo et al., 1999: 98; *Falsicingula mundana*: Kantor and Sysoev, 2006: 82, pl. 39 J].

Type locality: Sado Island, East Sea (fossil).

Distribution: Asian low-boreal species; I-VIII; abundant

Habitat: intertidal -20 m; rock and stones usually in algae

Family Iravadiidae Thiele, 1928

Genus **Iravadia** Blanford, 1867

Type species: *Iravadia ornata* Blanford, 1867.

105. ***Iravadia elegantula*** (A.Adams, 1861).

[*Onoba elegantula* A. Adams, 1861a: 136; *Iravadia (Fluviocingula) elegantula*: Higo et al., 1999: 96; *Fluviocingula elegantula*: Kantor and Sysoev, 2006: 112-113, pl. 51 C].

Type locality: Gulf of Pe-chili (Gulf of Chihli, China).

Distribution: Asian subtropical-low-boreal species; I, V; uncommon (brackish water lagoons, channels and river mouths)

Habitat: intertidal -0.3 m; muddy sand

Genus **Hyala** H. et A. Adams

Type species: *Turbo vitrea* Montagu, 1803.

106. ***Hyala adamsi*** Golikov et Kussakin in Golikov and Scarlato, 1971.

[*Hyala adamsi* Golikov and Scarlato, 1971: 193, fig. 5; Kantor and Sysoev, 2006: 112, pl. 51 A-B].

Type locality: Patrokl Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I rare.

Habitat: intertidal -2 m; muddy sand, epiphytes of *Zostera* sp.

Family Pomatiopsidae Stimpson, 1865

Genus **Cecina** A. Adams, 1861

Type species: *Cecina manchurica* A. Adams, 1861.

107. ***Cecina manchurica*** A. Adams, 1861.

[*Cecina manchurica* A. Adams, 1861b: 308; Higo et al., 1999: 96 (in part); Kantor and Sysoev, 2006: 115, pl. 52 D].

Type locality: Olga and Vladimir Bay (Tatar Strait, East Sea)

Distribution: Asian boreal species; I; uncommon.

Habitat: supratidal and intertidal; sand and muddy sand, in eelgrass and algae

108. ***Cecina scarlatoi*** Prosovora, 1996

[*Cecina scarlatoi* Prosovora, 1996: 661, fig. 1 (e); 3 (v); Kantor and Sysoev, 2006: 116, pl. 52 A].

Type locality: Posjet Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I; uncommon

Habitat: supratidal and intertidal; sand and muddy sand, in eelgrass and algae

109. ***Cecina tatarica*** (Schrenck, 1867)

[*Truncatella tatarica* Schrenck, 1867: 310-312, pl. XIV, figs. 10-13; *Cecina manchurica*: Higo et al., 1999: 96 (in part); Kantor and Sysoev, 2006: 116, pl. 52 G].

Type locality: Chihachev Bay, Tatar Strait.

Distribution: Asian low-boreal species; I; II; IV; uncommon.

Habitat: supratidal and intertidal; sand and muddy sand, in eelgrass and algae

Family Rissoidae Gray, 1847

Subfamily Rissoinae Gray, 1847

Genus ***Alvania*** Risso, 1826

Type species: *Alvania europea* Risso, 1826.

110. ***Alvania ferruginea*** A. Adams, 1861

[*Alvania ferruginea* A. Adams, 1861a: 138; Higo et al., 1999: 99; Kantor and Sysoev, 2006: 67, pl. 33 B].

Type locality: Hakodate Bay, Sangar Strait.

Distribution: Asian low-boreal species; I rare.

Habitat: 67-101 m; sand and muddy sand

111. ***Alvania reticulatum*** (Golikov, 1986)

[*Punctulum reticulatum* Golikov, 1986b: 88-89, fig. 4; Kantor and Sysoev, 2006: 75, pl. 36 J]

Type locality: Rynda Bay (Southern Primorye, East Sea).

Distribution: Asian low-boreal species; I, II; uncommon.

Habitat: 240-880 m; sandy mud

Genus ***Lucidestea*** Laseron, 1956

Type species: *Lucidestea vitrea* Laseron, 1956.

112. ***Lucidestea ornata*** Golikov and Kussakin in Golikov and Scarlato, 1967

[*Falsisetia ornata* Golikov and Scarlato, 1967: 35, fig. 26; *Lucidestea ornata*: Higo et al., 1999: 100; Kantor and Sysoev, 2006: 79, pl. 38 L-M].

Type locality: Posjet Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I; uncommon.

Habitat: intertidal -4 m; rock and stones in algae and eelgrass

Genus ***Mohrensternia*** Stoliczka, 1868

Type species: *Rissoa angulata* Eichwald, 1864.

113. ***Mohrensternia derjugini*** Golikov and Licharev in Golikov and Scarlato, 1985

[*Mohrensternia derjugini* Golikov and Scarlato, 1985: 387, fig. 2; Kantor and Sysoev, 2006: 70, pl. 34 B].

Type locality: Svobodny Cape, southeastern Sakhalin (Sea of Okhotsk)

Distribution: Asian low-boreal species; I rare.

Habitat: 100-190 m; muddy sand

Genus ***Onoba*** H. et A. Adams, 1852

Type species: *Turbo striatus* J. Adams, 1852.

114. ***Onoba auriwillii axicostata*** Golikov and Gulbin, 1978.

[*Onoba auriwillii axicostata* Golikov and Gulbin, 1978: 194-195, figs. 2, 9, 10 Higo et al., 1999: 100 Kantor and Sysoev, 2006: 71-72, pl. 34 I].

Type locality: South Kurile Islands.

Distribution: Asian low-boreal species; I, IV, VII, VIII; uncommon.

Habitat: 12-70 m; rock, stone and sand with shells

115. ***Onoba castanella*** (Dall, 1887)

[*Alvania castanella* Dall, 1887: 307-308, pl. 3, fig. 5

Onoba castanella: Higo et al., 1999: 100; *Alvania michelsi*: Kantor and Sysoev, 2006: 67-68, pl. 33 A].

Type locality: none designated.

Review of the Shell-bearing Gastropods in the Russian Waters of the East Sea. II. Caenogastropoda:
Sorbeoconcha, Hypsogastropoda

Distribution: Pacific boreal species I, III, V, VI, VIII; uncommon.

Habitat: 15–74 m; rock and stones, and pebbles and shells

116. ***Onoba compsa*** Barch in Golikov, Gulbin, and Sirenko, 1987.

[*Onoba compsa* Golikov, Gulbin, and Sirenko, 1987: 33–34, pl. III, fig. 2; Kantor and Sysoev, 2006: 72, pl. 35 I].

Type locality: Sovetskaya Gavan Bay, Tatar Strait

Distribution: Asian low-boreal species; II, III, IV, VIII; uncommon.

Habitat: 6–70 m; sand with shells

117. ***Onoba laticingulata*** Golikov and Kussakin, 1978.

[*Onoba laticingulata* Golikov and Kussakin, 1978: 115–116, fig. 75; Kantor and Sysoev, 2006: 73, pl. 35 F].

Type locality: Paramushir Island (North Kurile Islands).

Distribution: Pacific high-boreal species (mainly), extending to low-boreal waters IV, V rare.

Habitat: 5–9.5 m; stones and rock

118. ***Onoba tenuistriata*** Golikov, Gulbin, and Sirenko, 1987.

[*Onoba tenuistriata* Golikov, Gulbin, and Sirenko, 1987: 34, pl. III, fig. 3; Kantor and Sysoev, 2006: 74, pl. 36 E].

Type locality: Moneron Island (East Sea).

Distribution: Asian low-boreal species; VIII rare.

Habitat: 50–70 m; sand with shells

Genus ***Pusilina*** Monterosato, 1884

Type species: *Rissoa pusilla* Philippi, 1836.

119. ***Pusilina plicosa*** (Smith, 1875).

[*Hydrobia plicosa* Smith, 1875: 103–104; *Alvania plicosa*: Higo et al., 1999: 98; Min et al., 2004: 143, fig. 271; *Pusilina plicosa*: Kantor and Sysoev, 2006: 76, pl. 37 A].

Type locality: Aomori Bay (northeastern Honshu).

Distribution: Asian low-boreal species; I–VIII; abundant

Habitat: intertidal –32 m; rock and stones usually in algae

Genus ***Setia*** H. et A. Adams, 1852

Type species: *Rissoa pulcherrima* Jeffreys, 1848.

120. ***Setia candida*** A. Adams, 1861.

[*Setia candida* A. Adams, 1861a: 136; Kantor and Sysoev, 2006: 78, pl. 38 E].

Type locality: Sado Island (East Sea).

Distribution: Asian low-boreal species; I, III, VI rare.

Habitat: 11–33 m; sand and muddy sand

Family Stenothyridae Tryon, 1866

Genus ***Stenothyra*** Benson, 1856

Type species: *Nematura deltae* Benson, 1836.

121. ***Stenothyra hokkaidoensis*** Kuroda, 1962.

[*Stenothyra japonica hokkaidonis* Kuroda, 1962: 61, pl. 4, fig. 3; Higo et al., 1999: 108; Kantor and Sysoev, 2006: 116, pl. 51 M].

Type locality: Teshio (Hokkaido, East Sea).

Distribution: Asian low-boreal species; I, VI; uncommon (brackish water lagoon-estuarine species)

Habitat: 0–1 m; sand and muddy sand

122. ***Stenothyra recondite*** Lindholm, 1929

[*Stenothyra recondita* Lindholm, 1929: 305; Kantor and Sysoev, 2006: 116–117, pl. 51 L; *Stenothyra japonica*: Higo et al., 1999: 108].

Type locality: Estuary of Tumannaja River (southern Posjet Bay, Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I, VI; uncommon (brackish water lagoon-estuarine species)

Habitat: 0–1 m; sand and muddy sand

Family Truncatellidae Gray, 1840

Genus ***Teinostoma*** H. et A. Adams, 1853

Type species: *Teinostoma politum* A. Adams (SD Cossmann, 1918).

123. ***Teinostoma atomaria*** (A. Adams, 1861)

[*Ethalia atomaria* A. Adams, 1861b: 305–306;

Teinostoma atomaria: Kantor and Sysoev, 2006: 117, pl. 52 N].

Type locality: Shan-Tung (China).

Distribution: Asian subtropical-low-boreal species; I, V, VII rare.

Habitat: 0–33 m; sand and muddy sand

Superfamily Tonnaidea Suter, 1913 (1825)

Family Ranellidae Gray, 1854

Genus **Fusitriton** Cossmann, 1903

Type species: *Triton cancellatus* Lamarck, 1816.

124. *Fusitriton oregonensis* (Redfield, 1846)

[*Triton oregonense* Redfield, 1846: 165–166, pl. 11 fig.

2; *Fusitriton oregonensis*: Higo et al., 1999: 155; Min et al., 2004: 183, fig. 431; Kantor and Sysoev, 2006: 119, pl. 53 B; *Fusitriton galea*: Higo et al., 1999: 155; Min et al., 2004: 183, fig. 432].

Type locality: Straits of St.Juan de Fuca, Oregon (northwestern America).

Distribution: Pacific boreal species II, III, VI, VIII; uncommon.

Habitat: 15–120 m; rock and stones, sand with pebbles, and muddy sand

Superfamily Velutinoidea Gray, 1842

Family Velutinidae Gray, 1842

Subfamily Capulacmaeinae Golikov et Gulbin, 1990

Genus **Piliscus** Lovén, 1859

Type species: *Capulus radiatus*, Sars, 1851.

125. *Piliscus commodus* (Middendorff, 1851)

[*Pilidium commodum* Middendorff, 1851: 214, pl. 17, figs.4–11; *Capulacmaea commoda*: Higo et al., 1999: 142 (in part); *Piliscus commoda*: Kantor and Sysoev, 2006: 125, pl.57 E (in part)].

Type locality: Tugur Bay (southern coast of the Sea of Okhotsk).

Distribution: Pacific boreal species, mainly Asian species extending to Arctic waters I, VI, VII, VIII rare.

Habitat: 36–60 m; muddy sand, stones.

Note. Many malacologists combine this species with *Piliscus radiatus* Sars, 1851. However, the *Piliscus commodus* shell consists of 2 whorls. Its embryonic

shell is small and forms a complete whorl not pressed against the posterior slope of the shell. The *Piliscus radiatus* shell consists of 1.5 whorls, and its embryonic shell is large, forming a curved horn pressed against the posterior slope (Gulbin and Golikov, 1997).

Subfamily Velutininae Gray, 1840

Genus **Cartilagovelutina** Golikov et Gulbin, 1990

Type species: *Velutina beringensis* Derjugin, 1950.

126. *Cartilagovelutina chondrina* (Bartsch in Derjugin, 1950)

[*Velutina chondrina* Derjugin, 1950: 16–17, fig. 11 pl. 2, fig. 11 pl. 5, fig. 5; *Cartilagovelutina chondrina*: Kantor and Sysoev, 2006: 120, pl. 53 E].

Type locality: Peter the Great Bay (East Sea).

Distribution: Asian boreal species; I, II, VII rare.

Habitat: 63–150 m; muddy sand

127. *Cartilagovelutina cristata* (Derjugin, 1950)

[*Velutina cristata* Derjugin, 1950: 16–17, fig. 11 pl. 2, fig. 11 pl. 5, fig. 5; *Cartilagovelutina cristata*: Kantor and Sysoev, 2006: 120, pl. 53 D].

Type locality: Tauiskaya Inlet (Sea of Okhotsk).

Distribution: Asian boreal species (mainly Sea of Okhotsk) VII; one record only

Habitat: depth unknown, muddy sand

Genus **Ciliatovelutina** Golikov et Gulbin, 1990

Type species: *Velutina lanigera* Møller, 1842.

128. *Ciliatovelutina capillata* (Derjugin, 1950)

[*Velutina capillata* Derjugin, 1950: 14–15, fig. 9 pl. 1, fig. 9 pl. 4, fig. 5; *Ciliatovelutina capillata*: Kantor and Sysoev, 2006: 120–121, pl. 54 C].

Type locality: Sakhalin, Sea of Okhotsk.

Distribution: Asian boreal species (mainly Sea of Okhotsk) VII; one record only

Habitat: 15 m rock

129. *Ciliatovelutina nana* (Bartsch in Derjugin, 1950)

[*Velutina nana* Derjugin, 1950: 12–13, fig. 6 pl. 1, fig.

6 pl. 4, fig. 6; *Ciliatovelutina nana*: Kantor and Sysoev, 2006: 121, pl. 54 B].

Type locality: East Sea.

Distribution: Asian boreal species; I, II, III, VI, VIII; uncommon.

Habitat: 1-349 m; sand with pebbles and muddy sand, rock and stones (rare)

Genus **Limneria** H. et A. Adams, 1851

Type species: *Limneria caspiensis* H. and A. Adams, 1851.

130. **Limneria prolongata** (Carpenter, 1864)

[*Velutina prolongata* Carpenter, 1864: 628, 661
Limneria conica Higo et al., 1999: 142; Min et al., 2004: 169, fig. 383; *Limneria bulla*: Higo et al., 1999: 142; Min et al., 2004: 171, fig. 384; *Limneria prolongata*: Kantor and Sysoev, 2006: 121-122, pl. 54 F-F].

Type locality: Neach Bay, Vancouver Island.

Distribution: Pacific boreal species I, II, III, IV, VI, VII, VIII; common.

Habitat: 2-94 m; muddy sand, pebbles, rock

131. **Limneria undata bifasciata** (Derjugin, 1950)

[*Velutina undata bifasciata* Derjugin, 1950: 10-11, fig. 3 pl. 1, fig. 3-3a pl. 2, fig. 3; Kantor and Sysoev, 2006: 122, pl. 54 D].

Type locality: Peter the Great Bay, East Sea.

Distribution: Asian low-boreal subspecies of a boreal-arctic species; I, II rare.

Habitat: 33-355 m; muddy sand with pebbles

Genus **Marsenina** Gray, 1850

Type species: *Lamellaria prodita* Lovén, 1850 (= *Oxinoe glabra* Couthouy, 1838).

132. **Marsenina rhombica** (Dall, 1871)

[*Lamellaria rhombica* Dall, 1871: 122, pl. 15, figs. 4, 5; *Marsenina rhombica*: Higo et al., 1999: 142; Kantor and Sysoev, 2006: 123, pl. 55 C].

Type locality: Monterey, California.

Distribution: Pacific boreal species I, II rare.

Habitat: 2-5 m rock

133. **Marsenina uchidai** (Habe, 1958)

[*Lamellaria uchidai* Habe, 1958: 14-15, pl. 1, fig. 11; *Marsenina uchidai*: Higo et al., 1999: 142; Kantor and Sysoev, 2006: 123, pl. 55 D].

Type locality: Akkeshi Bay, Hokkaido

Distribution: Asian boreal species; I, II, IV, V, VI, VII, VIII; common

Habitat: 5-60 m; sand with pebbles and shells, stones

Genus **Velutella** Gray, 1847

Type species: *Bulla flexilis* Montagu, 1808 (= *Helix plicatilis* Müller, 1776).

134. **Velutella plicatilis** (Müller, 1776)

[*Helix plicatilis* Müller 1776: 242; *Velutina cryptospira* Middendorff, 1851: 216, pl. 25, figs. 8-10; Kantor and Sysoev, 2006: 124, pl. 57 A; *Velutina* (*Velutella*) *plicatilis* *plicatilis*: Higo et al., 1999: 141-142; Min et al., 2004: 169, fig. 380; *Velutina* (*Velutella*) *plicatilis* *cryptospira*: Higo et al., 1999: 142 (in part, non *V. coriacea* Pallas, 1847); Min et al., 2004: 169, fig. 381; *Velutella plicatilis*: Kantor and Sysoev, 2006: 123, pl. 55 F].

Type locality: Denmark and Norway.

Distribution: Boreal-arctic species; I, II, VII, VIII rare.

Habitat: 15-210 m; rock and stones

Genus **Velutina** Fleming, 1821

Type species: *Bulla velutina* Müller, 1776.

135. **Velutina bartschi** Derjugin, 1950

[*Velutina bartschi* Derjugin, 1950: 18-19, fig. 14 pl. 2, fig. 14 pl. 5, fig. 14; Kantor and Sysoev, 2006: 123, pl. 56 E]

Type locality: Medny Island (Commander Islands).

Distribution: Asian boreal species; VII; one record only

Habitat: 5 m; stones

136. **Velutina coriacea** Pallas, 1788

[*Helix coriacea* Pallas, 1788: 234, pl. VII, figs. 31, 32; *Velutina plicatilis cryptospira*: Higo et al., 1999: 142 (in part); *Velutina coriacea coriacea*: Kantor and Sysoev, 2006: 123-124, pl. 57 B]

Type locality: Kamchatka, Sea of Okhotsk.
Distribution: Asian boreal species I, III, VII, VIII rare.
Habitat: 7–20 m; rock and stones, and muddy sand

137. *Velutina pulchella* Derjugin, 1950
[*Velutina pulchella* Derjugin, 1950: 17, fig. 12 pl. 2, fig. 12 pl. 5, fig. 12; Kantor and Sysoev, 2006: 124, pl. 56 B].
Type locality: Laperus Strait (southern Sakhalin).
Distribution: Asian boreal species; I, II, VII, VIII rare.
Habitat: 5–110 m; rock and stones

138. *Velutina tarasovi* Derjugin, 1950
[*Velutina tarasovi* Derjugin, 1950: 15–16, fig. 10 pl. 2, fig. 10 pl. 4, fig. 10; Kantor and Sysoev, 2006: 125, pl. 56 A].
Type locality: Valentin Bay (southern Primorye, East Sea).
Distribution: Asian low-boreal species; II, III rare.
Habitat: 17–30 m; rock and stones

139. *Velutina velutina* (Müller, 1776)
[*Bulla velutina* Müller, 1776: 242; *Velutina laevigata*: Kantor and Sysoev, 2006: 124, pl. 56 D].
Type locality: Denmark and Norway.
Distribution: Boreal-arctic species; VII, one record only
Habitat: 15 m; rock

Subfamily Onchidiopsinae Golikov et Gulbin, 1990
Genus *Onchidiopsis* Berg, 1853

Type species: *Onchidiopsis groenlandica* Bergh, 1853.
140. *Onchidiopsis groenlandica* (Bergh, 1853)
[*Onchidiopsis groenlandica* Bergh, 1853: 106, pl. 2; Kantor and Sysoev, 2006: 127, pl. 58 J; *Onchidiopsis glacialis*: Higo *et al.*, 1999: 142].
Type locality: Greenland.
Distribution: Boreal-arctic species; VIII, one record only
Habitat: unknown

141. *Onchidiopsis gurjanovae* Derjugin, 1937

[*Onchidiopsis gurjanovae* Derjugin, 1937: 13–14, fig. 6 pl. 3, fig. 5 pl. 4, fig. 5; Kantor and Sysoev, 2006: 127, pl. 58 F].

Type locality: Rynda Bay (southern Primorye, East Sea)
Distribution: Asian boreal species; II, III rare.
Habitat: 86–187 m; sand with pebbles

142. *Onchidiopsis maculata* Derjugin, 1937
[*Onchidiopsis maculata* Derjugin, 1937: 14–15, figs. 7, 8 pl. 1, fig. 1 pl. 3, fig. 6 pl. 4, fig. 6 pl. 5, fig. 6; Kantor and Sysoev, 2005: 115: 127, pl. 58 H].
Type locality: Peter the Great Bay, East Sea.
Distribution: Asian boreal species; I, II, IV rare.
Habitat: 5–80 m; muddy sand with stones

143. *Onchidiopsis nadinae* Derjugin, 1937
[*Onchidiopsis nadinae* Derjugin, 1937: 17–18, figs. 10, 11 pl. 3, fig. 8 pl. 4, fig. 8 pl. 5, fig. 8; Kantor and Sysoev, 2006: 127, pl. 58 I].
Type locality: western Kamchatka, Sea of Okhotsk
Distribution: Asian boreal species VIII; one record only
Habitat: 80 m shells

144. *Onchidiopsis variegata* Derjugin, 1937
[*Onchidiopsis variegata* Derjugin, 1937: 15–16, fig. 9 pl. 2, fig. 1 pl. 3, fig. 7 pl. 4, fig. 7 pl. 5, fig. 7; Kantor and Sysoev, 2006: 128, pl. 58 C].
Type locality: Peter the Great Bay, East Sea.
Distribution: Asian boreal species I, II, VII rare.
Habitat: 110–630 m; muddy sand with stones

Group Ptenoglossa
Superfamily Epitoniidea Berry, 1910 (1812)

Family Epitoniidae Berry, 1910 (1812)
Genus *Acirsa* Mörch, 1857
Type species: *Turritella costulata* Mighels and Adams, 1842.

145. *Acirsa ochotensis* (Middendorff, 1848)
[*Scalaria ochotensis* Middendorff, 1848: 1; *Acirsa ochotensis*: Higo *et al.*, 1999: 174; Kantor and Sysoev, 2006: 128, pl. 59 A–B]
Type locality: Sea of Okhotsk (southern region)

Review of the Shell-bearing Gastropods in the Russian Waters of the East Sea. II. Caenogastropoda:
Sorbeoconcha, Hypsogastropoda

Distribution: Asian boreal species V; one record only
Habitat: 10 m; rock.

Genus **Boreoscala** Kobelt, 1902

Type species: *Scalaria greenlandica* Perry, 1811.

146. ***Boreoscala greenlandica*** (Perry, 1811)
[*Scalaria greenlandica* Perry, 1811: pl. 28, fig. 8;
Boreoscala greenlandica: Higo et al., 1999: 179; Min et al., 2004: 197, fig. 489; Kantor and Sysoev, 2006: 128, pl. 59 E].

Type locality: Greenland.

Distribution: Boreal-arctic species; I; one record only
Habitat: 62 m; muddy sand

147. ***Boreoscala rarecostulata*** Golikov in Golikov and Scarlato, 1967

[*Boreoscala rarecostulata* Golikov and Scarlato, 1967: 48, fig. 36; Kantor and Sysoev, 2006: 129, Pl. 59 D].
Type locality: Posjet Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I; one record only

Habitat: 17 m; muddy sand

Genus **Papyriscala** Boury, 1909

Type species: *Scalaria latifasciata* Sowerby, 1878

148. ***Papyriscala tricincta*** Golikov in Golikov and Scarlato, 1967

[*Papyriscala tricincta* Golikov and Scarlato, 1967: 49, fig. 37; Kantor and Sysoev, 2006: 130, pl. 59 C].
Type locality: Posjet Bay (Peter the Great Bay, East Sea).

Distribution: Asian low-boreal species; I; one record only

Habitat: 1.5–5 m; muddy sand

Superfamily Eulimoidea Philippi, 1853

Family Eulimidae Philippi, 1853

Genus **Amamibalcis** Kuroda et Habe, 1950

Type species: *Amamibalcis kawamurai* Kuroda and Habe, 1950.

149. ***Amamibalcis yessoensis*** Rybakov and Yakovlev, 1993

[*Amamibalcis yessoensis* Rybakov and Yakovlev, 1993: 47–49, fig. 1a–e; Higo et al., 1999: 191; Kantor and Sysoev, 2006: 130, pl. 59 C].

Type locality: Moneron Island, East Sea.

Distribution: Asian low-boreal species; I, III, VIII; single records only

Habitat: 15–20 m; parasitic in holothurians

Genus **Melanella** Bowdich, 1822

Type species: *Melanella dufrensei* Bowdich, 1822.

150. ***Melanella conspicuus*** (Golikov in Golikov and Scarlato, 1985)

[*Balcis conspicuuus* Golikov and Scarlato, 1985: 438, fig. 16; *Melanella conspicuous*: Kantor and Sysoev, 2006: 131, pl. 59 J].

Type locality: Tatar Strait (northern part)

Distribution: Asian low-boreal species; IV; one record only
Habitat: 34 m; mud

151. ***Melanella kuronamako*** (Habe, 1952)

[*Balcis kuronamako* Habe 1952: 73–74, pl. VI, fig. 20; *Melanella kuronamako*: Higo et al., 1999: 187; Kantor and Sysoev, 2006: 131].

Type locality: Kii Peninsula, central Honshu

Distribution: Asian low-boreal species; I, II, III, IV, VI, VII, VIII; common.

Habitat: intertidal ~40 m; rock and mud, parasitic on echinoderms

Superfamily Triphoroidea Gray, 1847

Family Cerithiopsidae H. et A. Adams, 1854

Genus **Cerithiopsis** Forbes et Hanley, 1849

Type species: *Murex tuberculatus* Montagu, 1803.

152. ***Cerithiopsis janira*** Bartsch in Golikov and Scarlato, 1967

[*Cerithiopsis janira* Golikov and Scarlato, 1967: 42–43, fig. 33; Kantor and Sysoev, 2006: 135, pl. 60 J].

Type locality: East Sea.

Distribution: Asian low-boreal species; I, II, IV, VIII; single records only

Habitat: intertidal ~100 m; sand with shells, boulders

153. *Cerithiopsis stejnegeri* Dall, 1884

[*Cerithiopsis stejnegeri* Dall, 1884: 345-346, pl. 2, fig. 4; Higo et al., 1999: 172; Min et al., 2004: 191, fig. 467; Kantor and Sysoev, 2006: 135, pl. 60 F].

Type locality: Kamchatka Island, Aleutian Islands.

Distribution: Pacific boreal species; I, VII, VIII; common.

Habitat: intertidal -20 m; rock and stones

Family Newtoniellidae Korobkov, 1955

Subfamily Eumetullinae Golikov and Starobogatov, 1975

Genus *Furukawaia* Kuroda and Habe, 1954

Type species: *Furukawaia fukuensis* Kuroda and Habe in Habe, 1961.

154. *Furukawaia fukuensis* Kuroda and Habe in Habe, 1961

[*Furukawaia fukuensis* Habe, 1961: 28, app.9, pl. 12, fig. 26; Higo et al., 1999: 173; Kantor and Sysoev, 2006: 138, pl. 61 L F].

Type locality: Fukui Prefecture, Honshu (East Sea).

Distribution: Asian subtropical-low-boreal species; II, III, VII, VIII; uncommon.

Habitat: 10-100 m; sand with pebbles and shells, stones

155. *Furukawaia habei* Golikov and Gulbin, 1978

[*Furukawaia habei* Golikov and Gulbin, 1978: 219-220, figs. 24, 8, 25 Higo et al., 1999: 173; 138, pl. 61 K].

Type locality: Zelenyi Island, southern Kuril Islands

Distribution: Asian boreal species IV, VIII rare.

Habitat: 70-115 m; sand, and sand with shells

Genus *Laskeya* Iredale 1919

Type species: *Tuerritella costulata* Møller, 1842.

156. *Laskeya striata* (Gulbin, 1982)

[*Eumetula striata* Gulbin, 1982: 70-71, text-fig.; Kantor and Sysoev, 2006: 137-138, pl.61 H].

Type locality: Bering Island, Commander Islands.

Distribution: Asian boreal species VIII; one record only

Habitat: 115 m; sand with shells

ACKNOWLEDGEMENTS

This study was supported by grants from Far Eastern Branch Russian Academy of Sciences, grant number DVO-1 N 09-I-P23-01 and DVO-1 N 09-I-P15-03.

REFERENCES

- Adams, A. (1861a). On some new species of Mollusca from the North of China and Japan. *Annals and Magazine of Natural History, series 3*, **8**: 135-142.
- Adams, A. (1861b). On some new genera and species of Mollusca from the North of China and Japan. *Annals and Magazine of Natural History, series 3*, **8**: 239-246, 299-309.
- Adams, A. (1867) Description of new species of shells from Japan. *Proceedings of the Zoological Society of London*, part 2: 309-315.
- Bergh, R. (1853) Bidrag tilen monographi of Marseniaderne, en Familie of de Gastraepode Mollusker. 119 pp. Kjøbenhavn, Biaco Lunos Bogtrykkeri.
- Broderip, W.J. and Sowerby, G.B. (1829) Observations on new or interesting Mollusca contained, for the most part, in the Museum of the Zoological Society. *Zoological Journal*, **4**: 359-379.
- Bouchet, Ph. and Rocroi, J.-P. (2005) Classification and nomenclature of gastropod families. *Malacologia*, **47**(1-2): 1-397.
- Carpenter, P.P. (1864) Supplementary report on the present state of our knowledge with regard to the Mollusca of the west coast of North America. In: Report of the British Association for the Advancement of Science, for 1863. 517-686 pp.
- Couthouy, J.P. (1838) Descriptions of new species of Mollusca and shells and remarks on several polypii found in Massachusetts bay. *Boston Journal of Natural History*, **2**(1): 53-111.
- Crosse, H. (1862) Description d'espèces marines recueillies par M.H.Cuming dans le Nord de la Chine. *Journal de Conchyliologie*, **10**: 51-57.
- Dall, W.H. (1871) Descriptions of sixty new forms of mollusks from the west coast of North America and the North Pacific Ocean, with notes on others already described. *American Journal of Conchology*, **7**(2): 93-160.
- Dall, W.H. (1884) Contributions to the history of Commander Islands. No. 3. Report on the Mollusca of the Commander Islands, Bering Sea, collected by Leonhard Stejneger in 1882 and 1883. *Proceedings of the United States National Museum*, **7**(442): 340-349.
- Dall, W.H. (1887 [1886]). Supplementary notes on some species of mollusks of the Bering Sea and vicinity. *Proceedings of the United States National Museum*,

Review of the Shell-bearing Gastropods in the Russian Waters of the East Sea. II. Caenogastropoda: Sorbeoconcha, Hypsogastropoda

- 9(571): 297-309.
- Dall, W.H. (1919) Descriptions of new species of Mollusca from the North Pacific Ocean in the collection of the United States National Museum. *Proceedings of the United States National Museum*, **56**(2295): 293-371.
- Derjugin, K.M. (1937) New data on systematic, morphology and biogeography of the genus *Onchidiopsis* (Gastropoda). *Scientific Notices Leningrad University*, **5**, 7-31. [in Russian].
- Derjugin, K.M. (1950). New data on systematic, morphology and biogeography of the genus *Velutina* Flem. (Mollusca, Gastropoda, Lamellariidae). *Explorations Far East seas USSR*, **2**, 7-27. [in Russian].
- Deshayes, G.P. (1839) Nouvelles especes de mollusques, provenant des cotes de la Californie, du Mexique, de Kamtschatka et de Nouvelle-Zelande. *Revue Zoologique de la Societe Cuvierienne*, **2**(32): 356-361.
- Egorov, R.V. and Alexeev, D.O. (1998) Trichotropidae. Treasure of Russian Shells, 2. Moscow, 36 p.
- Gmelin, J.F. (1791) Caroli a Linne Systema naturae per regna tria naturae. Editio decima tertia. Vol. 1(6), Vermes: 3021-3910. Lipsiae, R. Poll.
- Golikov, A.N. (1986a). Gastropods of family Trichotropidae in temperate and cold waters of the Northern Hemisphere. *Proceedings of the Zoological Institute, Leningrad*, **152**: 11-29. [in Russian].
- Golikov, A.N. (1986b). New species of gastropods from the suborder Rissoidae. *Proceedings of the Zoological Institute, Leningrad*, **148**: 86-89. [in Russian].
- Golikov, A.N. and Gulbin, V.V. (1978) Prosobranchial gastropods of the Kurile Islands. I. Orders Docoglossa - Entomostoma. In: Fauna of coastal zones of Kurile Islands, pp. 159-223. Nauka, Moscow. [in Russian]
- Golikov, A.N., Gulbin, V.V. and Sirenko, B.I. (1987) Prosobranchial gastropods of the shelf of Moneron Island (the Sea of Japan). I. Ordo Patelliformes - Calyptraeiformes. In: Fauna and distribution of mollusks: Northern Pacific and Polar Basin (ed. by Kafanov, A.I.), pp. 22-40. Far East Center of the USSR Academy of Sciences. Vladivostok. [in Russian].
- Golikov, A.N. and Kussakin, O.G. (1962) Fauna and ecology of prosobranchs (Gastropoda, Prosobranchia) of intertidal zone of Kurile Islands. *Explorations Far East seas USSR*, **8**: 248-346. [in Russian].
- Golikov, A.M. and Kussakin O.G. (1978) Shell-bearing gastropods of the intertidal zone of the seas of the USSR. Leningrad, Nauka. 292 p. [in Russian].
- Golikov, A.N. and Scarlato, O.A. (1967) Molluscs of the Possiet Bay (the Sea of Japan) and their ecology. *Proceedings of the Zoological Institute, Leningrad*. **42**: 5-154. [in Russian].
- Golikov, A.N. and Scarlato O.A. (1971) On the fauna of the molluscs of Possiet Bay of the Sea of Japan. *Explorations of the Fauna of the seas*, **8**: 188-205.
- [in Russian].
- Golikov, A.N. and Scarlato, O.A. (1985) Shell-bearing gastropod and bivalve molluscs of the shelf of southern Sakhalin and their ecology. *Explorations of the Fauna of the seas*, **30**: 360-487. [in Russian].
- Gould, A.A. (1846-1850) Shells collected by the United States Exploring Expedition under the command of Charles Wilkes. *Proceedings of the Boston Society of Natural History*, (1846), **2**: 141-145; 148-152; 153-156; 159-162; 165-167; 170-173; 175-176; 1847: 196-198; 200-203; 204-208; 209; 210-212; 214-215; 222-224; 225; 237-239; 251-252.
- Gould, A.A. (1860) Descriptions of shells collected in the North Pacific Exploring Expedition under Captains Ringgold and Rodgers. *Proceedings of the Boston Society for Natural History*, **7**: 323-336; 337-340; 382-384.
- Gulbin, V.V. 1982. New gastropod mollusc *Eumetula striata* from Commander Islands. *Biologiya morya*, **3**: 70-71. [in Russian]
- Gulbin, V.V. and Golikov, A.N. (1997) A review of the prosobranch family Velutinidae in cold and temperate waters of the Northern Hemisphere. I. Capulacmaeinae. *Ophelia*, **49**(1): 43-54.
- Habe, T. (1952) Parasitic gastropods found in echinoderms from Japan. *Publications of the Seta Marine Biological Laboratory*, **2**(2): 73-85.
- Habe, T. (1953) Descriptions of twelve new Japanese species. *Venus*, **17**(3): 130-144.
- Habe, T. (1958) The fauna of Akkeshi Bay XXV. Gastropoda. *Publications from the Akkeshi Marine Biological Station, Hokkaido University, Sapporo*, **8**: 1-40.
- Habe, T. (1961) Colored illustrations of the shells of Japan, vol. II. 183 pp. Osaka, Hoikusha.
- Habe, T. and Ito K. (1976). The new naticid species (Mollusca) from Hokkaido. *Bulletin of the National Science Museum, Series A (Zoology)*, **2**(2): 79-82.
- Higo, Sh., Callomon, P. and Gotō, Y. (1999) Catalogue and Bibliography of the Marine Shell-bearing Mollusca of Japan. 749 pp. Elle Scientific Publication. Ōsaka.
- Kantor, Yu.I. and Sysoev A.V. (2006) Marine and brackish water Gastropoda of Russia and adjacent countries: an illustrated catalogue. 371 pp. KMK Scientific Press Ltd. Moscow.
- Kuroda, T. (1962) Notes on Stenothyridae (aquatic Gastropoda) from Japan and adjacent regions. *Venus*, **22**(1): 59-69.
- Lindholm, W.A. (1929) Einige neue Mollusken (Pelecypoda und Gastropoda) aus den Gewässern Sudost-Sibiriens. *Доклады Академии Наук*. 12: 302-306.
- Middendorff, A.T. (1848) Vorläufige Anzeige einiger neuer Konchylien Russland's, aus den Geschlechtern: *Scalaria*, *Crepidula*, *Velutina*, *Trichotropis*, *Purpura* und *Pleurotoma*. *Bulletin de la Classe Physico-Mathematique de l' Academie Impériale des Sciences de Saint-Petersbourg*, (2): 1-4

- (pagination according to reprint).
- Middendorff, A.T. (1849) Beiträge zu einer Malacozoologia Rossica. II. Aufzählung und Beschreibung der zur Meeresfauna Russlands gehörigen Einschaler. *Mémoires de l'Académie Impériale des Sciences de Saint-Pétersbourg, Serie 6. Sciences Mathématique, Physique et Naturelles*, **8**(5-6): 329-516 (Reprint pp. 1-187)
- Middendorff, A.T. (1851) Mollusken. In: Reise in den dusserten Norden und Osten Sibiriens. Band H. Zoologie. Theil I. 163-464 pp. Wirbellose Thiere. St. Petersburg, Kaiserliche Akademie der Wissenschaften.
- Migheis, J.W. and Adams C.B. (1842) Descriptions of twenty-four species of the shells of New England. *Boston Journal of Natural History*, **4**: 37-54.
- Min, D.K., J.S. Lee, D.B. Koh and J.G. (2004) Mollusks in Korea. 566 pp. Min Molluscan Research Institute, Seoul. [in Korean].
- Møller, H.P.C (1842) Index Molluscorum Groenlandiae. *Naturhistorisk Tidsskrift*, **4**(1): 76-97.
- Müller, O.F. (1776) Zoologie Danicae prodromus, seu Animalium Daniae et Norvegiae indigenarum characters, nomina et synonyma imprimis popularium. 282 pp. Havniae.
- Pallas, P.S. (1788) Marina varia nova et rarioса. *Nova Ada Academia Petropolitanae*, **2**: 223-250.
- Perry, G. (1811) Conchology, or the natural history of shells, containing the new arrangement of the genera and species. 61 p. London.
- Philippi, R.A. (1844) Descriptiones testaceorum quorundam novorum, maxime chinensium. *Zeitschrift jur Malakozoolgie*, **1844**: 161-167.
- Philippi R.A. (1846) Descriptions of a new species of Trochus and eighteen species of *Littorina*, in the collections of H. Cuming, Esq. *Proceedings of the Zoological Society of London*, **13**: 138-143.
- Philippi, R.A. (1849) Centuria tertia Testacorum novorum. *Zeitschrift fur Malakozoolgie*, **5** (1848): 151-176, 186-192.
- Pilsbry, H.A. (1901) New Mollusca from Japan, the Loo-Choo islands, Formosa and the Philippines. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **53**: 385-408.
- Pilsbry, H.A. (1905) New Japanese Marine Mollusca. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **57**(1): 101-122.
- Pilsbry, H.A. (1911) New Japanese Naticidae and Scalariidae. *Nautilus*, **25**: 32-34.
- Prozorova, L.A. (1996) The species of the genus Cecina (Gastropoda, Pomatiopsidae) in Far East of Russia. *Zoological Zhurnal*, **75**(5): 653-663. [in Russian]
- Redfield, J.H. (1846) Description of some new species of shells (*Marginea flava*, *M. obesa*, *Triton oregonense*, *Ranella thersites* and *Cardium setosum*). *Annals of the Lyceum of Natural History of New York*, **4**: 163-168.
- Röding, P.F. (1789) Index, Museum Boltenianum. 48 p. Hamburg.
- Rybakov, A.V. and Yakovlev Yu.M. (1993) *Amamibalcis yessoensis* n. sp. (Gastropoda: Eulimidae) - parasite of holothurians from Sea of Japan. *Venus*, **52**(1): 47-49.
- Schrenek, L. (1867) Mollusken des Amur-Landes und des Nordjapanischen Meeres. In: Reisen und Forschungen im Amur-Lands in der Jahren 1854-1856, vol. 2: 259-976. SPb.
- Smith, E.A. (1875) A list of the Gastropoda collected in Japanese seas by commander H. C. St. John. R.N. *Annals and Magazine of Natural History*, series 4, **16**: 103-115,
- Sowerby, G.B. (1914) Descriptions of fifteen new Japanese marine mollusca. *Annals and Magazine of Natural History*, series 8, **14**: 33-39.
- Sowerby, G.B. (1915) Descriptions of new species of Mollusca from various localities. *Annals and Magazine of Natural History*, series 8, **16**(93): 164-170.
- Yokoyama, M. (1926) Fossil shells from Sado. *Journal of the Faculty of Science, Imperial University, Tokyo*, series 2, **1**(8): 249-312.