

INTEGRAL REPRESENTATIONS FOR SRIVASTAVA'S HYPERGEOMETRIC FUNCTION H_A

JUNESANG CHOI, ANVAR HASANOV AND MAMASALI TURAEV

Abstract. While investigating the Lauricella's list of 14 complete second-order hypergeometric series in three variables, Srivastava noticed the existence of three additional complete triple hypergeometric series of the second order, which were denoted by H_A , H_B and H_C . Each of these three triple hypergeometric functions H_A , H_B and H_C has been investigated extensively in many different ways including, for example, in the problem of finding their integral representations of one kind or the other. Here, in this paper, we aim at presenting further integral representations for the Srivastava's triple hypergeometric function H_A .

1. Introduction and Preliminaries

In the theory of hypergeometric functions of several variables, a remarkably large number of triple hypergeometric functions have been introduced and investigated. A comprehensive table of 205 distinct triple hypergeometric functions is provided in the work of Srivastava and Karlsson [15, Chapter 3]. Out of these 205 distinct triple hypergeometric functions, Lauricella [8, p. 114] introduced fourteen complete triple hypergeometric functions of the second order. He denoted his triple hypergeometric functions by the symbols F_1, \dots, F_{14} of which F_1, F_2, F_3 and F_9 correspond, respectively, to the three variable Lauricella functions $F_A^{(3)}$, $F_B^{(3)}$, $F_C^{(3)}$ and $F_D^{(3)}$ that are the three-variable cases of the n -variable Lauricella functions $F_A^{(n)}$, $F_B^{(n)}$, $F_C^{(n)}$ and $F_D^{(n)}$

Received January 25, 2012. Accepted March 7, 2012.

2000 Mathematics Subject Classification. Primary 33C20, 33C65; Secondary 33C05, 33C60, 33C70, 68Q40, 11Y35.

Key words and phrases. Multiple hypergeometric functions, Gauss hypergeometric function ${}_2F_1$, Confluent hypergeometric functions, Eulerian integrals, Laplace integrals, Srivastava's triple hypergeometric function H_A , Exton's functions, Humbert functions, Bessel functions, Beta and Gamma functions, Appell functions, Picard's integral formula..

(cf. [8, p. 113]; see also [1, p. 114, Equations (1) to (4)], [15, p. 33 *et seq.*] and [5, 6]). Saran [10] initiated a systematic study of these ten triple hypergeometric functions from Lauricella's set. Exton [4] introduced 20 distinct triple hypergeometric functions, which he denoted by X_1, \dots, X_{20} , and investigated their twenty Laplace integral representations whose kernels include the confluent hypergeometric functions ${}_0F_1$ and ${}_1F_1$, and the Humbert function hypergeometric functions Ψ_2 and Φ_2 of two variables. The four Appell hypergeometric functions F_1, \dots, F_4 of two variables are simply the special case of Lauricella's n -variable functions when $n = 2$, that is,

$$F_1 = F_D^{(2)}, \quad F_2 = F_A^{(2)}, \quad F_3 = F_B^{(2)} \quad \text{and} \quad F_4 = F_C^{(2)}.$$

While transforming Pochhammer's double-loop contour integrals associated with the functions F_8 and F_{14} (that is, F_G and F_F , respectively) belonging to Lauricella's set of hypergeometric functions of three variables, Srivastava [11, 12] discovered the existence of three additional complete triple hypergeometric functions H_A , H_B and H_C of the second order, of which H_A is defined as follows (see also [15, p. 43, Equation 1.5(11)]):

$$(1.1) \quad \begin{aligned} & H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) \\ &= \sum_{m, n, p=0}^{\infty} \frac{(a_1)_{m+p} (a_2)_{m+n} (a_3)_{n+p}}{(c_1)_m (c_2)_{n+p}} \frac{x^m}{m!} \frac{y^n}{n!} \frac{z^p}{p!} \\ & (|x| =: \mathbf{r} < 1; |y| =: \mathbf{s} < 1; |z| =: \mathbf{t} < (1 - \mathbf{r})(1 - \mathbf{s})), \end{aligned}$$

where, with \mathbb{C} and \mathbb{Z}_0^- denoting the set of complex numbers and the set of nonpositive integers, respectively, $(\lambda)_n$ is the Pochhammer symbol defined (for $\lambda \in \mathbb{C}$) by

$$(1.2) \quad (\lambda)_n := \frac{\Gamma(\lambda + n)}{\Gamma(\lambda)} = \begin{cases} 1 & (n = 0) \\ \lambda(\lambda + 1) \dots (\lambda + n - 1) & (n \in \mathbb{N} := \{1, 2, 3, \dots\}), \end{cases}$$

Γ being the well-known Gamma function. Of course, all 20 of Exton's triple hypergeometric functions X_1, \dots, X_{20} as well as Srivastava's triple hypergeometric functions H_A , H_B and H_C are included in the set of the aforementioned 205 distinct triple hypergeometric functions which were presented systematically by Srivastava and Karlsson [15, Chapter 3]. The above-stated three-dimensional region of convergence of the triple hypergeometric function in (1.1) for H_A was given by Srivastava [11, 12] (see also Srivastava and Karlsson [15, Section 3.4]).

Various multivariable generalizations and cases of reducibility of Srivastava's functions H_A , H_B and H_C have been investigated (see, for details, [15, pp. 43–44]). Turaev [17] studied the Srivastava function H_A . Hasanov *et al.* [7] reproduced Srivastava's integral representations for the functions [11, 12] H_A , H_B and H_C . Very recently, Choi *et al.* [2] also presented certain integral representations for the functions H_A , H_B and H_C .

Here, in this present sequel to some of the above-mentioned works, we aim at investigating *further* 19 integral representations for the Srivastava function H_A , for completeness, including the five ones in [2].

2. Integral Representations for Srivastava's hypergeometric function H_A

Theorem. *Each of the following integral representations for H_A holds true.*

(2.1)

$$H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(s)}{\Gamma(a_1)\Gamma(s-a_1)} \int_0^1 \xi^{a_1-1} (1-\xi)^{s-a_1-1} H_A(s, a_2, a_3; c_1, c_2; x\xi, y, z\xi) d\xi$$

$$(\Re(s) > \Re(a_1) > 0);$$

(2.2)

$$H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(s)}{\Gamma(a_2)\Gamma(s-a_2)} \int_0^1 \xi^{a_2-1} (1-\xi)^{s-a_2-1} H_A(a_1, s, a_3; c_1, c_2; x\xi, y\xi, z) d\xi$$

$$(\Re(s) > \Re(a_2) > 0);$$

(2.3)

$$H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(s)}{\Gamma(a_3)\Gamma(s-a_3)} \int_0^1 \xi^{a_3-1} (1-\xi)^{s-a_3-1} H_A(a_1, a_2, s; c_1, c_2; x, y\xi, z\xi) d\xi$$

$$(\Re(s) > \Re(a_3) > 0);$$

$$H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(c_1)}{\Gamma(c_1-s)\Gamma(s)} \cdot \int_0^1 \xi^{s-1} (1-\xi)^{c_1-s-1} H_A(a_1, a_2, a_3; s, c_2; x\xi, y, z) d\xi$$

(2.4)
$$(\Re(c_1) > \Re(s) > 0);$$

$$(2.5) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(c_2)}{\Gamma(c_2 - s)\Gamma(s)} \cdot \int_0^1 \xi^{s-1} (1 - \xi)^{c_2 - s - 1} H_A(a_1, a_2, a_3; c_1, s; x, y\xi, z\xi) d\xi$$

$$(\Re(c_2) > \Re(s) > 0);$$

$$(2.6) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(c_2)}{\Gamma(a_3)\Gamma(c_2 - a_3)} \int_0^1 \xi^{a_3 - 1} (1 - \xi)^{c_2 - a_3 - 1} \cdot (1 - y\xi)^{-a_2} (1 - z\xi)^{-a_1} {}_2F_1\left(a_1, a_2; c_1; \frac{x}{(1 - y\xi)(1 - z\xi)}\right) d\xi$$

$$(\Re(c_2) > \Re(a_3) > 0);$$

$$(2.7) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(c_2)(1 + \lambda)^{a_3}}{\Gamma(a_3)\Gamma(c_2 - a_3)} \cdot \int_0^1 \xi^{a_3 - 1} (1 - \xi)^{c_2 - a_3 - 1} \cdot (1 + \lambda\xi)^{a_1 + a_2 - c_2} [1 + \lambda\xi - (1 + \lambda)\xi y]^{-a_2} [1 + \lambda\xi - (1 + \lambda)\xi z]^{-a_1} \cdot {}_2F_1\left(a_1, a_2; c_1; \frac{x(1 + \lambda\xi)^2}{[1 + \lambda\xi - (1 + \lambda)\xi y][1 + \lambda\xi - (1 + \lambda)\xi z]}\right) d\xi$$

$$(\Re(c_2) > \Re(a_3) > 0; \lambda > -1);$$

$$(2.8) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(c_2)}{\Gamma(a_3)\Gamma(c_2 - a_3)} \frac{(\beta - \gamma)^{a_3} (\alpha - \gamma)^{c_2 - a_3}}{(\beta - \alpha)^{c_2 - a_1 - a_2 - 1}} \cdot \int_\alpha^\beta (\beta - \xi)^{c_2 - a_3 - 1} (\xi - \alpha)^{a_3 - 1} (\xi - \gamma)^{a_1 + a_2 - c_2} \cdot [(\beta - \alpha)(\xi - \gamma) - (\beta - \gamma)(\xi - \alpha)y]^{-a_2} \cdot [(\beta - \alpha)(\xi - \gamma) - (\beta - \gamma)(\xi - \alpha)z]^{-a_1} \cdot {}_2F_1(a_1, a_2; c_1; \sigma x) d\xi$$

$$(\Re(c_2) > \Re(a_3) > 0; \gamma < \alpha < \beta),$$

where σ is defined by

$$\sigma = \frac{(\beta - \alpha)^2 (\xi - \gamma)^2}{[(\beta - \alpha)(\xi - \gamma) - (\beta - \gamma)(\xi - \alpha)y][(\beta - \alpha)(\xi - \gamma) - (\beta - \gamma)(\xi - \alpha)z]};$$

$$\begin{aligned}
 H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) &= \frac{\Gamma(c_2)}{\Gamma(a_3)\Gamma(c_2 - a_3)} \frac{(\gamma - \beta)^{a_3} (\gamma - \alpha)^{c_2 - a_3}}{(\beta - \alpha)^{c_2 - a_1 - a_2 - 1}} \\
 (2.9) \quad &\cdot \int_{\alpha}^{\beta} (\beta - \xi)^{c_2 - a_3 - 1} (\xi - \alpha)^{a_3 - 1} (\gamma - \xi)^{a_1 + a_2 - c_2} \\
 &\cdot [(\beta - \alpha)(\gamma - \xi) - (\gamma - \beta)(\xi - \alpha)y]^{-a_2} \\
 &\cdot [(\beta - \alpha)(\gamma - \xi) - (\gamma - \beta)(\xi - \alpha)z]^{-a_1} \\
 &\cdot {}_2F_1(a_1, a_2; c_1; \sigma x) d\xi \\
 &\quad (\Re(c_2) > \Re(a_3) > 0; \alpha < \beta < \gamma),
 \end{aligned}$$

where σ is defined by

$$\sigma = \frac{(\beta - \alpha)^2 (\gamma - \xi)^2}{[(\beta - \alpha)(\gamma - \xi) - (\gamma - \beta)(\xi - \alpha)y][(\beta - \alpha)(\gamma - \xi) - (\gamma - \beta)(\xi - \alpha)z]};$$

$$\begin{aligned}
 (2.10) \quad &H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
 &= \frac{\Gamma(c_2)}{\Gamma(a_3)\Gamma(c_2 - a_3)} \int_0^{\infty} \xi^{a_3 - 1} (1 + \xi)^{a_1 + a_2 - c_2} (1 + \xi - y\xi)^{-a_2} \\
 &\cdot (1 + \xi - z\xi)^{-a_1} {}_2F_1\left(a_1, a_2; c_1; \frac{x(1 + \xi)^2}{(1 + \xi - y\xi)(1 + \xi - z\xi)}\right) d\xi \\
 &\quad (\Re(c_2) > \Re(a_3) > 0);
 \end{aligned}$$

$$\begin{aligned}
 (2.11) \quad &H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
 &= \frac{\Gamma(c_2)}{\Gamma(a_3)\Gamma(c_2 - a_3)} \int_0^{\infty} (e^{-\xi})^{a_3} (1 - e^{-\xi})^{c_2 - a_3 - 1} \\
 &\cdot (1 - ye^{-\xi})^{-a_2} (1 - ze^{-\xi})^{-a_1} {}_2F_1\left(a_1, a_2; c_1; \frac{x}{(1 - ye^{-\xi})(1 - ze^{-\xi})}\right) d\xi \\
 &\quad (\Re(c_2) > \Re(a_3) > 0);
 \end{aligned}$$

$$\begin{aligned}
& H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
&= \frac{2\Gamma(c_2)}{\Gamma(a_3)\Gamma(c_2 - a_3)} \int_0^{\frac{\pi}{2}} (\sin^2 \xi)^{a_3 - \frac{1}{2}} (\cos^2 \xi)^{c_2 - a_3 - \frac{1}{2}} \\
(2.12) \quad & \cdot (1 - y \sin^2 \xi)^{-a_2} (1 - z \sin^2 \xi)^{-a_1} \\
& \cdot {}_2F_1 \left(a_1, a_2; c_1; \frac{x}{(1 - y \sin^2 \xi)(1 - z \sin^2 \xi)} \right) d\xi \\
& \quad (\Re(c_2) > \Re(a_3) > 0);
\end{aligned}$$

$$\begin{aligned}
(2.13) \quad & H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
&= \frac{2\Gamma(c_2)(1 + \lambda)^{a_3}}{\Gamma(a_3)\Gamma(c_2 - a_3)} \int_0^{\frac{\pi}{2}} \frac{(\sin^2 \xi)^{a_3 - \frac{1}{2}} (\cos^2 \xi)^{c_2 - a_3 - \frac{1}{2}}}{(1 + \lambda \sin^2 \xi)^{c_2 - a_1 - a_2}} \\
& \cdot [1 + \lambda \sin^2 \xi - (1 + \lambda)y \sin^2 \xi]^{-a_2} [1 + \lambda \sin^2 \xi - (1 + \lambda)z \sin^2 \xi]^{-a_1} {}_2F_1 \\
& \cdot \left(a_1, a_2; c_1; \frac{x(1 + \lambda \sin^2 \xi)^2}{[1 + \lambda \sin^2 \xi - (1 + \lambda)y \sin^2 \xi][1 + \lambda \sin^2 \xi - (1 + \lambda)z \sin^2 \xi]} \right) \\
& \cdot d\xi \quad (\Re(a_3) > 0; \Re(c_2 - a_3) > 0; \lambda > -1);
\end{aligned}$$

$$\begin{aligned}
(2.14) \quad & H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
&= \frac{2\Gamma(c_2)\lambda^{a_3}}{\Gamma(a_3)\Gamma(c_2 - a_3)} \int_0^{\frac{\pi}{2}} \frac{(\sin^2 \xi)^{a_3 - \frac{1}{2}} (\cos^2 \xi)^{c_2 - a_3 - \frac{1}{2}}}{(\cos^2 \xi + \lambda \sin^2 \xi)^{c_2}} \\
& \cdot \left(\frac{\cos^2 \xi + \lambda \sin^2 \xi - \lambda y \sin^2 \xi}{\cos^2 \xi + \lambda \sin^2 \xi} \right)^{-a_2} \left(\frac{\cos^2 \xi + \lambda \sin^2 \xi - \lambda z \sin^2 \xi}{\cos^2 \xi + \lambda \sin^2 \xi} \right)^{-a_1} \\
& \cdot {}_2F_1 \left(a_1, a_2; c_1; \frac{x(\cos^2 \xi + \lambda \sin^2 \xi)^2}{[\cos^2 \xi + \lambda \sin^2 \xi - \lambda y \sin^2 \xi][\cos^2 \xi + \lambda \sin^2 \xi - \lambda z \sin^2 \xi]} \right) \\
& \cdot d\xi \quad (\Re(a_3) > 0; \Re(c_2 - a_3) > 0; \lambda > 0);
\end{aligned}$$

$$\begin{aligned}
 (2.15) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) &= \frac{\Gamma(a_1 + a_2) \Gamma(c_2)}{\Gamma(a_1) \Gamma(a_2) \Gamma(a_3) \Gamma(c_2 - a_3)} \\
 &\cdot \int_0^1 \int_0^1 \xi^{a_1-1} \eta^{a_3-1} (1-\xi)^{a_2-1} (1-\eta)^{c_2-a_3-1} (1-y\eta - z\xi\eta + y\xi\eta)^{-a_1-a_2} \\
 &\cdot {}_2F_1\left(\frac{a_1+a_2}{2}, \frac{a_1+a_2}{2} + \frac{1}{2}; c_1; \frac{4x\xi(1-\xi)}{(1-y\eta - z\xi\eta + y\xi\eta)^2}\right) d\xi d\eta \\
 &(\Re(a_1) > 0; \Re(a_2) > 0; \Re(c_2) > \Re(a_3) > 0);
 \end{aligned}$$

$$\begin{aligned}
 (2.16) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) &= \frac{\Gamma(a_1 + a_2) \Gamma(c_2) (1+\lambda)^{a_3}}{\Gamma(a_1) \Gamma(a_2) \Gamma(a_3) \Gamma(c_2 - a_3)} \\
 &\cdot \int_0^1 \int_0^1 \xi^{a_1-1} \eta^{a_3-1} (1-\xi)^{a_2-1} (1-\eta)^{c_2-a_3-1} (1+\lambda\eta)^{a_1+a_2-c_2} \\
 &\cdot [(1+\lambda\eta) - (1+\lambda)z\xi\eta - (1+\lambda)y(1-\xi)\eta]^{-a_1-a_2} {}_2F_1 \\
 &\cdot \left(\frac{a_1+a_2}{2}, \frac{a_1+a_2}{2} + \frac{1}{2}; c_1; \frac{4x\xi(1-\xi)(1+\lambda\eta)^2}{[(1+\lambda\eta) - (1+\lambda)z\xi\eta - (1+\lambda)y(1-\xi)\eta]^2}\right) \\
 &\cdot d\xi d\eta \quad (\Re(a_1) > 0; \Re(a_2) > 0; \Re(c_2) > \Re(a_3) > 0; \lambda > -1);
 \end{aligned}$$

$$\begin{aligned}
 (2.17) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) &= \frac{\Gamma(a_1 + a_2)}{\Gamma(a_1) \Gamma(a_2)} \int_0^1 \xi^{a_1-1} (1-\xi)^{a_2-1} \\
 &\cdot H_4[a_1 + a_2, a_3; c_1, c_2; x\xi(1-\xi), y(1-\xi) + z\xi] d\xi \\
 &(\Re(a_1) > 0; \Re(a_2) > 0);
 \end{aligned}$$

$$\begin{aligned}
 (2.18) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) &= \frac{\Gamma(a_1 + a_2) \Gamma(c_2)}{\Gamma(a_1) \Gamma(a_2) \Gamma(s) \Gamma(c_2 - s)} \\
 &\cdot \int_0^1 \int_0^1 \xi^{a_1-1} \eta^{s-1} (1-\xi)^{a_2-1} (1-\eta)^{c_2-s-1} \\
 &\cdot X_4[a_1 + a_2, a_3; c_1, c_2 - s, s; x\xi(1-\xi), y(1-\xi)(1-\eta), z\xi\eta] d\xi d\eta \\
 &(\Re(a_1) > 0; \Re(a_2) > 0; \Re(c_2) > \Re(s) > 0);
 \end{aligned}$$

$$\begin{aligned}
(2.19) \quad & H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
& = \frac{\Gamma(a_1 + a_2 + a_3)}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)} \int_0^1 \int_0^1 \xi^{a_1-1} \eta^{a_1+a_2-1} (1-\xi)^{a_2-1} (1-\eta)^{a_3-1} F^{(3)} \\
& \cdot \left[\begin{array}{c} \frac{a_1+a_2+a_3}{2}, \frac{a_1+a_2+a_3}{2} + \frac{1}{2} \quad \vdots \quad -; \quad -; \quad -; \quad -; \quad -; \quad -; \\ - \quad \vdots \quad -; \quad c_2; \quad -; \quad c_1; \quad -; \quad -; \end{array} \sigma_1 x, \sigma_2 y, \sigma_3 z \right] \\
& \cdot d\xi d\eta \quad (\Re(a_i) > 0; i = 1, 2, 3),
\end{aligned}$$

where σ_1, σ_2 and σ_3 are defined by

$$\sigma_1 = 4\xi(1-\xi)\eta^2, \quad \sigma_2 = 4(1-\xi)\eta(1-\eta) \quad \text{and} \quad \sigma_3 = 4\xi\eta(1-\eta),$$

and ${}_2F_1, H_4, X_4$ and $F^{(3)}$ denote, respectively, Gauss hypergeometric function, Horn's function, Exton function and generalized Srivastava hypergeometric function defined by

$${}_2F_1(a, b; c; x) = \sum_{m=0}^{\infty} \frac{(a)_m (b)_m}{(c)_m m!} x^m,$$

$$H_4(a_1, a_2; c_1, c_2; x, y) = \sum_{m, n=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_n}{(c_1)_m (c_2)_n m! n!} x^m y^n,$$

$$X_4(a_1, a_2; c_1, c_2, c_3; x, y, z) = \sum_{m, n, p=0}^{\infty} \frac{(a_1)_{2m+n+p} (a_2)_{n+p}}{(c_1)_m (c_2)_n (c_3)_p m! n! p!} x^m y^n z^p$$

and

$$F^{(3)}[x, y, z] = F^{(3)} \left[\begin{array}{c} (a) \quad \vdots \quad (b); \quad (b'); \quad (b''); \quad (c); \quad (c'); \quad (c''); \\ (e) \quad \vdots \quad (g); \quad (g'); \quad (g''); \quad (g); \quad (g'); \quad (g''); \end{array} x, y, z \right].$$

Proof. The integral representation (2.6) was derived by Srivastava himself [11, p. 100] as an intermediate result in his demonstration of the following integral representation ([11, p. 100, Equation (3.3)]):

$$\begin{aligned}
(2.20) \quad & H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \frac{\Gamma(c_1)\Gamma(c_2)}{\Gamma(c_1)\Gamma(c_2)\Gamma(c_1-a_2)\Gamma(c_2-a_3)} \\
& \cdot \int_0^1 \int_0^1 \xi^{a_2-1} \eta^{a_3-1} (1-\xi)^{c_1-a_2-1} (1-\eta)^{c_2-a_3-1} (1-y\eta)^{-a_2} \\
& \cdot (1-x\xi-z\eta)^{-a_1} \left(1 - \frac{xy\xi\eta}{(1-y\eta)(1-x\xi-z\eta)} \right)^{-a_1} d\xi d\eta \\
& (\Re(c_1) > \Re(a_2) > 0; \Re(c_2) > \Re(a_3) > 0).
\end{aligned}$$

In fact, Srivastava's derivation of (2.6) involved writing the triple hypergeometric series in (1.1) as a single series of the Appell function F_1 as follows:

$$(2.21) \quad H_A(a_1, a_2, a_3; c_1, c_2; x, y, z) = \sum_{m=0}^{\infty} \frac{(a_1)_m (a_2)_m}{(c_1)_m} F_1[a_3, a_2 + m, a_1 + m; c_2; y, z] \frac{x^m}{m!}$$

and then applying Picard's integral formula [1, p. 29, Equation (4)]:

$$(2.22) \quad F_1[\alpha, \beta, \beta'; \gamma; x, y] = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma - \alpha)} \cdot \int_0^1 \tau^{\alpha-1} (1 - \tau)^{\gamma-\alpha-1} (1 - x\tau)^{-\beta} (1 - y\tau)^{-\beta'} d\tau$$

$(\Re(\gamma) > \Re(\alpha) > 0; \gamma \in \mathbb{C} \setminus \mathbb{Z}_0^-)$

to each term on the right-hand side of (2.21). The transition from (2.6) to Srivastava's final result (2.20) was made by appealing to the following *classical* result (see, for details, [11, pp. 99–100]):

$$(2.23) \quad {}_2F_1(\alpha, \beta; \gamma; z) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma - \alpha)} \int_0^1 \tau^{\alpha-1} (1 - \tau)^{\gamma-\alpha-1} (1 - z\tau)^{-\beta} d\tau$$

$(\Re(\gamma) > \Re(\alpha) > 0; \gamma \in \mathbb{C} \setminus \mathbb{Z}_0^-)$.

Each of the integral representations (2.1) to (2.19) can also be proved *directly* by expressing the series definition of the involved special function in each integrand and changing the order of the integral sign and the summation, and finally using the following well-known relationship between the Beta function $B(\alpha, \beta)$, the Gamma function Γ and their various associated Eulerian integrals (see, for example, [3, pp. 9–11], [13, 14, Section 1.1] and [16, p. 26 and p. 86, Problem 1]):

$$(2.24) \quad B(\alpha, \beta) = \begin{cases} \int_0^1 t^{\alpha-1} (1 - t)^{\beta-1} dt & (\Re(\alpha) > 0; \Re(\beta) > 0) \\ \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)} & (\alpha, \beta \in \mathbb{C} \setminus \mathbb{Z}_0^-), \end{cases}$$

$$(2.25) \quad B(\alpha, \beta) = 2 \int_0^{\frac{\pi}{2}} (\sin \theta)^{2\alpha-1} (\cos \theta)^{2\beta-1} d\theta = \int_0^{\infty} \frac{\tau^{\alpha-1}}{(1 + \tau)^{\alpha+\beta}} d\tau$$

$(\Re(\alpha) > 0; \Re(\beta) > 0)$

and

(2.26)

$$\begin{aligned} B(\alpha, \beta) &= \frac{(b-c)^\alpha (a-c)^\beta}{(b-a)^{\alpha+\beta-1}} \int_a^b \frac{(t-a)^{\alpha-1} (b-t)^{\beta-1}}{(t-c)^{\alpha+\beta}} dt \quad (c < a < b) \\ &= (1+\lambda)^\alpha \int_0^1 \frac{t^{\alpha-1} (1-t)^{\beta-1}}{(1+\lambda t)^{\alpha+\beta}} dt \quad (\lambda > -1) \\ &\quad (\Re(\alpha) > 0; \Re(\beta) > 0). \end{aligned}$$

□

3. Concluding Remarks

Integral representations for most of the special functions of mathematical physics and applied mathematics have been investigated in the existing literature. Here we have presented only some illustrative integral representations for the Srivastava's function H_A . A variety of integral representations of H_A , which may be different from those presented here, can also be provided. Integral representations (2.6), (2.7), (2.8), (2.10) and (2.12) here include the integral representations (2.1), (2.2), (2.3), (2.4) and (2.5), respectively.

Acknowledgements

The work was supported by the *Basic Science Research Program through the National Research Foundation of the Republic of Korea* funded by the Ministry of Education, Science and Technology (2012-0002957).

References

- [1] P. Appell and J. Kampé de Fériet, *Fonctions Hypergeometriques et Hyper-spheriques; Polynomes d'Hermite*, Gauthier - Villars, Paris, 1926.
- [2] J. Choi, A. Hasanov, H. M. Srivastava and M. Turaev, Integral representations for Srivastava's triple hypergeometric functions, *Taiwanese J. Math.* **15** (2011), 2751–2762.
- [3] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher Transcendental Functions*, Vol. I, McGraw-Hill Book Company, New York, Toronto and London, 1953.
- [4] H. Exton, Hypergeometric functions of three variables, *J. Indian Acad. Math.* **4** (1982), 113–119.

- [5] A. Hasanov and H. M. Srivastava, Some decomposition formulas associated with the Lauricella function $F_A^{(r)}$ and other multiple hypergeometric functions, *Appl. Math. Lett.* **19** (2006), 113-121.
- [6] A. Hasanov and H. M. Srivastava, Decomposition formulas associated with the Lauricella multivariable hypergeometric functions, *Comput. Math. Appl.* **53** (2007), 1119-1128.
- [7] A. Hasanov, H. M. Srivastava and M. Turaev, Decomposition formulas for some triple hypergeometric functions, *J. Math. Anal. Appl.* **324** (2006), 955-969.
- [8] G. Lauricella, Sulle funzioni ipergeometriche a più variabili, *Rend. Circ. Mat. Palermo* **7** (1893), 111-158.
- [9] P. A. Padnanabham, Two results on three variable hypergeometric function, *Indian J. Pure Appl. Math.* **30** (1999), 1107-1109.
- [10] S. Saran, Hypergeometric functions of three variables, *Ganita* **5** (1954), 71-91.
- [11] H. M. Srivastava, Hypergeometric functions of three variables, *Ganita* **15** (1964), 97-108.
- [12] H. M. Srivastava, Some integrals representing triple hypergeometric functions, *Rend. Circ. Mat. Palermo* **16** (1967), 99-115.
- [13] H. M. Srivastava and J. Choi, *Series Associated with the Zeta and Related Functions*, Kluwer Academic Publishers, Dordrecht, Boston and London, 2001.
- [14] H. M. Srivastava and J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.
- [15] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.
- [16] H. M. Srivastava and H. L. Manocha, *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1984.
- [17] M. Turaev, Decomposition formulas for Srivastava's hypergeometric function H_A on Saran functions, *J. Comput. Appl. Math.* **233** (2009), 842-846.

Junesang Choi
Department of Mathematics, Dongguk University,
Gyeongju 780-714, Korea.
E-mail: junesang@mail.dongguk.ac.kr

Anvar Hasanov
Department of Mathematics, Dongguk University,
Gyeongju 780-714, Korea.
E-mail: ahasanov@dongguk.ac.kr

Mamasali Turaev
Department of Mathematics, Dongguk University,
Gyeongju 780-714, Korea.
E-mail: mturaev@dongguk.ac.kr