

**CONTINUOUS ORDER REPRESENTABILITY PROPERTIES
OF TOPOLOGICAL SPACES
AND ALGEBRAIC STRUCTURES**

MARÍA JESÚS CAMPIÓN, JUAN CARLOS CANDEAL, ESTEBAN INDURÁIN,
AND GHANSHYAM BHAGVANDAS MEHTA

ABSTRACT. In the present paper, we study the relationship between continuous order-representability and the fulfillment of the usual covering properties on topological spaces. We also consider the case of some algebraic structures providing an application of our results to the social choice theory context.

1. Introduction

A topology τ defined on a nonempty set X is said to satisfy the continuous representability property (CRP) if every continuous total preorder \lesssim defined on X admits a numerical representation by means of a continuous real-valued order-monomorphism (i.e., a continuous map $u : X \rightarrow \mathbb{R}$ such that $x \lesssim y \iff u(x) \leq u(y)$ for every $x, y \in X$). There are different motivations to study this order-representability property (CRP). It is interesting from a topological point of view because it can be used to characterize other topological properties of the given space X . To cite two recent examples, we can say that it has already been used to analyze order-extension properties of topological spaces (see Yi [37] or Campión et al. [5, 7]), as well as to characterize various classical topological properties of a Banach space (see Campión et al. [4]) in functional analysis.

The study of covering properties on topological totally ordered spaces is classical in the mathematical theory of ordered structures (see e.g. Lutzer and Bennet [29]). A further step consists in considering topological spaces (not necessarily endowed a priori with any ordering) and then investigate the family of all the continuous total preorders that may be defined there. This is equivalent to start from a topology τ on a nonempty set X and consider the family of all its subtopologies that are preorderable. Obviously, properties of these

Received April 5, 2010; Revised October 27, 2011.

2010 *Mathematics Subject Classification.* 54F05.

Key words and phrases. ordered structures on topological spaces, order-representability properties, covering properties, preorderable subtopologies, algebraic order-representability, social choice theory.

preorderable subtopologies will reflect properties of the given topological space (X, τ) , and vice versa. In the present paper we work in this direction, paying special attention to covering properties of the preorderable subtopologies of a given topology, in order to analyze the structure, closely related to those covering properties, of topological spaces that satisfy the continuous order-representability property (CRP).

The structure of the paper goes as follows: after the introduction (Section 1) and preliminaries (Section 2), we include a preparatory Section 3 containing results on the topological properties of order topologies, that will be used in the main Section 4 to achieve a characterization of topological spaces satisfying CRP. The key characterization given in Theorem 4.1 is stated in terms of the fulfillment of the second countability axiom for every preorderable subtopology of the given topology. This implies that any topological property that makes every preorderable subtopology to be second countable provides a sufficient condition for CRP to be held. Properties of this kind appear throughout Section 4. In Section 5, an extension of CRP in the algebraic setting, called continuous algebraic representability property (shortly, CARP), is introduced. Basically, whereas CRP asks for continuous numerical representations, CARP asks for those which are continuous and, in addition, preserve the algebraic structure. In Section 6, we develop an application of the previous algebraic approach to social choice theory.

2. Preliminaries

A *preorder* \preceq on an arbitrary nonempty set X is a binary relation on X which is reflexive and transitive.

An antisymmetric preorder is said to be an *order*. A *total preorder* \preceq on a set X is a preorder such that if $x, y \in X$, then $[x \preceq y]$ or $[y \preceq x]$.

If \preceq is a preorder on X , then as usual we denote the associated *asymmetric* relation by \prec and the associated *equivalence* relation by \sim and these are defined, respectively, by $[x \prec y \iff (x \preceq y) \wedge \neg(y \preceq x)]$ and $[x \sim y \iff (x \preceq y) \wedge (y \preceq x)]$. Also, the associated *dual* preorder \preceq_d is defined by $[x \preceq_d y \iff y \preceq x]$.

Let (X, \preceq) be a totally preordered set and let X/\sim be the set of equivalence classes. If $x \in X$ we denote the equivalence class of x by $[x]$. The preorder \preceq on X induces a natural order \preceq on X/\sim defined by $[x] \preceq [y] \iff x \preceq y$.

Let $[x], [y]$ be two equivalence classes in X/\sim . Then we say that the ordered pair $([x], [y]) \in (X/\sim) \times (X/\sim)$ is a *jump* if there is no $[z] \in X/\sim$ such that $[x] < [z] < [y]$, where $<$ denotes the asymmetric part of \preceq . If $([x], [y])$ is a jump, then we sometimes abuse notation and say that (x, y) is a jump in X . But it should be remembered that a jump is only defined for the corresponding equivalence classes.

A totally preordered set (X, \preceq) is said to be *densely ordered* if it has no jumps. A subset Z of X is said to be *order-dense* in X with respect to \preceq , if

$x, y \in X$ and $x \prec y$ imply that there exists $z \in Z$ such that $x \preceq z \preceq y$. (X, \preceq) is said to be *order-separable* if it has a countable order-dense subset.

A totally preordered set (X, \preceq) is said to be *Dedekind-complete* if each nonempty subset F that has an upper bound has a least upper bound.

Let (X, \preceq) be a totally preordered set. A subset $Z \subseteq X$ is said to be *cointial* (respectively, *cofinal*) in X if for every $x \in X$ there exists some $z \in Z$ such that $z \preceq x$ (respectively, such that $x \preceq z$). The preorder \preceq is said to be *countably bounded* if there exists a countable subset $Z \subseteq X$ that is cointial and cofinal in X .

If (X, \preceq) is a preordered set, then a real-valued function $u : X \rightarrow \mathbb{R}$ is said to be

- (i) *increasing* if for every $x, y \in X$, $[x \preceq y \Rightarrow u(x) \leq u(y)]$,
- (ii) *order-preserving* if f is increasing and $[x \prec y \Rightarrow u(x) < u(y)]$.

An order-preserving function is also said to be an *order-monomorphism*.

If a nonempty set X is endowed with a topology τ , then a *total* preorder \preceq on X is said to be *continuously representable* if there exists an order-monomorphism $u : X \rightarrow \mathbb{R}$ that is continuous.

Let (X, \preceq) be a totally preordered set. The family of all sets of the form $L(x) = \{a \in X : a \prec x\}$ and $G(x) = \{a \in X : x \prec a\}$, where $x \in X$ is a subbasis for a topology τ_{\preceq} on X . The pair (X, τ_{\preceq}) is called the *order topology* on X . The pair (X, τ_{\preceq}) is called a *preordered topological space*.

If (X, \preceq) is a preordered set and τ is a topology on X , then the preorder \preceq is said to be τ -continuous on X if for each $x \in X$ the sets $\{a \in X : x \preceq a\}$ and $\{b \in X : b \preceq x\}$ are τ -closed in X . If \preceq is a total preorder on X , then it is easy to prove that the continuity of \preceq amounts to the fact that the graph of \preceq (i.e., $\{(x, y) \in X \times X; x \preceq y\}$) is a closed subset of $X^2 = X \times X$ endowed with the product topology $\tau_X \times \tau_X$. A topology τ on (X, \preceq) is said to be *natural* if \preceq is τ -continuous. Thus, if \preceq is continuous with respect to a topology τ on X , then $\tau_{\preceq} \subseteq \tau$.

Given a nonempty set X endowed with a topology τ (i.e., (X, τ) is a *topological space*), the topology τ on X is said to have the *continuous representability property* (CRP) if every continuous total preorder \preceq defined on X admits a representation by means of a continuous order-monomorphism. Topologies of this kind were introduced by Herden [25] under the name of “useful topologies” (see also Herden and Pallack [26]). Among the topologies that have CRP are the second countable ones (see Debreu [17]), the connected plus separable ones (see Eilenberg [19]) and the locally connected plus separable ones (see Campión et al. [8]).

Given a topological space (X, τ) , the topology τ is said to be *separably connected* if for every two points $a, b \in X$ there exists a connected and separable subset $C_{a,b} \subseteq X$ that includes a and b , and it is said to be *preorderable* if it is the order topology τ_{\preceq} of some *total preorder* \preceq defined on X .

Remark 2.1. At this point, it is important to explain why this theory of order-representability of topological spaces deals with *total preorders* instead of (just) total orders. The main reason is that on many classical topological spaces (X, τ) there is *no continuous total order* (A well known *example* of this situation is the real plane \mathbb{R}^2 endowed with its usual Euclidean metric and topology, see Theorem 4 in Candeal and Induráin [10] for details).

3. Properties of order topologies

Before studying order-representability properties on a (general) topological space, it seems necessary to begin with a totally preordered set (X, \preceq) endowed with its corresponding order topology τ_{\preceq} , in order to understand better what else must happen for the order topology τ_{\preceq} to have some classical topological property.

Having this idea in mind, we include this *preparatory Section 3*, in which we quote several helpful results on ordered sets, to be used in the sequel. Some proofs have been omitted because they are either straightforward or well-known (see e.g. Bridges and Mehta [3], Th. 1.6.11 and Th. 3.2.9; Campión et al. [5], Lemma 3.9 and Corollary 3.10; Steen and Seebach [35], pp. 67–68).

Lemma 3.1. *Let X be a nonempty set endowed with a total preorder \preceq . Let τ_{\preceq} be the order topology on X . Then the following conditions are equivalent:*

- (i) *The total preorder \preceq is continuously representable through an order-isomorphism.*
- (ii) *The order topology τ_{\preceq} is second countable.*
- (iii) *The totally preordered set (X, \preceq) is order-separable.*

Lemma 3.2. *Let X be a nonempty set endowed with a total order \preceq .*

- (i) *The order topology τ_{\preceq} is connected if and only if (X, \preceq) is densely ordered and Dedekind complete.*
- (ii) *The order topology τ_{\preceq} on X is second countable if and only if it is metrizable and separable.*
- (iii) *If the order topology τ_{\preceq} on X is connected and separable, then it is second countable.*

Remark 3.3. Separability cannot be dropped in part (ii) of Lemma 3.2. A counterexample is $X = (0, 1) \times (0, 1) \subset \mathbb{R}^2$ endowed with the lexicographic order \leq_L defined by $(x, y) \leq_L (z, t) \iff [(x < z) \vee (x = z, y \leq t)]$ for every $0 < x, y, z, t < 1$ (see also Theorem 2 in Candeal and Induráin [12]).

Lemma 3.4. *Let X be a nonempty set endowed with a total order \preceq . Then the order topology τ_{\preceq} on X is path-connected if and only if it is separably connected.*

Proof. See part (iv) of Remark 2 in Candeal et al. [9]. □

Remark 3.5. On a general topological space (X, τ) it is clear that connected plus separable implies separably connected. The converse may fail to be true

(consider, e.g., a non-separable Banach space endowed with the norm topology). It is also true that separably connected implies connected, but, even on metric spaces, the converse is not true in general (see, e.g., Simon [34]). There are also examples of separably connected topological spaces that are not path-connected: Consider, for instance, the subset of \mathbb{R}^2 given by $(\{0\} \times [-1, 1]) \cup \{(x, \sin(\frac{1}{x})) : x > 0\}$.

Lemma 3.6. *Let X be a nonempty set endowed with a total order \preceq that is countably bounded. Suppose that the order topology τ_{\preceq} on X is separably connected. Then τ_{\preceq} is second countable.*

*Proof.*¹ Since X is countably bounded, it can be written as the union of a countable number of nested intervals $[a_n, b_n] \subseteq X$. Since \preceq is continuous and τ_{\preceq} is separably connected each of these intervals is contained in a separable and connected subset of X . That means that X is the union of a countable number of connected separable subsets, all of which intersect, so it follows that the order topology τ_{\preceq} on X is separable and connected. Therefore τ_{\preceq} is second countable by Lemma 3.2(iii). \square

Corollary 3.7. *Let X be a nonempty set endowed with a topology τ such that (X, τ) is separably connected. Let \preceq be a countably bounded and τ -continuous total preorder defined on X . Then \preceq is continuously representable through an order-monomorphism.*

Proof. Since the topological space (X, τ) is separably connected, the order topology τ_{\preceq} , that satisfies $\tau_{\preceq} \subseteq \tau$ by hypothesis, is also separably connected. It is obvious that the total order \preceq on the quotient space through indifference X/\sim is also countably bounded, and its corresponding order topology is separably connected. Thus, as a direct consequence of Lemma 3.6 and Lemma 3.1, it follows that \preceq is continuously representable (considering the order topology τ_{\preceq} on X/\sim and the usual topology on \mathbb{R}) by means of an order-monomorphism. Consequently, \preceq is continuously representable (now considering the topology τ_{\preceq} on X and the usual topology on \mathbb{R}) through an order-monomorphism ϕ . Indeed, since $\tau_{\preceq} \subseteq \tau$, the map ϕ is also continuous if we consider on X the given topology τ and on \mathbb{R} the usual topology. \square

Theorem 3.8. *Let X be a nonempty set endowed with a total order \preceq . Suppose that the order topology τ_{\preceq} on X is separably connected. Then the following conditions are equivalent:*

- (i) \preceq is countably bounded.
- (ii) \preceq is continuously representable.
- (iii) τ_{\preceq} is second countable.
- (iv) τ_{\preceq} is metrizable and second countable.
- (v) τ_{\preceq} is metrizable and separable.
- (vi) τ_{\preceq} is separable.

¹We owe this proof of Lemma 3.6 to an anonymous referee.

- (vii) τ_{\preceq} is Lindelöf.
- (viii) τ_{\preceq} satisfies the countable chain condition (CCC).
- (ix) τ_{\preceq} is σ -compact.

Proof. We follow the scheme (i) \Rightarrow (iii); (ii) \iff (iii) \iff (iv) \iff (v); (v) \Rightarrow (vi) \Rightarrow (ii); (iii) \Rightarrow (vii) \Rightarrow (i); (iii) \Rightarrow (viii) \Rightarrow (i) \Rightarrow (ix) \Rightarrow (vii).

The fact (i) \Rightarrow (iii) has been proved in Lemma 3.6.

The equivalence (ii) \iff (iii) was stated in Lemma 3.1. The order topology τ_{\preceq} is connected because, by hypothesis, it is separably connected.

The equivalences (iii) \iff (iv) \iff (v) as well as the implication (v) \Rightarrow (vi) follow now immediately from Lemma 3.2(ii)-(iii) and Th. 5.6 in Dugundji [18].

Assume now that (vi) holds. Let $D \subseteq X$ be a countable subset that meets every nonempty τ_{\preceq} -open subset. Let $a, b \in X$ be such that $a \prec b$. Since the topology τ_{\preceq} is connected by hypothesis, it follows by Lemma 3.2(i) that (X, \preceq) is densely ordered. Hence there exists $c \in X$ such that $a \prec c \prec b$, so that in particular the τ_{\preceq} -open subset $(a, b) = \{x \in X : a \prec x \prec b\}$ is nonempty. Since D meets every nonempty τ_{\preceq} -open subset, there also exists $d \in D$ such that $a \prec d \prec b$. In particular, we have $a \preceq d \preceq b$. Hence D is order-dense in X . Condition (ii) follows now from Lemma 3.1.

The fact (iii) \Rightarrow (vii) is immediate.

Assume now that (vii) holds. If X has both a first element a and a last element b , then it is plain that \preceq is countably bounded. If X is unbounded, assuming that X has no last element with respect to \preceq , it happens that the family \mathcal{L} of all sets of the form $L(x) = \{a \in X : a \prec x\}$ is a τ_{\preceq} -open covering of X . By hypothesis, there exists a countable subcovering $\{L(x_n) : n \in \mathbb{N}\}$ of \mathcal{L} . Similarly, if X has no first element with respect to \preceq , then the family \mathcal{G} of all sets of the form $G(y) = \{b \in X : y \prec b\}$ is a τ_{\preceq} -open covering of X , and there exists a countable subcovering $\{G(y_k) : k \in \mathbb{N}\}$ of \mathcal{G} . Now it is straightforward to see that the countable subset $D = \{x_n : n \in \mathbb{N}\} \cup \{y_k : k \in \mathbb{N}\} \cup \{a, b\} \subseteq X$, where a (respectively, b) denotes the first element if any (respectively, the last element if any) of X , is coinital and cofinal in X with respect to \preceq . Therefore, \preceq is countably bounded, so that we get (i).

The fact (iii) \Rightarrow (viii) is clear because every second countable space is separable, and every separable space satisfies the countable chain condition (see e.g. Corollary 2.3.18 in Engelking [20]).

Assume now that (viii) holds. Suppose that there is no countable subset $D \subset X$ that is cofinal in X . Then (X, \preceq) contains a subset Y such that (Y, \preceq) can be identified to the first uncountable ordinal ω_1 endowed with the ordinal inclusion \leq . Thus, there exists a bijective map $F : \omega_1 \rightarrow Y$ such that $z \preceq t \iff F^{-1}(z) \leq F^{-1}(t)$ ($z, t \in Y$). For every $\alpha < \omega_1$, let $\alpha + 1$ be the ordinal that follows α . Since the order topology τ_{\preceq} on X is separably connected, hence connected, the collection $\{(F(\alpha), F(\alpha+1)) : \alpha < \omega_1\}$ would then violate CCC, which is a contradiction. Therefore, there exist countable

subsets $C, E \subseteq X$ such that C is coinitial in X and E is cofinal in X with respect to \preceq . The union $D = C \cup E$ is countable, coinitial and cofinal in X . Thus, \preceq is countably bounded, so that we get (i) again.

Assume now that (i) holds. Let $D = \{x_n : n \in \mathbb{N}\}$ be a coinitial and cofinal subset of X . For every $k \neq l \in \mathbb{N}$ such that $x_k \prec x_l$, we observe that the subset $[x_k, x_l] = \{y \in X : x_k \preceq y \preceq x_l\}$ is compact in the order topology τ_{\preceq} of X , by Lemma 3.2(i) and Theorem 27.1 in [30]. Therefore $X = \bigcup_{\{k, l \in \mathbb{N}, k \neq l\}} [x_k, x_l]$ is obviously σ -compact. Thus we get (ix). Moreover, since a σ -compact topological space is in particular Lindelöf we also get (vii). \square

Remarks 3.9. (i) New conditions could be added to the statement of Theorem 3.8, as a consequence of some results in Sections 2 and 3 of Lutzer and Bennet [29]. These results state that:

Let X be a nonempty set endowed with a total order \preceq . Consider on X the order topology τ_{\preceq} . Then it holds that

- a) the order topology τ_{\preceq} satisfies CCC if and only if it is hereditarily Lindelöf,
- b) the order topology τ_{\preceq} is separable if and only if it is hereditarily separable.

Notice in addition that in this result it is *not necessary* to ask the order topology τ_{\preceq} to be *separably connected*.

(ii) The hypothesis of being separably connected that appears in the statement of Theorem 3.8 cannot be replaced by the weaker one of connectedness. An *example* is the lexicographically ordered set $L = \{[0, \omega_1) \times J\} \cup \{(\omega_1, 0)\}$ where $J = \{x \in \mathbb{R} : 0 \leq x < 1\}$ and $\{[0, \omega_1)$ denotes the *long line* (see, e.g., Steen and Seebach [35]).

4. The continuous representability property on topological spaces

In the present section we search for topological conditions on a topological space (X, τ) , in order for τ to satisfy the continuous representability property (CRP). We pay an special attention to conditions related to covering properties.

The results of Section 3 are decisive in this process, due to the following *key fact* stated in the next Theorem 4.1, whose proof appears in Campión et al. [6].

Theorem 4.1. *Let (X, τ) be a topological space. Then the topology τ satisfies CRP if and only if all its preorderable subtopologies are second countable.*

A straightforward strengthening of Lemma 3.2(ii) shows that, given a topological space (X, τ) , any preorderable subtopology is second countable if and only if it is separable and pseudometrizable. This fact immediately leads to the following corollary.

Corollary 4.2. *Let (X, τ) be a topological space. Then the topology τ satisfies CRP if and only if all its preorderable subtopologies are separable and pseudometrizable.*

Preorderable topologies were characterized in Campión et al. [6] completing the panorama on *orderability of topologies* (see Van Dalen and Wattel [36], Purisch [31] or Gutev [24]).

Suppose now that we are looking for conditions on a topological space (X, τ) that imply that the topology τ satisfies CRP. *Which kind of conditions should we analyze first?*

A glance to Theorem 4.1 gives us an idea to begin with: Any topological condition *that implies second countability and is inherited by subtopologies* fits well our purposes. We can say even more, observing that *any topological property that implies that every preorderable subtopology is second countable* also implies CRP.

Now we obtain important results about the property CRP on topological spaces.

Theorem 4.3. *Let (X, τ) be a topological space. Each of the following conditions implies that (X, τ) satisfies CRP:*

- (i) τ is connected and separable (see Eilenberg [19]),
- (ii) τ is separably connected and satisfies CCC (see Campión et al. [4]),
- (iii) τ is separably connected and compact,
- (iv) τ is separably connected and σ -compact,
- (v) τ is separably connected and satisfies the Lindelöf property,
- (vi) τ is second countable (see Debreu [17]),
- (vii) τ is locally connected and separable (see Candeal et al. [15]).

Proof. Some of these results are already known in the literature. However, we offer here an alternative proof for some of them (parts (i) to (v)) that follows from the previous results of Section 3 and the observation above. Thus, notice that if τ is connected and separable, then every preorderable subtopology is also connected and separable. Hence, by Lemma 3.2(iii), the preorderable subtopologies of τ are indeed second countable. In the same way, if τ is separably connected, then every preorderable subtopology of τ is also separably connected. If, in addition, τ satisfies CCC, or the Lindelöf property, or it is compact or σ -compact, the same happens for every preorderable subtopology of τ . In each of these situations the preorderable subtopologies are actually second countable, as a consequence of Theorem 3.8.

By the way, part (i) is a well-known classical result in the theory of ordered structures (see Eilenberg [19]). The same happens to part (vi), that is known as *Debreu's theorem* (see Debreu [17] or Bridges and Mehta [3], Ch. 3). Finally, part (ii) also appears in Campión et al. [4], and a proof of part (vii) can be seen in Candeal et al. [14]. \square

Remark 4.4. Part (ii) of the above Theorem 4.3 shows that in separably connected topological spaces CCC implies CRP. However, the converse is *not* true. An example is the *Alexandroff topology* on $[0, 1] \times [0, 1]$ (see Steen and Seebach [35], pp. 120–121 for details) that is compact and path-connected, so that it is separably connected, and, by part (iii) of Theorem 4.3, it satisfies CRP. Moreover, it does not satisfy CCC and fails to be first countable.

In the category of *metric spaces* the next Theorem 4.5, and its subsequent Corollary 4.6, provide not only sufficient conditions, but actually *characterizations* of the fulfillment of the continuous representability property CRP. As a matter of fact, Theorem 4.5 is well-known in this literature (see e.g. Estévez and Hervés [21] or Candeal et al. [9]).

Theorem 4.5. *Let (X, d) be a metric space. Then the metric topology τ_d satisfies CRP if and only if it is separable.*

Corollary 4.6. (i) *Let (G, τ) be a first countable topological group. Then the topology τ satisfies CRP if and only if it is separable.*

(ii) *Let (G, τ) be a locally compact topological group. Let $C_0(G)$ be the space of continuous complex functions defined on G which vanish at infinity. Suppose that the weak topology on $C_0(G)$ satisfies the Lindelöf property. Then the topology τ satisfies CRP if and only if it is separable.*

(iii) *Let (G, τ) be a locally compact topological group such that the weak topology on $C_0(G)$ is normal. Then the topology τ satisfies CRP if and only if it is separable.*

(iv) *In a separably connected metric space CCC and CRP are equivalent conditions.*

Proof. (i). The classical Birkhoff-Kakutani theorem (see Birkhoff [2], Kakutani [27] or Kelley [28], p. 186) states that on a topological group (G, τ) the topology τ is first countable if and only if it is metrizable. The result follows now by Theorem 4.5.

(ii) and (iii). This also follows from Theorem 4.5, making use of Theorem 2 in Corson [16] where it is proved that on a locally compact topological group (G, τ) the following conditions are equivalent:

- (a) τ is a metrizable topology on G ,
- (b) the weak topology on $C_0(G)$ satisfies the Lindelöf property,
- (c) the weak topology on $C_0(G)$ is normal.

(iv). It is a consequence of Theorem 4.3(ii) and Theorem 4.5, since separability implies CCC. \square

Remarks 4.7. (i) We cannot drop “first countable” in the statement of Corollary 4.6. In other words, for a topological group (G, τ) it is *not true*, in general, that CRP is equivalent to separability. To cite an example, consider an infinite dimensional nonseparable Banach space X (e.g. the Hilbert space of uncountable basis $\ell_2(\mathbb{R})$) endowed with the weak topology ω . Observe that (X, ω) is, in particular, a topological group. The topology ω is not first countable since otherwise X would be finite dimensional (see e.g. Theorem 6.30 in Aliprantis and Border [1]). On the other hand, ω is not separable because on a Banach space, and as an easy consequence of Hahn-Banach theorem, the separability of the norm topology is equivalent to the separability of the weak topology ω . Finally, notice that by Theorem 4.3(ii) ω satisfies CRP, since it satisfies CCC, as proved in Campión et al. [4].

(ii) It could be interesting to say that there are classical topologies on remarkable families of topological spaces that do satisfy CRP. Sometimes this information is easily obtained from the results just introduced in this Section 4. Observe, for instance, that in the category of *topological vector spaces* any compatible topology is path-connected, hence separably connected. This allows us to use Theorem 4.3 to obtain particular results about the fulfillment of CRP in those spaces.

An important particular case is that of *Banach spaces*. Thus, if X is a real Banach space, it can be proved that the norm topology $\tau_{\|\cdot\|}$ on X satisfies CRP if and only if it is separable, as a direct consequence of Theorem 4.5. In addition, the weak topology ω on X always satisfies CRP (see Campión et al. [4]). Hence any topology τ on X such that $\tau \subseteq \omega$ satisfies CRP, and also the weak star topology ω^* on the Banach space X^* (the topological dual of X) satisfies CRP. Finally, in Candeal et al. [13], pp. 57–59, it has been proved that if the norm topology $\tau_{\|\cdot\|}$ on X is separable, then any topology τ on the dual Banach space X^* such that $\tau \subseteq m(X^*, X)$ satisfies CRP, where $m(X^*, X)$ stands here for the Mackey topology that the dual pair (X^*, X) induces on X^* .

We conclude this section with a useful result concerning the behaviour of CRP through continuous surjections of topological spaces.

Theorem 4.8. *CRP is invariant under continuous surjections.*

Proof. Let (X, τ_X) be a topological space such that τ_X satisfies CRP. Let (Y, τ_Y) be a topological space such that there exists a continuous surjection $\pi : X \rightarrow Y$. We observe that a total preorder \lesssim_Y on Y induces a total preorder \lesssim_X on X by declaring that $x_1 \lesssim_X x_2 \iff \pi(x_1) \lesssim_Y \pi(x_2)$. It is clear that, if the preorder \lesssim_Y is τ_Y -continuous, then the corresponding preorder \lesssim_X on X is τ_X -continuous, as a consequence of the continuity of the projection π : Given an element $x \in X$, it follows that $\{z \in X : z \prec_X x\} = \{z \in X : \pi(z) \prec_Y \pi(x)\} = \pi^{-1}(\{s \in Y : s \prec_Y \pi(x)\})$. Similarly $\{t \in X : x \prec_X t\} = \{t \in X : \pi(x) \prec_Y \pi(t)\} = \pi^{-1}(\{r \in Y : \pi(x) \prec_Y r\})$. In addition, since π is a surjection we have that given an element $y \in Y$ there exists at least one element $x_y \in X$ such that $\pi(x_y) = y$. Moreover $\pi(\{z \in X : z \prec_X x_y\}) = \{s \in Y : s \prec_Y y\}$ and $\pi(\{t \in X : x_y \prec_X t\}) = \{r \in Y : y \prec_Y r\}$. Therefore, if we consider on X the order topology induced by \lesssim_X and on Y the order topology induced by \lesssim_Y , then with respect to these topologies the map π is not only a continuous surjection, but also an *open* map. Since τ_X satisfies CRP, the order topology that \lesssim_X induces on X is second countable by Theorem 4.1. Hence the order topology that \lesssim_Y induces on Y is also second countable, because second countability is indeed invariant under continuous *open* surjections (see e.g. Dugundji [18], Th. 6.2 on p. 174). Again by Theorem 4.1, we conclude that τ_Y satisfies CRP. \square

Corollary 4.9. (i) *CRP is invariant under topological quotients.*

(ii) If the product topology τ_p of a product $\prod_{i \in I} (X_i, \tau_i)$ of topological spaces satisfies CRP, then the topology τ_i ($i \in I$) of each factor also satisfies CRP.

Remark 4.10. The converse of part (ii) of Corollary 4.9 is an *open question*.

In this direction, we say that a property \mathcal{P} on topological spaces is said to be *multiplicative* if for any topological spaces (X, τ_X) and (Y, τ_Y) that satisfy \mathcal{P} , the topological space $(X \times Y, \tau_X \times \tau_Y)$ also satisfies \mathcal{P} . Similarly, \mathcal{P} is said to be a *square property* if for any topological space (X, τ_X) that satisfies \mathcal{P} , the square topological space $(X \times X, \tau_X \times \tau_X)$ also satisfies \mathcal{P} . Of course, multiplicative properties are in particular square properties but the converse is not true, in general. Now, the *open problem* is the following:

- (i) Is CRP a multiplicative property?
- (ii) Is CRP a square property?

Of course, there are *partial answers* to the aforementioned question. For instance, as a consequence of Theorem 4.5, on *metric spaces* CRP is a multiplicative property because separability and metrizability are indeed multiplicative properties. Related to the open problem just introduced we may *conjecture* that CRP is a multiplicative property on *separably connected* topological spaces.

5. The continuous algebraic representability property

In this section, we explore the algebraic version of CRP. In the topological setting, CRP appears whenever we look for ordinal representations of total preorders that, in addition, preserve a nice topological property: namely, the continuity. In the algebraic context, in addition to the continuity property for an order-preserving function, we will ask for a new demanding requirement: that of being an algebraic-homomorphism. Of course, this imposes some kind of compatibility among order, topology, and algebra involved. Although we could begin with a very simple algebraic ordered structure (e.g., think of a totally preordered semigroup), we will focus on richer algebraic systems. In particular, we will pay attention to *totally preordered real algebras*. The reason for this choice will be clarified later on in Section 6 when all of this algebraic machinery will be applied to characterize *strongly dictatorial social welfare functionals*, an important issue within social choice theory. We now introduce some basic definitions.

Definition 5.1. A *real algebra* $(X, +, \cdot_{\mathbb{R}}, *)$ is a set X endowed with three binary operations so that:

- (i) $(X, +, \cdot_{\mathbb{R}})$ is a real vector space.
- (ii) $(X, +, *)$ is a ring.
- (iii) $\lambda \cdot (x * y) = (\lambda \cdot x) * y = x * (\lambda \cdot y)$ for all $x, y \in X, \lambda \in \mathbb{R}$.

Notation. The null element of X with respect to “+” will be denoted by $\mathbf{0}$.

Let $(X, +, \cdot_{\mathbb{R}}, *)$, $(\bar{X}, \bar{+}, \bar{\cdot}_{\mathbb{R}}, \bar{*})$ be two real algebras. A function $v : X \rightarrow \bar{X}$ is said to be an *algebra-homomorphism* if the following three conditions are met:

- (a) $v(x + y) = v(x) \bar{+} v(y)$ for all $x, y \in X$ (i.e., it is *additive*),
- (b) $v(\lambda \cdot x) = \lambda \bar{\cdot} v(x)$ for all $x \in X, \lambda \in \mathbb{R}$ (i.e., it is *homogeneous*),
- (c) $v(x * y) = v(x) \bar{*} v(y)$ for all $x, y \in X$ (i.e., it is *multiplicative*).

Remark 5.2. Condition (a) in combination to condition (b) can be given in a single formula in the following terms:

$v(\lambda \cdot x + \beta \cdot y) = \lambda \bar{\cdot} v(x) \bar{+} \beta \bar{\cdot} v(y)$ for all $x, y \in X, \lambda, \beta \in \mathbb{R}$ (i.e., it is *linear*).

In the next definition, some usual compatibility conditions between order and algebra are introduced. This leads to the concept of a *totally preordered real algebra*, or a *totally ordered real algebra* if the preorder is an order.

Definition 5.3. A *totally preordered real algebra* $(X, \preceq, +, \cdot_{\mathbb{R}}, *)$ is a real algebra equipped with a total preorder \preceq which is compatible with the operations “+”, “ $\cdot_{\mathbb{R}}$ ” and “*”, i.e., if the following three conditions are met:

- (1) $x \preceq y$ implies $x + z \preceq y + z$ for all $z \in X$ (*translation-invariance*),
- (2) $x \preceq y, 0 \leq \lambda$ imply $\lambda \cdot x \preceq \lambda \cdot y$ (*homotheticity*),
- (3) $x \preceq y, \mathbf{0} \preceq z$ imply $z * x \preceq z * y$ and $x * z \preceq y * z$ (*multiplicative-invariance*).

Notation. The class of all total preorders defined on a real algebra X that satisfy the conditions (1) to (3) of Definition 5.3 above will be denoted by \mathcal{P}_a , i.e., $\mathcal{P}_a = \{\preceq : \preceq \text{ is a translation-invariant, homothetic and multiplicative-invariant total preorder defined on } X\}$.

Remark 5.4. An easy example of a totally preordered real algebra is the n -dimensional Euclidean space $X = \mathbb{R}^n, n \geq 1$, endowed with the usual binary operations, “+”, “ $\cdot_{\mathbb{R}}$ ” and “*”, defined componentwise (i.e., $(a_i) + (b_i) = (a_i + b_i), \lambda \cdot (a_i) = (\lambda a_i), (a_i) * (b_i) = (a_i b_i)$ for all $(a_i), (b_i) \in \mathbb{R}^n, \lambda \in \mathbb{R}$), and the total preorder defined as follows: Let $j \in \{1, \dots, n\}$ be fixed. Then, for every $(x_i), (y_i) \in \mathbb{R}^n$, define $(x_i) \preceq (y_i)$ if and only if $x_j \leq y_j$. This total preorder only pays attention to the j -th coordinate of the corresponding vectors. This is the reason why we call it a *projective* total preorder on \mathbb{R}^n . This example can be straightforwardly generalized to include infinite-dimensional sequence spaces like $X = c_0$ or $X = l_1$, standard notations which correspond to the Banach space of real sequences vanishing at infinity and the Banach space of absolutely sumable real sequences, respectively.

An obvious example of a non-projective total preorder \preceq defined on $X = \mathbb{R}^n$ which satisfies the conditions (1) to (3) of Definition 5.3 is the trivial one (i.e., $(x_i) \sim (y_i)$ for every $(x_i), (y_i) \in \mathbb{R}^n$). Notice that this particular binary relation belongs to \mathcal{P}_a for any real algebra X . We will use the notation \preceq_t to refer to this total preorder (i.e., $x \sim_t y$ for all $x, y \in X$). Now, we present a much

more sophisticated example of a total preorder defined on a sequence space that belongs to \mathcal{P}_a and is non-projective. Consider the space $X = l_\infty$, which consists of all bounded real sequences. Endowed with the usual operations defined coordinatewise l_∞ is a real algebra. Let us denote by $\beta(\mathbb{N})$ the Stone-Ćech compactification of the set of the natural numbers \mathbb{N} (see, e.g., Dugundji [18]). By the Stone-Ćech's theorem every $x \in l_\infty$ extends in a unique manner to a continuous real-valued function, say, \tilde{x} , defined on $\beta(\mathbb{N})$. Let $p \in \beta(\mathbb{N}) \setminus \mathbb{N}$ be fixed. Then, it can be easily proved that the relation defined as $x \lesssim_p y$ if and only if $\tilde{x}(p) \leq \tilde{y}(p)$, ($x, y \in l_\infty$), is a total preorder on l_∞ that makes it to be a totally preordered real algebra. Clearly, \lesssim_p , so defined, is not projective.

Next we define the notion of a *topological real algebra*.

Definition 5.5. A real algebra $(X, +, \cdot_{\mathbb{R}}, *)$ equipped with a topology τ is said to be a *topological real algebra* if $(X, \tau, +, \cdot_{\mathbb{R}})$ is a topological vector space and $*$ is a continuous binary operation on $X \times X$.

Remark 5.6. Examples of topological real algebras are \mathbb{R}^n , $n \geq 1$, with the usual binary operations mentioned above and the Euclidean topology. Also, they are c_0 , l_∞ and l_1 endowed with the topology given by the usual norms; i.e., $\|(x_n)\|_\infty = \sup\{|x_n| : n \in \mathbb{N}\}$ for any $(x_n) \in c_0$ or l_∞ , and $\|(x_n)\|_1 = \sum |x_n|$, for any $(x_n) \in l_1$. Actually, all of these are examples of richer mathematical structures; namely, they are Banach algebras (see, e.g., Rickart [32] for an excellent account on this topic).

Notation. In order to shorten the notation, we will simply write X to denote a real algebra. If there is no ambiguity, we will keep the same notation throughout this section even though X is a totally preordered real algebra or a totally preordered topological real algebra.

Before introducing the concept of CARP in this algebraic environment a new notion is needed. A total preorder \lesssim on X is said to be *zero* if $x * y \sim \mathbf{0}$ for all $x, y \in X$. Otherwise \lesssim is said to be *non-zero*. If \lesssim is a total order, then \sim is the equality relation $=$. In this case, since the previous condition reduces to $x * y = \mathbf{0}$ for all $x, y \in X$, we will say that X is *zero* or *non-zero*, respectively.

Definition 5.7. Let X be a real algebra and let τ be a topology on X . Then τ satisfies the *continuous algebraic representability property* (shortly, CARP) if every continuous, translation-invariant, homothetic, multiplicative-invariant and non-zero total preorder \lesssim defined on X admits a continuous order-preserving function which is an algebra-homomorphism (shortly, a continuous algebraic order-preserving function).

Remarks 5.8. (i) It should be observed that the only zero total preorder defined on a real algebra X that admits an algebraic order-preserving function is the trivial one \lesssim_t (i.e., $x \sim_t y$ for every $x, y \in X$) for which the algebraic order-preserving function turns out to be the null function.

(ii) The condition of being non-zero plays an important role in Definition 5.7. Indeed, by Remark 5.8(i) and from the point of view of the existence of an algebraic order-preserving function, zero and non-trivial total preorders on a real algebra X are pathological. It is remarkable that this kind of total preorders can exist even though X is endowed with a nice topology. This is the reason why they are not included in Definition 5.7. Let us see an example of a topological totally preordered real algebra $(X, \tau, \preceq, +, \cdot_{\mathbb{R}}, *)$ such that \preceq is zero, continuous and non-trivial. Consider $X = \mathbb{R}^2$, endowed with the usual Euclidean topology and the usual binary operations, “+” and “ $\cdot_{\mathbb{R}}$ ” defined componentwise. Then, define both the binary operation “*” and the binary relation \preceq as follows: $(x_1, x_2) * (y_1, y_2) = (x_1 y_1, 0)$; $(x_1, x_2) \preceq (y_1, y_2)$ if and only if $x_2 \leq y_2$. It is simple to see that X , with “*” so defined, is a topological real algebra. In addition, it can be easily shown that \preceq , so defined, is a continuous, zero and non-trivial total preorder on X which belongs to \mathcal{P}_a . Notice that, although it admits a linear order-preserving function (e.g., $\psi(x_1, x_2) = x_2$ for every $(x_1, x_2) \in \mathbb{R}^2$, is such a function), there is no such a function which is multiplicative.

(iii) Projective preorders on \mathbb{R}^n , or on some sequence spaces like c_0 or l_1 , have the interesting property that they are the only non-trivial total preorders that admit a continuous algebraic order-preserving function (see, Remark 5.15 below). However, this is not the case for any arbitrary totally preordered topological real algebra. Indeed, consider again the Banach algebra $X = l_{\infty}$ and let $p \in \beta(\mathbb{N}) \setminus \mathbb{N}$ be fixed (see the second paragraph of Remark 5.4 above). Then, \preceq_p is a non-trivial total preorder on l_{∞} that is not projective. Observe that the evaluation map at p , $x \in l_{\infty} \rightsquigarrow e_p(x) = \tilde{x}(p) \in \mathbb{R}$, is a continuous algebraic order-preserving function for \preceq_p .

Notation. Similarly to the notation used above, we now introduce two important sub-classes of binary relations defined on X . These are, respectively, $\mathcal{P} = \{\preceq : \preceq \text{ is a continuous, translation-invariant, homothetic and multiplicative-invariant total preorder defined on } X\}$ and $\tilde{\mathcal{P}} = \{\preceq : \preceq \text{ is a non-zero, continuous, translation-invariant, homothetic and multiplicative-invariant total preorder defined on } X\}$. Obviously, $\tilde{\mathcal{P}} \subseteq \mathcal{P}$.

Next lemma will be useful to prove the main result of this section.

Lemma 5.9. *Let X be a non-zero² topological real algebra and let $\preceq \in \mathcal{P}$ a total order defined on X . Then there is a continuous algebraic order-preserving function for \preceq . Moreover, this function is unique, onto and open.*

Proof. First of all, we are going to show that the order topology τ_{\preceq} coincides with τ on X . Notice that, with respect to τ , X is a Hausdorff space since, by continuity of \preceq , $\tau_{\preceq} \subseteq \tau$ and clearly X is a Hausdorff space with respect

²We are indebted to an anonymous referee for drawing our attention to include this possibility.

to τ_{\preceq} . Thus, $(X, \tau, +, \cdot_{\mathbb{R}})$ is a Hausdorff topological vector space. Since \preceq is a continuous, translation-invariant and homothetic total (pre-)order on X , by a result of Candeal and Induráin [11], there is a linear and continuous order-preserving function, say ψ for \preceq . Notice also that, since \preceq is a total order on X , in addition to be both an algebraic and an order isomorphism, ψ is a homeomorphism from X onto the reals. As a direct consequence $(X, +, \cdot_{\mathbb{R}})$ can be identified with $(\mathbb{R}, +, \cdot_{\mathbb{R}})$, τ with the usual Euclidean topology on \mathbb{R} and \preceq with the usual order \leq on \mathbb{R} or its dual \leq_d (i.e., $a \leq_d b \Leftrightarrow b \leq a$ for every $a, b \in \mathbb{R}$). In particular, the order topology on X , τ_{\preceq} , coincides with τ .

Now, observe that $(X, \preceq, +, *)$ is a non-zero totally ordered ring³ too. In addition, by the argument of the previous paragraph, \preceq is Archimedean (i.e., for any $x, y \in X$, such that $\mathbf{0} \prec x \prec y$, there is $n \in \mathbb{N}$ such that $y \prec nx$). So, by a result of Pickert and Hion (see Fuchs [23], p. 126), there is an order-preserving function, say ϕ , for \preceq which is additive (i.e., $\phi(x + y) = \phi(x) + \phi(y)$ for every $x, y \in X$) and multiplicative. Let us see that ϕ meets all of the properties given in the statement of Lemma 5.9. In particular, it remains to prove that, in addition to satisfy the functional equation $\phi(\lambda \cdot x) = \lambda\phi(x)$ ($\lambda \in \mathbb{R}, x \in X$), ϕ is continuous, onto, open and unique.

(1) ϕ is continuous: Indeed, by the argument of the first paragraph above, (X, \preceq) can be identified with (\mathbb{R}, \leq) or (\mathbb{R}, \leq_d) and $\tau = \tau_{\preceq}$ with the Euclidean topology on \mathbb{R} . Suppose that the first occurs, the other situation being entirely similar. Then ϕ turns out to be an increasing function from \mathbb{R} into \mathbb{R} (with the usual meaning of an increasing real-valued function of one real variable). Now, it is well-known that an increasing function from \mathbb{R} into \mathbb{R} is almost everywhere continuous and, in addition, if it is additive, then it is continuous at every point. Therefore, ϕ is continuous.

(2) ϕ satisfies the functional equation $\phi(\lambda \cdot x) = \lambda\phi(x)$ ($\lambda \in \mathbb{R}, x \in X$): Indeed, a standard reasoning proves that, since ϕ is additive and continuous, it satisfies this homogeneity equation.

(3) ϕ is onto: It follows from linearity.

(4) ϕ is open: To show openness let us observe that, since ϕ is order-preserving, the image of every open \preceq -interval of X is an open \leq -interval of \mathbb{R} . Now, since $\tau = \tau_{\preceq}$ and τ_{\preceq} is the Euclidean topology on \mathbb{R} , it directly follows that ϕ is open.

(5) ϕ is unique: To prove this, we use the fact that ϕ is multiplicative. First, it should be noted that X has an identity, say 1_X (i.e., there is an element, say $1_X \in X$, such that $1_X * x = x * 1_X = x$ for every $x \in X$). Indeed, $1_X = \phi^{-1}(1)$ is the identity. In addition, since $(X, +, \cdot_{\mathbb{R}})$ is a one-dimensional vector space, for every $x \in X$ there is $\lambda_x \in \mathbb{R}$; $x = \lambda_x \cdot 1_X$. Suppose that there is another function $\tilde{\phi}$ that meets the conditions of Lemma 5.9. Let us see that $\tilde{\phi} \equiv \phi$. Indeed, note that, since $\tilde{\phi}$ is multiplicative too, $\tilde{\phi}(1_X) = \phi(1_X) = 1$. Then,

³A totally (pre-)ordered ring is a ring endowed with a (left and right) translation-invariant and multiplicative-invariant total (pre-)order.

by linearity, we have that $\tilde{\phi}(x) = \tilde{\phi}(\lambda_x \cdot 1_X) = \lambda_x \tilde{\phi}(1_X) = \lambda_x = \lambda_x \phi(1_X) = \phi(\lambda_x \cdot 1_X) = \phi(x)$ for every $x \in X$. So, $\tilde{\phi} \equiv \phi$. \square

Remark 5.10. Consider \mathbb{R} with the usual binary operations, the usual order \leq and the Euclidean topology. As direct consequences of Lemma 5.9 the following results are obtained:

(i) The only (up to a unique order-preserving, homeomorphism and algebraic isomorphism) non-zero totally ordered topological real algebra is \mathbb{R} . In particular, a non-zero totally ordered topological real algebra is a commutative field.

(ii) Let $(X, \preceq, +, \cdot_{\mathbb{R}}, *)$ be a non-zero totally ordered real algebra. Then, τ_{\preceq} is the only topology on X that makes it to be a non-zero totally ordered topological real algebra.

The main result of this section will be a characterization of the continuous algebraic representability property (CARP); but, this result must be prefaced by a remark.

Remark 5.11. The following well-known facts will be used in the sequel. They take part of the folklore of the theory of topological vector spaces and topological algebras (see, e.g., Schaefer [33] and Rickart [32]).

Let X be a real vector space and let $S \subseteq X$ be a linear subspace of X . Consider the equivalence relation R_S on X defined as follows: $xR_S y$ if and only if $x - y \in S$. Then, the binary operations “+” and “ $\cdot_{\mathbb{R}}$ ” on X are stable under R_S and so the quotient space X/S is a real vector space too (the corresponding binary operations on this quotient space are denoted also by “+” and “ $\cdot_{\mathbb{R}}$ ”). If, in addition, X is a topological vector space, with topology τ , then the quotient space X/S is also a topological vector space endowed with the quotient topology, denoted by τ_q . Notice that X/S is Hausdorff if and only if S is closed. Consider the projection map, denoted by p , defined as follows: $x \in X \rightsquigarrow p(x) = [x] \in X/S$. It is well-known that p is linear, continuous and open (with respect to the topology τ in X and τ_q in X/S).

Moreover, if X is a real algebra and $S \subseteq X$ is, in addition to being a linear subspace, an ideal⁴ of X , then the operation “ $*$ ” is also stable under R_S which means that the quotient space X/S is a real algebra (the binary operation of the quotient will also be denoted by “ $*$ ”). If X is a topological real algebra, with topology τ , and $S \subseteq X$ is a linear subspace and an ideal of X , then X/S is also a topological real algebra with respect to the quotient topology τ_q . Notice that, in this algebraic context, the projection map p is also multiplicative.

Theorem 5.12. *Let X be a real algebra and let τ be a topology on X . Then the following assertions are equivalent:*

- (i) τ satisfies CARP.

⁴An additive subgroup I of a ring $(A, +, *)$ is said to be an ideal if $x * a \in I$, $a * x \in I$ for every $x \in I$, $a \in X$.

- (ii) *Endowed with the order topology τ_{\lesssim} , X is a topological real algebra for every $\lesssim \in \tilde{\mathcal{P}}$.*

Proof. For the proof of this result, given a total preorder \lesssim on X , it will be useful to consider the following set $I(\mathbf{0}) = \{x \in X : x \sim \mathbf{0}\}$.

(i) \implies (ii). Let $\lesssim \in \tilde{\mathcal{P}}$. Let us see that, endowed with the order topology τ_{\lesssim} , X is a topological real algebra. Since, by hypothesis, τ satisfies CARP there is a continuous algebraic order-preserving function, say $\phi : X \rightarrow \mathbb{R}$ for \lesssim . Notice that $\phi \neq 0$ since \lesssim is non-zero. Then, it is routine to check that $I(\mathbf{0}) = \phi^{-1}(0)$ is a real vector subspace of X which, in addition, is an ideal of X . Consider now the quotient space $X/I(\mathbf{0})$. It is simple to see that, actually, $X/I(\mathbf{0})$ is a totally ordered real algebra with the usual binary operations induced by “+”, “ $\cdot_{\mathbb{R}}$ ” and “ $*$ ” in the quotient $X/I(\mathbf{0})$, and the total order \preceq defined as $[x] \preceq [y] \Leftrightarrow x \lesssim y$. It should be noted that $X/I(\mathbf{0})$ is non-zero since \lesssim is non-zero (hence there are $x, y \in X$ such that $x * y \approx \mathbf{0}$ and, therefore, $[x] * [y] = [x * y] \neq [\mathbf{0}]$). Notice that, by Remark 5.11 above, equipped with the quotient topology τ_q , $X/I(\mathbf{0})$ is a Hausdorff topological real algebra. Clearly, \preceq is a continuous total order on $X/I(\mathbf{0})$. Thus, by Remark 5.10(i) and (ii), $X/I(\mathbf{0})$ can be identified with the reals and $\tau_{\preceq} \equiv \tau_q$. Therefore, endowed with the order topology τ_{\preceq} , $X/I(\mathbf{0})$ is a non-zero topological real algebra. Let us show that X is a topological real algebra too with respect to τ_{\lesssim} .

To that end, consider the projection map $p : X \rightarrow X/I(\mathbf{0})$. By Remark 5.11 p is linear, multiplicative, continuous and open. Also, it is very simple to see that p is order-preserving (i.e., $x \lesssim y \Leftrightarrow [x] \preceq [y]$ for every $x, y \in X$). Furthermore, p is also continuous and open whenever X is endowed with the order topology τ_{\lesssim} and $X/I(\mathbf{0})$ is equipped with the order topology τ_{\preceq} .

Let us show that, whenever X is equipped with the order topology, the binary operation $+ : X \times X \rightarrow X$ is a continuous map. The proofs for the remaining operations, namely “ $\cdot_{\mathbb{R}}$ ” and “ $*$ ”, are similar. Let then $C \subseteq X$ be a τ_{\lesssim} -open subset of X . We have to find τ_{\lesssim} -open subsets, say $A, B \subseteq X$, such that $A + B = \{a + b : a \in A, b \in B\} \subseteq C$. Since, as mentioned above, $p : X \rightarrow X/I(\mathbf{0})$ is a τ_{\lesssim} to τ_{\preceq} continuous and open map, it follows that $p(C)$ is a τ_{\preceq} -open subset of $X/I(\mathbf{0})$. Now, since, by above, $X/I(\mathbf{0})$ is a non-zero topological real algebra with respect to τ_{\preceq} , there are τ_{\preceq} -open subsets, say E, F of $X/I(\mathbf{0})$ such that $E + F \subseteq p(C)$. Consider $A = p^{-1}(E)$ and $B = p^{-1}(F)$ and let us prove that $A, B \subseteq X$ are the required subsets. Indeed, A, B are τ_{\lesssim} -open subsets of X since p is continuous and E, F are τ_{\preceq} of $X/I(\mathbf{0})$. It remains to see that $A + B \subseteq C$. To that end, let $a \in A, b \in B$ arbitrarily chosen. Then, there are $e \in E, f \in F$ such that $p(a) = e$ and $p(b) = f$. By linearity, $p(a + b) = p(a) + p(b) = e + f \in p(C)$. This is, there is $c \in C$ such that $p(a + b) = p(c)$ and therefore $a + b \sim c$. This straightforwardly implies that $a + b \in C$ since C is a τ_{\lesssim} -open subset of X which, in particular means that if $c \in C$, then $c' \in C$ for every $c' \sim c$.

For the converse, (ii) \implies (i), we argue as follows. Let $\preceq \in \tilde{\mathcal{P}}$. Let us prove that there is a continuous algebraic order-preserving function for \preceq . Consider the order topology τ_{\preceq} on X , so that $(X, \tau_{\preceq}, +, \cdot_{\mathbb{R}}, *)$ is a topological real algebra, by hypothesis.

Let us see first that $I(\mathbf{0})$ is both a linear subspace and an ideal of X , for which we need to prove the following two properties:

- (a) $I(\mathbf{0})$ is a real vector subspace of X .
- (b) For every $x \in I(\mathbf{0})$, $y \in X$, it holds that $x * y \in I(\mathbf{0})$ and $y * x \in I(\mathbf{0})$.

Let $x, y \in I(\mathbf{0})$. Since \preceq is translation-invariant, it follows that $x + y \sim x + \mathbf{0} = x \sim \mathbf{0}$. So, in order to prove (a), it is sufficient to see that, given $x \in I(\mathbf{0})$ and $\lambda \in \mathbb{R}$, then $\lambda \cdot x \sim \mathbf{0}$. If $\lambda \geq 0$, then, by homotheticity, $\lambda \cdot x \sim \mathbf{0}$. If $\lambda < 0$, then $(-\lambda) \cdot x \sim \mathbf{0}$. But $(-\lambda) \cdot x = -\lambda \cdot x$ and, by translation invariance of \preceq , $\lambda \cdot x \sim \mathbf{0}$.

To prove (b), let $y \in X$ and $x \in I(\mathbf{0})$. If $\mathbf{0} \preceq y$, then, because \preceq is multiplicative-invariant, $x * y \preceq \mathbf{0} * y = \mathbf{0}$ and $\mathbf{0} = \mathbf{0} * y \preceq x * y$. Therefore, $x * y \sim \mathbf{0}$. If $y \preceq \mathbf{0}$, then $\mathbf{0} \preceq -y$ and so $-x * y \sim \mathbf{0}$. Since $x * y = -(-x) * y$, and $I(\mathbf{0})$ is a vector subspace of X , it holds that $x * y \sim \mathbf{0}$. The case $y * x \in I(\mathbf{0})$ is proved in a similar way. Thus (b) holds and therefore $I(\mathbf{0})$ is an ideal of X .

Consider again the quotient space $X/I(\mathbf{0})$. It is now routine to see that $X/I(\mathbf{0})$ is a totally ordered real algebra with the usual binary operations induced by “+”, “ $\cdot_{\mathbb{R}}$ ” and “*” in the quotient $X/I(\mathbf{0})$, and the total order \preceq defined as $[x] \preceq [y] \iff x \preceq y$. As above, $X/I(\mathbf{0})$ is non-zero since, by hypothesis, \preceq is non-zero. So, by Lemma 5.9, there is a τ_{\preceq} -continuous algebraic order-preserving function $u : X/I(\mathbf{0}) \rightarrow \mathbb{R}$ for \preceq . Consider again the projection map $p : X \rightarrow X/I(\mathbf{0})$. Recall that p is linear, multiplicative, continuous and order-preserving. Then, by considering the composition $\psi = u \circ p : X \rightarrow \mathbb{R}$ we obtain a τ_{\preceq} -continuous (hence τ -continuous since \preceq is continuous), linear and multiplicative order-preserving function for \preceq . Thus τ satisfies CARP. \square

Remark 5.13. If X has an identity, say $1 \in X$, then the only zero total preorder on X is the trivial one \preceq_t . So, if X has an identity, then $\mathcal{P} = \tilde{\mathcal{P}} \cup \{\preceq_t\}$. Note that the null function is a continuous algebraic order-preserving function for \preceq_t . Moreover, for this preorder \preceq_t , the quotient space $X/I(\mathbf{0})$ is the zero vector (i.e., $X/I(\mathbf{0}) = \{\mathbf{0}\}$) and therefore the quotient topology amounts to the order topology (which coincides with the discrete topology, $\tau_d = \{\emptyset, X\}$, on $X/I(\mathbf{0})$). Then, Theorem 5.12 can be re-formulated in the following terms.

Let X be a real algebra with an identity and let τ be a topology on X . Then the following assertions are equivalent:

- (i) Every $\preceq \in \mathcal{P}$ admits a continuous algebraic order-preserving function.
- (ii) Endowed with the order topology τ_{\preceq} , X is a topological real algebra for every $\preceq \in \mathcal{P}$.

If τ is a topology on X that makes it to be a topological real algebra, then we have the following interesting consequence.

Corollary 5.14. *Let X be a real algebra equipped with a topology τ . If X is a topological real algebra, then τ satisfies CARP.*

Proof. According to the statement of Theorem 5.12, we only have to prove that, equipped with the order topology τ_{\lesssim} , X is a topological real algebra for every $\lesssim \in \tilde{\mathcal{P}}$. Let then be \lesssim such a total preorder on X and consider again the quotient space $X/I(\mathbf{0})$. As seen in the proof of Theorem 5.12, $X/I(\mathbf{0})$ is a non-zero totally ordered topological real algebra. Then, the result follows directly from Remark 5.10(i) and (ii) above and the argument provided in the last paragraph of the proof of the implication (i) \implies (ii) of Theorem 5.12. \square

Remark 5.15. In particular, Corollary 5.14 applies to the n -dimensional Euclidean space \mathbb{R}^n endowed with the usual binary operations, “+”, “ $\cdot_{\mathbb{R}}$ ” and “ \cdot^* ”, defined componentwise, and the Euclidean topology. Moreover, in this case it can be easily seen that the non-null algebra-homomorphisms are of the form $\psi(x_1, \dots, x_j, \dots, x_n) = x_j$ for some $j \in \{1, \dots, n\}$. Actually, in this case, continuity and order-preserving are redundant properties. Indeed, let $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$ such a function. Then, since ψ is linear, it is of the form $\psi(x_1, \dots, x_i, \dots, x_n) = \sum_i a_i x_i$, where $a_i \in \mathbb{R}$ for every $i \in \{1, \dots, n\}$. Now, let us denote by $(e^i)_{i=1}^n$ the canonical basis of \mathbb{R}^n . Then, since ψ is multiplicative too, in particular we have that $\psi(e^i * e^j) = \psi(e^i)\psi(e^j) = a_i a_j$ for all i, j . But, if $i \neq j$, $\psi(e^i * e^j) = \psi(\mathbf{0}) = 0 = a_i a_j$ and $a_i^2 = \psi(e^i)\psi(e^i) = \psi(e^i * e^i) = \psi(e^i) = a_i$. Therefore, since ψ is non-null, there is $j \in \{1, \dots, n\}$ such that $a_j = 1$ and $a_i = 0$ for $i \neq j$. Thus, $\psi(x_1, \dots, x_j, \dots, x_n) = x_j$. By using a similar argument, it is not difficult to prove that this kind of result also holds true for the infinite-dimensional cases $X = c_0$ or $X = l_1$, but not for $X = l_\infty$ (see, Remark 5.4 and Remarks 5.8(iii)).

Note that, by Corollary 5.14, we have proved that any non-zero (or, equivalently in this case, non-trivial) continuous total preorder defined on \mathbb{R}^n which is translation-invariant, homothetic and multiplicative-invariant is projective.

6. Application to social choice theory

In this section, we develop an application of the algebraic approach presented in the previous Section 5 to the context of utility theory in the social choice framework. A social welfare functional is a map that assigns a preference relation (total preorder) to any profile of individual utilities. In this literature, a utility function refers to any map $u : X \rightarrow \mathbb{R}$, where X is the choice set. It should be noted that a utility function u generates a total preorder on X , denoted by \lesssim_u , defined as follows: $x \lesssim_u y$ if and only if $u(x) \leq u(y)$ for every $x, y \in X$. We characterize strongly dictatorial social welfare functionals in terms of invariance and continuity properties of such rules, in the spirit of the literature of utility measurability and (inter/intra) personal comparability (see, e.g., Fleurbaey and Hammond [22]). Before presenting the application some definitions and notations are needed. Other insights regarding the potential use

of algebraic techniques in social choice theory can be seen in Candeal, Induráin and Molina [15].

Let X be a nonempty set (usually called in this context the *set of alternatives* or the *choice set*). Let us denote by \mathfrak{R} the class of all total preorders (or preference relations as are usually called in this setting) defined on X . Let $n > 1$ be a natural number (number of agents or individuals in the society). The set of all functions from X into \mathbb{R} will be denoted by U . A utility function for the agent i will be denoted by $u_i \in U$. A *profile* of utility functions, one for each agent, will be denoted by (u_1, \dots, u_n) , i.e., it is an element of the usual product of n -copies of U , denoted by U^n , which is usually referred to as the set of all possible profiles. The set which consists of the n first natural numbers will be denoted by N , i.e., $N = \{1, \dots, n\}$.

A *social welfare functional* is a rule $F : U^n \rightarrow \mathfrak{R}$ that assigns a preference relation $F(u_1, \dots, u_n) \in \mathfrak{R}$, interpreted as the social preference relation, to any profile (u_1, \dots, u_n) in the domain U^n . In order to shorten the notation, for a profile (u_1, \dots, u_n) we will use the notation (u_i) . Also, $F(u_i)_s$ stands for the strict preference (asymmetric relation) associated with $F(u_i)$.

A social welfare functional $F : U^n \rightarrow \mathfrak{R}$ is said to be *Paretian* if, for any pair of alternatives $x, y \in X$ and any profile $(u_i) \in U^n$, we have that $u_i(x) \leq u_i(y)$ for all $i \in N$ implies $x F(u_i) y$ and also that $u_i(x) < u_i(y)$ for all $i \in N$ implies $x F(u_i)_s y$.

A social welfare functional $F : U^n \rightarrow \mathfrak{R}$ satisfies the *binary independence of irrelevant alternatives* condition if, for any pair of alternatives $x, y \in X$ and any pair of profiles $(u_i), (v_i) \in U^n$ with the property that $u_i(x) = v_i(x)$ and $u_i(y) = v_i(y)$ for all $i \in N$, we have that $x F(u_i) y$ if and only if $x F(v_i) y$.

A social welfare functional $F : U^n \rightarrow \mathfrak{R}$ is called *strongly dictatorial* if there is an agent $j \in N$ (the dictator) such that $F(u_i)$ coincides with \succsim_{u_j} for every $(u_i) \in U^n$, where \succsim_{u_j} denotes the preference relation on X induced by the utility function of the j -agent, u_j .

A social welfare functional $F : U^n \rightarrow \mathfrak{R}$ is called *anonymous* if for any pair of profiles $(u_i), (v_i) \in U^n$ such that (v_i) is derived from (u_i) by permuting the individuals' utility functions, one has that $F(u_i)$ and $F(v_i)$ coincide.

A social welfare functional $F : U^n \rightarrow \mathfrak{R}$ is called *continuous* if $\{(a_i), (b_i)\} \in \mathbb{R}^n \times \mathbb{R}^n$: there exist $(u_i) \in U^n$, $x, y \in X$ with $u_i(x) = a_i$, $u_i(y) = b_i$ and such that $x F(u_i) y$ is an Euclidean-closed subset of $\mathbb{R}^n \times \mathbb{R}^n$.

Finally, a social welfare functional $F : U^n \rightarrow \mathfrak{R}$ satisfies *information invariance with respect to cardinal measurability* if, for any profile $(u_i) \in U^n$, and any n -tuple of functions (ϕ_1, \dots, ϕ_n) such that, for each $i \in N$, the map $\phi_i : \mathbb{R} \rightarrow \mathbb{R}$ is of the form $\phi_i(t) = a_i + b_i t$ ($t \in \mathbb{R}$) with $a_i \in \mathbb{R}$ and $b_i > 0$, it holds that $F(u_i)$ and $F(\phi_i \circ u_i)$ coincide.

We reach the following result.

Theorem 6.1. *Suppose that X contains at least three elements and let $F : U^n \rightarrow \mathfrak{R}$ be a social welfare functional. Then the following assertions are equivalent:*

- (i) *F is Paretian, satisfies the binary independence of irrelevant alternatives condition, continuity, and information invariance with respect to cardinal measurability.*
- (ii) *F is strongly dictatorial.*

Proof. Let us prove that (ii) implies (i). Suppose then that F is strongly dictatorial and assume without loss of generality that the j -th agent acts as the dictator. It is straightforward to see that F is Paretian and also that it satisfies the binary independence of irrelevant alternatives condition. In order to prove that it satisfies information invariance with respect to cardinal measurability, let $(u_i) \in U^n$ a profile of utility functions and let (ϕ_1, \dots, ϕ_n) be a n -tuple of functions so that, for every $i \in N$, $\phi_i(t) = a_i + b_i t$ ($t \in \mathbb{R}$) with $a_i \in \mathbb{R}$ and $b_i > 0$. Since, by hypothesis, the j -th agent is the dictator for F , it follows that $F(u_i) = \succsim_{u_j}$. Now, consider the profile $(\phi_i \circ u_i) \in U^n$. Then, $F(\phi_i \circ u_i) = \succsim_{\phi_j \circ u_j} = \succsim_{u_j}$, since $u_j(x) \leq u_j(y)$ if and only if $\phi_j \circ u_j(x) \leq \phi_j \circ u_j(y)$ for every $x, y \in X$ which means that u_j and $\phi_j \circ u_j$ define the same total preorder in \mathfrak{R} . Therefore, $F(u_i) = F(\phi_i \circ u_i)$ and so F satisfies information invariance with respect to cardinal measurability. It remains to prove that F is continuous. To that end, observe that $\{(a_i), (b_i)\} \in \mathbb{R}^n \times \mathbb{R}^n$: there exist $(u_i) \in U^n$, $x, y \in X$ with $u_i(x) = a_i$, $u_i(y) = b_i$ and $x F(u_i) y\} = \{(a_i), (b_i)\} \in \mathbb{R}^n \times \mathbb{R}^n$: there exist $(u_i) \in U^n$, $x, y \in X$ with $u_i(x) = a_i$, $u_i(y) = b_i$ and $u_j(x) \leq u_j(y)\} = \{(a_i), (b_i)\} \in \mathbb{R}^n \times \mathbb{R}^n$: $a_j \leq b_j\}$, which, clearly, is a Euclidean-closed subset of $\mathbb{R}^{2n} = \mathbb{R}^n \times \mathbb{R}^n$.

To prove the converse, consider the binary relation \succsim^* defined on \mathbb{R}^n as follows: Given $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n) \in \mathbb{R}^n$, then $a \succsim^* b$ if and only if there exist $x, y \in X$ and a profile $(u_i) \in U^n$ such that for every $i \in N$ it holds that $u_i(x) = a_i$, $u_i(y) = b_i$ and $x F(u_i) y$. Since F satisfies the binary independence of irrelevant alternatives condition, and X contains at least three elements, it can be shown that \succsim^* is a total preorder that “generates” F . Indeed, by definition, \succsim^* is obviously reflexive. It is total since the domain of F is U^n . Let us see that it is transitive too. To that end, let $a = (a_i), b = (b_i), c = (c_i) \in \mathbb{R}^n$ such that $a \succsim^* b$ and $b \succsim^* c$. We want to show that $a \succsim^* c$. Since $a \succsim^* b$ there are $x, y \in X$ and a profile of utility functions $(u_i) \in U^n$ such that $u_i(x) = a_i$, $u_i(y) = b_i$ for all i , and $x F(u_i) y$. Similarly, since $b \succsim^* c$, there are $z, t \in X$ and a profile of utility functions $(v_i) \in U^n$ such that $v_i(z) = b_i$, $v_i(t) = c_i$ for all i and $z F(v_i) t$. Consider a profile of utility functions $(w_i) \in U^n$ such that, for each $i \in N$, $w_i(x) = a_i$, $w_i(y) = w_i(z) = b_i$ and $w_i(t) = c_i$. Note that such a profile $(w_i) \in U^n$ does exist since U consists of all functions from X into \mathbb{R} and, by hypothesis, X has at least three elements (note that it might be that $y = z$). Then, by the binary independence of irrelevant alternatives condition and the fact that $x F(u_i) y$ it follows $x F(w_i) y$. Similarly, we have

that $yF(w_i)t$. Now, $F(w_i) \in \mathfrak{R}$, hence it is a transitive binary relation defined on X . Thus, $xF(w_i)t$, for such a profile $(w_i) \in U^n$ for which $w_i(x) = a_i$ and $w_i(t) = c_i$. Therefore, by definition of \succsim^* , $a \succsim^* c$. So, \succsim^* is transitive. In addition, it should be noted that, since F is Paretian, \succsim^* is non-trivial and increasing⁵. Moreover, for every $a, b \in \mathbb{R}^n$ such that $a \ll b$ it holds that $a \prec^* b$. Let us show that $\succsim^* \in \tilde{\mathcal{P}}$, where the binary operations on \mathbb{R}^n “+”, “ $\cdot_{\mathbb{R}}$ ” and “*” are the usual ones defined componentwise.

Indeed, \succsim^* is non-zero since 1_n is the identity of \mathbb{R}^n with respect to “*” and $0_n \prec^* 1_n$, hence \succsim^* is non-trivial, because F is Paretian. Continuity of \succsim^* , referred to the Euclidean topology in \mathbb{R}^n , follows directly from the continuity of F (see in Section 2 the equivalent formulation of continuity for a total preorder). Let us prove now that \succsim^* is translation-invariant. To that end, let $a = (a_i), b = (b_i), c = (c_i) \in \mathbb{R}^n$ such that $a \succsim^* b$. Then there are $x, y \in X$ and a profile of utility functions $(u_i) \in U^n$ such that $u_i(x) = a_i, u_i(y) = b_i$ for all i and $xF(u_i)y$. For every i , consider the functions $v_i(z) = u_i(z) + c_i$ for all $z \in X$. By information invariance with respect to cardinal measurability, it holds that $F(v_i)$ coincides with $F(u_i)$. In particular, $xF(v_i)y$ or, equivalently, $(v_i(x)) \succsim^* (v_i(y))$. Thus, $a + c = (a_i + c_i) \succsim^* b + c = (b_i + c_i)$.

Similarly, let us prove that \succsim^* is homothetic. To see this, let $a = (a_i), b = (b_i) \in \mathbb{R}^n$, with $0 \leq \lambda$, such that $a \succsim^* b$. If $\lambda = 0$, obviously $\lambda \cdot a \succsim^* \lambda \cdot b$. So, suppose that $0 < \lambda$. Then, there are $x, y \in X$ and a profile of utility functions $(u_i) \in U^n$, such that $u_i(x) = a_i, u_i(y) = b_i$ for all i and $xF(u_i)y$. For each i , consider the (utility) function $v_i = \lambda u_i$. By information invariance with respect to cardinal measurability, it holds that $F(v_i)$ and $F(u_i)$ coincide. In particular, $xF(v_i)y$ or, equivalently, $(v_i(x)) \succsim^* (v_i(y))$. Thus, $\lambda \cdot a = (\lambda a_i) \succsim^* \lambda \cdot b = (\lambda b_i)$. Notice that, by translation invariance and homotheticity of \succsim^* it follows, as in the proof of Theorem 5.12, that $I(0_n) = \{z \in \mathbb{R}^n : z \sim^* 0_n\}$ is a vector subspace of \mathbb{R}^n .

The fact that \succsim^* is multiplicative-invariant, with respect to the binary operation $*$ defined componentwise (i.e., $(a_i) * (b_i) = (a_i b_i)$ for every $(a_i), (b_i) \in \mathbb{R}^n$), is a little more tricky. Since \succsim^* is translation-invariant it should be noted that, in order to prove that \succsim^* is multiplicative-invariant, it is sufficient to show that for any pair $a, b \in \mathbb{R}^n$ such that $0_n \succsim^* a, b$ it holds that $0_n \succsim^* a * b$. To show this it will be useful to observe the following facts:

(1) Given any $a = (a_i) \in \mathbb{R}^n$, such that $0_n \succsim^* a = (a_i)$, it holds that $0_n \succsim^* a * c$ for all $0_n \ll c = (c_i)$ (In particular, this will imply that if $0_n \sim^* a$, then $0_n \sim^* a * c$ for all $0_n \ll c$). Indeed, since $0_n \succsim^* a$, there are $x, y \in X$ and a profile of utility functions $(u_i) \in U^n$, such that $u_i(x) = 0, u_i(y) = a_i$ for all i and $xF(u_i)y$. For each i , consider the (utility) function

⁵A total preorder \preceq defined on \mathbb{R}^n is said to be *increasing* if $x \leq y$ implies $x \preceq y$ for every $x, y \in \mathbb{R}^n$ (Here, $x = (x_1, \dots, x_n) \leq y = (y_1, \dots, y_n)$ means that $x_i \leq y_i$ for all $i \in N$). “ \ll ” will stand for the strict partial order in \mathbb{R}^n (i.e., $x \ll y$ if and only if $x_i < y_i$ for all $i \in N$). The zero vector in \mathbb{R}^n and the vector with all its coordinates equal one will be denoted by 0_n and 1_n , respectively.

$v_i = c_i u_i$. By information invariance with respect to cardinal measurability, it holds that $F(v_i)$ and $F(u_i)$ coincide. In particular, $x F(v_i) y$ or, equivalently, $(v_i(x)) \succ^* (v_i(y))$. Thus, $0_n \succ^* a * c = (a_i c_i)$.

(2) Let then $a = (a_i) \in \mathbb{R}^n$ such that $0_n \succ^* a$ and denote by $(e^i)_{i \in N}$ the canonical basis in \mathbb{R}^n . Let us see that, for each $i \in N$, $0_n \succ^* a_i \cdot e^i$ and, in addition, if $a_i \leq 0$, then $0_n \sim^* a_i \cdot e^i$. Indeed, let $i \in N$ be fixed. If $a_i \geq 0$, then the fact that $0_n \succ^* a_i \cdot e^i$ follows directly since \succ^* is increasing and so $0_n \leq a_i \cdot e^i$ implies that $0_n \succ^* a_i \cdot e^i$. If $a_i < 0$ the argument requires a further consideration. On the one hand, the fact that $a_i \cdot e^i \succ^* 0_n$ follows by the previous reasoning again. To prove that $0_n \succ^* a_i \cdot e^i$ consider, for every $m \in \mathbb{N}$, the vector $c^m = (c_k^m) \in \mathbb{R}^n$ defined as follows: $c_k^m = \frac{1}{m}$, if $k \neq i$ and $c_i^m = 1$. Clearly, the sequence $(c^m)_{m=1}^\infty$ converges to e^i , as m goes to ∞ , and therefore the sequence $(a * c^m)_{m=1}^\infty$ converges to $a_i \cdot e^i$, as m goes to ∞ . Moreover, for every $m \in \mathbb{N}$, $0_n \ll c^m$ and, since $0_n \succ^* a$, it follows also, from the argument of the previous paragraph, that $0_n \succ^* a * c^m$. Now, the continuity of \succ^* clearly implies that $0_n \succ^* a_i \cdot e^i$. This fact, in combination with $a_i \cdot e^i \succ^* 0_n$, yields $0_n \sim^* a_i \cdot e^i$. Thus, $0_n \sim^* a_i \cdot e^i$ provided that $a_i \leq 0$ (the case $a_i = 0$ is trivial).

Let then $a = (a_i), b = (b_i) \in \mathbb{R}^n$ such that $0_n \succ^* a, b$ and let us prove that $0_n \succ^* a * b$. Notice that $a * b = \sum (a_i b_i) \cdot e^i$. Let us show that, for all $i \in N$, $0_n \succ^* (a_i b_i) \cdot e^i$. Then, the result would follow from the fact that \succ^* is translation-invariant. Let $i \in N$ be fixed. If $0 \leq a_i b_i$, then $0_n \succ^* (a_i b_i) \cdot e^i$ since \succ^* is increasing. It remains to analyze what occurs if $a_i b_i < 0$. Assume that $a_i < 0$ and $b_i > 0$, the other case (i.e., $a_i > 0$ and $b_i < 0$) being similar. Then, by the argument above, $0_n \sim^* a_i \cdot e^i$, i.e., $a_i \cdot e^i \in I(0_n)$. Now, since $I(0_n)$ is a vector subspace of \mathbb{R}^n , it follows that $(a_i b_i) \cdot e^i = b_i \cdot (a_i \cdot e^i) \in I(0_n)$. In particular, we have that $0_n \succ^* (a_i b_i) \cdot e^i$ as desired.

Thus, \succ^* is multiplicative-invariant. Hence all the conditions of Remark 5.15 are satisfied and therefore \succ^* is a projective total preorder on \mathbb{R}^n . This clearly implies that F is a strongly dictatorial social welfare functional. \square

Remark 6.2. In the proof of Theorem 6.1 above we have used the fact that, with respect to the usual binary operations “+”, “ $\cdot_{\mathbb{R}}$ ” and “*”, defined componentwise, $I(0_n)$ is a vector subspace of \mathbb{R}^n . In fact, as it was proved in the proof of Theorem 5.12, $I(0_n)$ is also an ideal of \mathbb{R}^n . So, a fortiori, $I(0_n)$ is of the form $I(0_n) = \sum_{i \in N, i \neq j} \lambda_i e^i$ for some $j \in N$. But this fact is a consequence of Theorem 5.12 and Remark 5.15

As a direct consequence of the previous Theorem 6.1, we obtain the following impossibility result.

Corollary 6.3. *Suppose that X contains at least three elements. Then there is no continuous social welfare functional $F : U^n \rightarrow \mathfrak{R}$ that is Paretian, satisfies the binary independence of irrelevant alternatives condition, information invariance with respect to cardinal measurability, and anonymity.*

Acknowledgements. Thanks are given to an anonymous referee for the excellent revision done. His/her suggestions and comments have led to substantial improvements in the manuscript.

This work has been supported by the research projects MTM 2007-62499, ECO2008-01297 and MTM2009-12872-C02-02 (Spain).

References

- [1] C. D. Aliprantis and K. C. Border, *Infinite Dimensional Analysis*, A Hitchhiker's guide, Springer, Berlin, 1999.
- [2] G. Birkhoff, *A note on topological groups*, Compos. Math. **3** (1936), 427–430.
- [3] D. S. Bridges and G. B. Mehta, *Representation of Preference Orderings*, Springer, Berlin, 1995.
- [4] M. J. Campión, J. C. Candeal, and E. Induráin, *The existence of utility functions for weakly continuous preferences on a Banach space*, Math. Social Sci. **51** (2006), no. 2, 227–237.
- [5] ———, *On Yi's extension property for totally preordered topological spaces*, J. Korean Math. Soc. **43** (2006), no. 1, 159–181.
- [6] ———, *Preorderable topologies and order-representability of topological spaces*, Topology Appl. **156** (2009), no. 18, 2971–2978.
- [7] ———, *Semicontinuous planar total preorders on non-separable metric spaces*, J. Korean Math. Soc. **46** (2009), no. 4, 701–711.
- [8] M. J. Campión, J. C. Candeal, E. Induráin, and G. B. Mehta, *Representable topologies and locally connected spaces*, Topology Appl. **154** (2007), no. 10, 2040–2049.
- [9] J. C. Candeal, C. Hervés, and E. Induráin, *Some results on representation and extension of preferences*, J. Math. Econom. **29** (1998), no. 1, 75–81.
- [10] J. C. Candeal and E. Induráin, *Utility functions on chains*, J. Math. Econom. **22** (1993), no. 2, 161–168.
- [11] ———, *A note on linear utility*, Econom. Theory **6** (1995), no. 3, 519–522.
- [12] ———, *Lexicographic behaviour of chains*, Arch. Math. (Basel) **72** (1999), no. 2, 145–152.
- [13] J. C. Candeal, E. Induráin, and G. B. Mehta, *Order preserving functions on ordered topological vector spaces*, Bull. Austral. Math. Soc. **60** (1999), no. 1, 55–65.
- [14] ———, *Utility functions on locally connected spaces*, J. Math. Econom. **40** (2004), no. 6, 701–711.
- [15] J. C. Candeal, E. Induráin, and J. A. Molina, *Numerical representability of ordered topological spaces with compatible algebraic structure*, To appear in Order. DOI 10.1007/s11083-011-9202-8.
- [16] H. H. Corson, *The weak topology of a Banach space*, Trans. Amer. Math. Soc. **101** (1961), 1–15.
- [17] G. Debreu, *Representation of a preference ordering by a numerical function*, In R. Thrall, C. Coombs and R. Davies (Eds.), Decision processes, John Wiley. New York, 1954.
- [18] J. Dugundji, *Topology*, Allyn and Bacon, Boston, 1966.
- [19] S. Eilenberg, *Ordered topological spaces*, Amer. J. Math. **63** (1941), 39–45.
- [20] R. Engelking, *General Topology*, Revised and Completed Edition. Heldermann Verlag, Berlin, 1989.
- [21] M. Estévez and C. Hervés, *On the existence of continuous preference orderings without utility representations*, J. Math. Econom. **24** (1995), no. 4, 305–309.

- [22] M. Fleurbaey and P. J. Hammond, *Interpersonally comparable utility*, Chapter 21 of the Handbook of Utility Theory (Vol. 2), Edited by Barberà S, Hammond PJ and Seidl C, Kluwer Academic Publishers. Amsterdam, 2004.
- [23] L. Fuchs, *Partially Ordered Algebraic Systems*, Pergamon Press, Reading, Massachusetts, 1963.
- [24] V. Gutev, *Weak orderability of second countable spaces*, Fund. Math. **196** (2007), no. 3, 275–287.
- [25] G. Herden, *Topological spaces for which every continuous total preorder can be represented by a continuous utility function*, Math. Social Sci. **22** (1991), no. 2, 123–136.
- [26] G. Herden and A. Pallack, *Useful topologies and separable systems*, Appl. Gen. Topol. **1** (2000), no. 1, 61–82.
- [27] S. Kakutani, *Über die Metrisation der topologischen Gruppen*, Proc. Imp. Acad. **12** (1936), no. 4, 82–84.
- [28] J. L. Kelley, *General Topology*, Van Nostrand. New York, 1955.
- [29] D. J. Lutzer and H. R. Bennet, *Separability, the countable chain condition and the Lindelöf property on linearly ordered spaces*, Proc. Amer. Math. Soc. **23** (1969), 664–667.
- [30] J. R. Munkres, *Topology*, (2nd ed.), Prentice Hall. New York, 2000.
- [31] S. Purisch, *A history of results on orderability and suborderability*, In Handbook of the history of General Topology, Vol. 2, pp. 689–702. Kluwer. Dordrecht, 1998.
- [32] C. E. Rickart, *General Theory of Banach Algebras*, Van Nostrand. Princeton, N.J., 1960.
- [33] H. H. Schaefer, *Topological Vector Spaces*, (3rd ed.), Springer Verlag. New York, 1971.
- [34] P. Simon, *A connected, not separably connected metric space*, Rend. Istit. Mat. Univ. Trieste **32** (2001), suppl. 2, 127–133.
- [35] L. A. Steen and J. A. Jr. Seebach, *Counterexamples in Topology*, (2nd ed.), Springer Verlag. New York, 1978.
- [36] J. van Dalen and E. Wattel, *A topological characterization of ordered spaces*, General Topology and Appl. **3** (1973), 347–354.
- [37] G. Yi, *Continuous extension of preferences*, J. Math. Econom. **22** (1993), no. 6, 547–555.

MARÍA JESÚS CAMPIÓN
DEPARTAMENTO DE MATEMÁTICAS
UNIVERSIDAD PÚBLICA DE NAVARRA
E-31006 PAMPLONA, SPAIN
E-mail address: mjesus.campion@unavarra.es

JUAN CARLOS CANDEAL
DEPARTAMENTO DE ANÁLISIS ECONÓMICO
UNIVERSIDAD DE ZARAGOZA
E-50005 ZARAGOZA, SPAIN
E-mail address: canddeal@unizar.es

ESTEBAN INDURÁIN
DEPARTAMENTO DE MATEMÁTICAS
UNIVERSIDAD PÚBLICA DE NAVARRA
E-31006 PAMPLONA, SPAIN
E-mail address: steiner@unavarra.es

GHANSHYAM BHAGVANDAS MEHTA
DEPARTMENT OF ECONOMICS
UNIVERSITY OF QUEENSLAND
BRISBANE 4072, AUSTRALIA
E-mail address: mehtagb@gmail.com