



Effect of 24 h Fasting on Gene Expression of AMPK, Appetite Regulation Peptides and Lipometabolism Related Factors in the Hypothalamus of Broiler Chicks

Liu Lei and Zhu Lixian*

College of Animal Science and Technology, Shandong Agricultural University, Taian, Shandong, 271018, China

ABSTRACT: The 5'-adenosine monophosphate-activated protein kinase (AMPK) is a key part of a kinase-signaling cascade that acts to maintain energy homeostasis. The objective of this experiment was to investigate the possible effects of fasting and refeeding on the gene expression of hypothalamic AMPK, some appetitive regulating peptides and lipid metabolism related enzymes. Seven-day-old male broiler (Arbor Acres) chicks were allocated into three equal treatments: fed *ad libitum* (control); fasted for 24 h; fasted for 24 h and then refed for 24 h. Compared with the control, the hypothalamic gene expression of AMPK α 2, AMPK β 1, AMPK β 2, AMPK γ 1, Ste20-related adaptor protein β (STRAD β), mouse protein 25 α (MO25 α) and agouti-related peptide (AgRP) were increased after fasting for 24 h. No significant difference among treatments was observed in mRNA levels of AMPK α 1, AMPK γ 2, LKB1 and neuropeptide Y (NPY). However, the expression of MO25 β , pro-opiomelanocortin (POMC), corticotropin-releasing hormone (CRH), ghrelin, fatty acid synthase (FAS), acetyl-CoA carboxylase α (ACC α), carnitine palmitoyltransferase 1 (CPT-1) and sterol regulatory element binding protein-1 (SREBP-1) were significantly decreased. The present results indicated that 24 h fasting altered gene expression of AMPK subunits, appetite regulation peptides and lipometabolism related factors in chick's hypothalamus; the hypothalamic FAS signaling pathway might be involved in the AMPK regulated energy homeostasis and/or appetite regulation in poultry. (**Key Words:** Metabolism, Nutrition, Peptides, Lipid, Hypothalamus, Broiler)

INTRODUCTION

AMPK is the central component of a cellular signaling system that monitors cellular energy change, acting as 'metabolic master switch' to regulate ATP concentrations in the face of stress. AMPK can be activated by allosteric effect of AMP and/or by threonine phosphorylation of one subunit in response to AMPK kinases (Hong et al., 2003; Hawley et al., 2005; Hurley et al., 2005; Hardie, 2006). In some cases, activation of AMPK requires phosphorylation of T-172 by an upstream protein kinase such as LKB1 (Hawley et al., 1996). LKB1 is a serine/threonine protein kinase which was first discovered in the study of Peutz-Jeghers syndrome (Hemminki et al., 1998; Jenne et al., 1998). LKB1 is a complex with two accessory proteins, pseudokinase STRAD (alpha or beta isoforms) and scaffold

protein MO25 (alpha and beta isoforms) (Bass et al., 2003; Boudeau et al., 2003). AMPK can be activated in a variety of physiological circumstances, including hypoglycemia (da Silva Xavier et al., 2000), ischemia (Marsin et al., 2000), heat shock (Corton et al., 1994) and exercise (Mu et al., 2001; Musi et al., 2001). AMPK is also affected by several orexigenic and anorexigenic signals in the hypothalamus (e.g. thyroid hormones, ghrelin, insulin and leptin) (Kola et al., 2006). AMPK is widely expressed in the brain, including the areas that control feed intake and neuroendocrine function. Immunostaining revealed that various AMPK isoforms distributed in hypothalamus and the hindbrain (Turnley et al., 1999). In mammals, AMPK exists as a heterotrimeric enzyme complex consisting of one catalytic (alpha) subunit and two regulatory (beta and gamma) subunits (Mitchell et al., 1994). There are two known alpha subunit isoforms (alpha-1 and alpha-2, Stapleton et al., 1996), two beta subunit isoforms (beta-1 and beta-2, Stapleton et al., 1997) and three gamma subunit isoforms (gamma-1, gamma-2 and gamma-3, Cheung et al., 2000).

* Corresponding Author: Zhu Lixian. College of Food Science and Engineering, Shandong Agricultural University, Taian, Shandong, 271018 China. Tel: +86-5388241419, Fax: +86-538 8246029, E-mail: zhlix@sdau.edu.cn

Submitted Mar. 22, 2012; Accepted Jun. 2, 2012; Revised Jun. 13, 2012

Previous studies in mammals showed that hypothalamic AMPK integrates nutrient and hormonal signals from both anorexigenic and orexigenic pathways to regulate feed intake (Obici et al., 2003; Andersson et al., 2004; Minokoshi et al., 2004). Minokoshi et al. (2004) reported that fasting resulted in increased hypothalamic AMPK activity, whereas refeeding inhibited it (Minokoshi et al. 2004). AMPK is considered as a major regulator of fatty acid synthesis pathway due to its function to phosphorylate and inactivate key enzymes in fatty acid metabolism such as acetyl-CoA carboxylase (ACC). Pharmacological inhibition of fatty acid synthase (FAS) produces anorexia in mice (Loftus et al., 2000). It is reported that the activation of AMPK in the hypothalamus is associated with activation of mitochondrial enzyme carnitine palmitoyltransferase-1 (CPT1), leading to a decrease in the cellular levels of long-chain acyl-CoAs and increase feed intake (Andersson et al., 2004). The modest reduction of long-chain acyl-CoAs within the arcuate nucleus (ARC) would increase the expression of both agouti-related peptide (AgRP) and neuropeptide Y (NPY) and enhance feed intake (Obici et al., 2003).

There have been numerous reports showing that AMPK is involved in appetite regulation in mammals (Claret et al., 2007; Dzamko et al., 2010). Because feeding behavior and energy homeostasis are basic processes crucial to the survival of all animals (Richards and Proszkowiec-Weglarz, 2007), it is logical to assume that AMPK should be involved in poultry feed intake control. Gene expression of AMPK in chicken embryos has been explored (Proszkowiec-Weglarz and Richards, 2009), but information of its expression in chicks hypothalamus remains scarce. Previous studies in mammals showed that isoform composition of AMPK subunit changes during physiological and pathological conditions (Gregor et al., 2006; Kim et al., 2009). The aim of the present experiment was to investigate if AMPK subunits (α , β and γ) gene expression is modulated by 24 h fasting in meat-type chicks and to determine the interrelationship between AMPK and the mRNA levels of some hypothalamic feeding regulatory neuropeptides (i.e. corticotropin-releasing hormone (CRH), neuropeptide Y (NPY), agouti-related peptide (AgRP), pro-opiomelanocortin (POMC)) and fatty acid synthase (FAS).

MATERIALS AND METHODS

Animals, experimental protocol and sample collection

Male broiler (Arbor Acres) chicks were obtained from a local hatchery at 1 d of age and reared in an environmentally controlled room. The brooding temperature was maintained at 35°C (65% RH) for the first 2 d, and then decreased gradually to 31°C (45% RH) until 10 d. The light

regime was 23 L/1 D. All chicks received a starter diet with 21.5% crude protein and 12.33 MJ/kg of metabolizable energy (Zhao et al., 2009). All the birds had free access to feed and water during the rearing period. The study was approved by the University and carried out in accordance with the "Guidelines for Experimental Animal" of Ministry of Science and Technology (Beijing, P. R. China). All the chicks were cared for in accordance with the Guide to the Care and Use of Experimental Animals (Olfert et al., 1993).

Seven-day-old male broiler chicks were allocated into three equal treatments: control, fed *ad libitum* (C); fasted for 24 h (S24); fasted for 24 h and then refed for 24 h (S24R24). At the end of each treatment period, birds were sacrificed by exsanguination (Close, 1997), and then the whole hypothalamus was collected. After snap frozen in liquid nitrogen, the tissue samples were stored at -80°C for RNA extraction.

RNA isolation and analysis

The expression of genes in hypothalamus was quantified using quantitative real-time PCR with SYBR Green I labeling. Total RNA was isolated using the guanidinium isothiocyanate method with Trizol Reagent (Invitrogen, San Diego, CA, USA). The quality of the RNA was tested by electrophoresis on an agarose-gel and the quantity of the RNA was determined with biophotometer (Eppendorf, Germany).

RT reactions (10 μ l) consisted of 500 ng total RNA, 5 mmol/L MgCl₂, 1 μ l RT buffer, 1 mmol/L dNTP, 2.5 U AMV, 0.7 nmol/L oligo d(T) and 10 U Ribonuclease inhibitor (TaKaRa Biotechnology, Co., Ltd. Dalian, P. R. China). Real-time PCR analysis was conducted using the Applied Biosystems 7500 Real-time PCR System (Applied Biosystems, Foster, CA, USA). Each RT-reaction served as a template in a 20 μ l PCR reaction containing 0.2 μ mol/L of each primer and SYBR green master mix (Takara Biotechnology, Co., Ltd. Dalian, P. R. China). Primer-set sequences are described in Table 1. Real-time PCR reactions were performed at 95°C for 10 s, followed by 40 cycles at 95°C for 5 s and 60°C for 34 s. SYBR green fluorescence was detected at the end of each cycle to monitor the amount of PCR product. When calculating the efficiency of qPCR primers, a standard curve was made in 10 fold dilutions, and its slope was used to calculate efficiency.

The relative amount of mRNA for a gene was calculated according to the method of Livak and Schmittgen (Livak and Schmittgen, 2001). The mRNA levels of these genes were normalized to 18s rRNA levels (Δ CT) (Proszkowiec-Weglarz et al., 2006). The Δ CT was calibrated against an average of the control chicks. The linear amount of target molecules relative to the calibrator was calculated by $2^{-\Delta\Delta CT}$.

Table 1. Gene-specific primers used for the analysis of chick gene expression

Gene	GenBank accession no.	Primer sequences (5'-3')	Orientation	Product size (bp)
18s	AF173612	ATAACGAACGAGACTCTGGCA	Forward	136
		CGGACATCTAAGGGCATCACA	Reverse	
LKB1	NM_001045833	TGAGAGGGATGCTTGAATACGA	Forward	158
		ACTTGTCTTTGTTTCTGGGC	Reverse	
AMPK α 1	DQ302133	CGGAGATAAACAGAAGCACGAG	Forward	266
		CGATTCAGGATCTTCACTGCAAC	Reverse	
AMPK α 2	DQ340396	GGGACCTGAAACCAGAGAACG	Forward	215
		ACAGAGGAGGGCATAGAGGATG	Reverse	
AMPK β 1	XM_415278	ATGGTGGACTCCCAGAAGTG	Forward	254
		GAGCACCATCACTCCATCCT	Reverse	
AMPK β 2	BG713266	CTGTCATGGGGAACACCAC	Forward	363
		GGTCCAGGATAGCGACAAAG	Reverse	
AMPK γ 1	DQ133597	AGCTGCAGATCGGTACCTACA	Forward	200
		CGTCACGTCCAGGTTGTTGT	Reverse	
AMPK γ 2	DQ212708	ATCGGCATTACCTGTTGTGG	Forward	231
		ACCACCAAACGATGAACCTC	Reverse	
NPY	M87294	GAGGCACTACATCAACCTCATCAC	Forward	101
		TGTTTTCTGTGCTTCCCTCAA	Reverse	
AgRP	NM_001031457	GGAACCGCAGGCATTGTC	Forward	163
		GTAGCAGAAGGCGTTGAAGAA	Reverse	
POMC	NM_001031098	CGCTACGGCGCTTTGACGAT	Forward	88
		TCTTGTAGGCGCTTTGACGAT	Reverse	
CRH	NM_001123031	CTCCCTGGACCTGACTTTCC	Forward	86
		TGTTGCTGTGGGCTTGCT	Reverse	
Ghrelin	AB075215	CCTTGGGACAGAACTGCTC	Forward	203
		CACCAATTTCAAAGGAACG	Reverse	
POMC	NM_001031098	AGAACAGCAAGTGCCAGGAC	Forward	162
		TGCGGAAATGCCTCATCACG	Reverse	
FAS	J03860	CTATCGACACAGCCTGCTCCT	Forward	107
		CAGAATGTTGACCCCTCCTACC	Reverse	
ACC α	NM_205505	AATGGCAGCTTTGGAGGTGT	Forward	136
		TCTGTTTGGGTGGGAGGTG	Reverse	
CPT1	AY675193	GGAGAACCCAAGTGAAAGTAATGAA	Forward	135
		GAAACGACATAAAGGCAGAACAGA	Reverse	
SREBP-1	AY029224	GAGGAAGGCCATCGAGTACA	Forward	392
		GGAAGACAAAGGCACAGAGG	Reverse	
MO25 α	XM_422642	GTGGAGATGTCGACGTTTGA	Forward	439
		ATTGTTTCATCCTCGGTCTG	Reverse	
MO25 β	NM_0010006272	CTGGAATCTGCTTCCCATC	Forward	252
		GCAGGATTTTGTGCGATTT	Reverse	
STRAD β	XM_421938	CTCCATTTCATGGCCTATGGT	Forward	558
		TCGTCATTGTGCGTGCATA	Reverse	

18s = 18s rRNA; LKB1 = Liver kinase B1; AMPK = AMP-activated protein kinase; NPY = Neuropeptide Y; AgRP = Agouti-related peptide; CRH = Corticotropin-releasing hormone; POMC = Pro-opiomelanocortin; FAS = Fatty acid synthase; ACC α = Acetyl-CoA carboxylase α ; CPT-1 = Carnitine palmitoyltransferase 1; SREBP-1 = Sterol regulatory element binding protein-1; MO25 = Mouse protein 25; STRAD = STE20 related adaptor protein; STRAD β .

Therefore, all gene transcription results are reported as the n-fold difference relative to the calibrator. Specificity of the amplification product was verified with melt curve.

Statistical analysis

Data are presented as means \pm SEM. Homogeneity of variances among groups was confirmed using Bartlett's test.

All data were subjected to one-way ANOVA analysis testing the main effect of the treatment ((Version 8e, SAS Institute, Cary, NC, USA). When the main effect of treatment was significant, differences between means were assessed by Duncan's multiple range analysis. $p < 0.05$ was considered significant.

RESULTS

Expression of AMPK and related genes of hypothalamus during fasting and refeeding

Fasting for 24 h increased the gene expression of AMPK α 2, AMPK β 1, AMPK β 2, AMPK γ 1, STRAD β and MO25 α ($p < 0.05$), but showed no significant effects on the gene expression of AMPK α 1, AMPK γ 2 and LKB1 ($p > 0.05$). The gene expression of MO25 β was decreased ($p < 0.05$) by

24 h fasting. Refeeding for 24 h restored the gene expression of AMPK α 2, AMPK γ 1 and MO25 α to the control levels (Figure 1).

Expression of feed intake regulation peptides of hypothalamus during fasting and refeeding

Fasting for 24 h had no significant ($p > 0.05$) effects on gene expression of NPY. However, mRNA levels of POMC, CRH and ghrelin were significantly ($p < 0.05$) lower during




Figure 1. Effect of fasting and refeeding on mRNA levels of AMP-activated protein kinase AMPK α 1 (A), AMPK α 2 (A), AMPK β 1 (B), AMPK β 2 (B), AMPK γ 1 (C), AMPK γ 2 (C), LKB1 (D), STRAD β (D), MO25 α (E), MO25 β (E) in hypothalamus of broiler chicks. Values are means \pm SEM ($n = 4$). ^{a, b} Means with different letters differ significantly ($p < 0.05$).

fasting compared to fed control. AgRP gene expression was significantly increased ($p < 0.05$) by 24 h fasting. After 24 h of refeeding, mRNA levels of POMC and ghrelin were significantly increased ($p < 0.05$). Refeeding decreased mRNA levels of NPY and restored gene expression of AgRP to control levels ($p < 0.05$, Figure 2).

Expression of lipid metabolism-related genes in hypothalamus during fasting and refeeding

Fasting for 24 h significantly ($p < 0.05$) decreased ACC, FAS, CPT-1 and SREBP-1 mRNA levels compared to *ad libitum* fed controls (Figure 3). Refeeding for 24 h restored ACC, FAS, CPT-1 and SREBP-1 gene expression; in some cases (ACC, FAS, SREBP-1), increased expression levels were significantly above the control levels, which indicated that there was a classical 'overshoot' response ($p < 0.05$).

DISCUSSION

Effect of 24 h fasting on gene expression of AMPK subunits in chick's hypothalamus

AMPK has emerged as a nutrient and glucose sensor in the hypothalamus (Momcilovic et al., 2006). It is demonstrated that hypothalamic AMPK activity increases during fasting and decreases during refeeding in mammals (Culmsee, 2001; Minokoshi et al., 2004). AMPK switches on the oxidative metabolism of glucose and fatty acids to generate ATP, while switching off ATP consuming pathways. It achieves this metabolic switching both by direct phosphorylation of metabolic enzymes and via effects on transcription (Hardie and Hawley, 2001; Hardie et al., 2003). However the information of avian AMPK remains obscure, especially its definite roles in avian energy and appetite homeostasis.

In the present study, the expression of AMPK subunits increased after fasting for 24 h, except for $\alpha 1$ and $\gamma 2$. It has been shown that AMPK subunit genes have tissue-specific expression and both alpha subunit isoforms (alpha-1 and alpha-2) were co-expressed in all tissues examined (Proszkowiec-Weglarz et al., 2006). The current results were inconsistent with some previous reports in chicken liver (Proszkowiec-Weglarz et al., 2009). Whether this




Figure 2. Effect of fasting and refeeding on mRNA levels of POMC (A), CRH (A), ghrelin (A), NPY (B) and AgRP (B) in hypothalamus of broiler chicks. Values are means \pm SEM ($n = 4$). ^{a, b} Means with different letters differ significantly ($p < 0.05$).




Figure 3. Effect of fasting and refeeding on mRNA levels of ACC (A), CPT-1 (A), FAS (B) and SREBP-1 (B) in hypothalamus of broiler chicks. Values are means \pm SEM ($n = 4$). ^{a, b} Means with different letters differ significantly ($p < 0.05$).

difference derived from the tissue specific expression of AMPK needs further investigation. Based on the present results, AMPK $\alpha 1$ was less sensitive to different nutritional conditions than AMPK $\alpha 2$ in the chick's hypothalamus. When chicks were refed, the increased gene expression of AMPK subunits were reduced except AMPK $\gamma 1$. The integration of these data implied that the different AMPK subunits did not respond to the fasting in the same way, which revealed the complexity of the function of AMPK in the chick's hypothalamus. However, phosphorylation reaction of AMPK protein(s) would be a key of the activation and inactivation of AMPK signaling and the relationships among changes in gene expression of AMPK subunits and its protein phosphorylation during fasting needs further investigation.

LKB1 is one of the AMPK upstream protein kinases (Hawley et al., 1996). Hawley et al. (2003) demonstrated that complexes of LKB1 tumor suppressor, STRAD and MO25 are upstream kinases in the AMPK cascade. In chicken, a functional LKB1/AMPK pathway, similar to the corresponding pathway in mammals has been suggested (Proszkowiec-Weglarz et al., 2006). The hypothalamus of chicken expresses high amount of LKB1, MO25 α and both isoforms of STRAD (Proszkowiec-Weglarz et al., 2006). In the present study, the gene expression of MO25 β was decreased with 24 h fasting; however, the gene expression of STRAD β and MO25 α were increased. The increased gene expression of STRAD β and MO25 α might result in the up-regulation of AMPK $\alpha 2$ in the chick's hypothalamus.

Effect of 24 h fasting on gene expression of appetite regulation peptides in chick's hypothalamus

Multiple neuron populations distributed throughout the brain influence the decision to seek and consume food (Morton et al., 2006). It is known that hypothalamic melanocortin system (HMS) has a crucial role in feeding regulatory neural circuitry. HMS is composed of two different populations of neurons, one set that expresses neuropeptides Y (NPY) and agouti-related protein (AgRP) and a second set that expresses proopiomelanocortin (POMC), a precursor containing α -melanocyte-stimulating hormone (Richards et al., 2010). In the present study fasting broiler chicks for 24 h did not change mRNA level of NPY in the hypothalamus of broiler chicks, however the mRNA level of AgRP and POMC were increased and decreased respectively (Figure 2).

It is demonstrated that AgRP increase food intake when injected into the brain (Ollmann et al., 1997). Moreover, Takahashi and Cone (2005) discovered that the firing rate of AgRP neurons is elevated in brain slices from food-deprived mice. Hahn et al. (1998) suggested that hypothalamic AgRP neuron constitute a unique cell type that is activated by fasting to stimulate food intake via a

simultaneous decrease of melanocortin. In contrast, genetic and pharmacologic evidence suggest that POMC neurons inhibit feeding by releasing α -melanocyte-stimulating hormone, a melanocortin receptor agonist (Tsujii and Bray, 1989; Yaswen et al., 1999). It is suggested that opposite trend of AgRP and POMC gene expression implies that AgRP neurons inhibit POMC neurons in ARC (Cowley et al., 2001; Roseberry et al., 2004).

In the present study, fasting resulted in a decrease in the mRNA level of ghrelin and CRH in the hypothalamus of broiler chicks. It is known that ghrelin acts as orexigenic peptide in mammals when injected centrally and peripherally (Nakazato et al., 2001; Date et al., 2002). In the case of mammals, orexigenic activity of ghrelin is mediated by activation of other orexigenic peptides such as NPY, AgRP and orexin (Toshinai et al., 2003). In neonatal chickens, intracerebroventricular (icv) administration of ghrelin potently inhibits food intake in a dose dependent manner (Khan et al., 2006). Saito et al. (2005) suggested that anorexigenic effects of central ghrelin mediated via CRH in chickens.

The potential relationship between AMPK, appetite regulation peptides and lipid metabolism related genes

All isoforms of AMPK are expressed in neuronal tissues, including these involved in the control of food intake and neuroendocrine function, such as the hypothalamus and the hindbrain (Turnley et al., 1999; Kola, 2008). Pharmacological activation of AMPK in rodent hypothalamus with 5-aminoimidazole-4-carboxamide riboside (AICAR) causes an increase in food intake (Xue and Kahn, 2006). In line with this, it has been also recently reported that AMPK $\beta 1$ KO mice reduce food intake, either under low fat (LFD) or HFD (Dzamko, 2010). Deletion of AMPK $\alpha 2$ in AgRP neurons led to the development of an age-dependent lean phenotype (Claret et al., 2007; Lim et al., 2010). In the present study, the enhanced gene expression of $\alpha 2$, $\beta 1$, $\beta 2$ and $\gamma 1$ isoforms of AMPK was accompanied by a reduction in the mRNA levels of POMC and ghrelin and an enhancement of the AgRP mRNA levels in the hypothalamus of broiler chicks after 24 h fasting. Based on these results, it was speculated that AMPK might be involved in the appetite regulation in the hypothalamus of the chicks. AMPK might have enhanced the appetite of chicks via increasing the orexigenic gene expression of AgRP and decreasing the anorexigenic gene expression of POMC in chicks with 24 h fasting.

The level of metabolic flux through the fatty acid biosynthetic pathway in hypothalamic neurons as regulated by AMPK determines the levels of two key metabolites, malonyl-CoA, and long chain fatty acyl-CoA, which in turn lead to changes in feed intake and energy expenditure by altering the expression of orexigenic and anorexigenic

neuropeptides in melanocortin system neurons (Lam et al., 2005; Lane et al., 2005; He et al., 2006). AMPK can regulate the transcription and expression of FAS and ACC α in hypothalamus of mammals and avian species (Xue and Kahn, 2006; Xu et al., 2011a; Xu et al., 2011b). The AMPK inhibits the activity of ACC, the rate-limiting enzyme involved in the production of malonyl-CoA used for fatty acyl-CoA biosynthesis, and causes a reduction in this reaction, which stimulates CPT1 and reduces the flux of substrates in the fatty acid anabolic pathway (Carling et al., 2008; Lage et al., 2008). In this study, although we anticipated an increase in the mRNA level of CPT1, however, 24 h fasting significantly decreased CPT1 gene expression in the hypothalamus of broiler chicks. The underlying mechanism needs further investigation.

In our experiment, the mRNA levels of FAS and SREBP-1 were reduced after 24 h fasting and refeeding caused an overshooting of FAS and SREBP-1 gene expression. FAS inhibitor reduces food intake in mammals and chickens (Dridi et al., 2006). Feed restriction decreased gene expression of FAS in an AMPK-dependent manner in mammals (López et al., 2008) and chickens (Dridi et al., 2006; Proszkowiec-Weglarz et al., 2006). AMPK can directly decrease FAS gene transcription via regulating the gene expression of SREBP-1 in mammals and avian (Foretz et al., 2005; Shaw et al., 2005; Proszkowiec-Weglarz et al., 2009). Bennett et al. (2008) recently reported that the lipogenic enzyme gene expression ‘overshoot’ in response to refeeding with a high carbohydrate diet in mice following a fast, which was attributable to enhanced binding of SREBP-1c to SRE sites located in lipogenic enzyme target gene promoters, as well as increased co-regulatory protein recruitment. In line with previous reports, the present findings revealed that the AMPK/SREBP-1/FAS signal pathway might be involved in the 24 h fasting induced appetite regulation in chicks.

In conclusion, 24 h fasting altered gene expression of AMPK subunits, appetite regulation peptides and lipometabolism related factors in chick’s hypothalamus; the hypothalamic FAS signal pathway might be involved in the AMPK regulated energy homeostasis and/or appetite regulation in poultry.

ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (31172226) and the Yangtze River Scholar and Innovation Research Team Development Program (Grant no. IRT0945).

REFERENCES

Bass, A. F., J. Boudeau, G. P. Sapkota, L. Smit, R. Medema, N. A.

- Morrice, D. R. Alessi and H. C. Clevers. 2003. Activation of the tumor suppressor kinase LKB1 by the STE20-like pseudokinase STRAD. *EMBO J.* 22:3062-3072.
- Bennett, M., Y. Seo, S. Datta, D. Shin and T. Osborne. 2008. Selective binding of sterol regulatory element-binding protein isoforms and co-regulatory proteins to promoters for lipid metabolic genes in liver. *J. Biol. Chem.* 283:15628-15637.
- Boudeau, J., A. F. Bass, M. Deak, N. A. Morrice, A. Kieloch, M. Schutkowski, A. R. Prescott, H. C. Clevers and D. R. Alessi. 2003. MO25 α/β interact with STRAD α/β enhancing their ability to bind, activate and localize LKB1 in the cytoplasm. *EMBO J.* 22:5102-5114.
- Carling, D., M. Sanders and A. Woods. 2008. The regulation of AMP-activated protein kinase by upstream kinases. *Int. J. Obes. (Lond).* 32:S55-S59.
- Cheung, P., I. Salt, S. Davies, D. Hardie and D. Carling. 2000. Characterization of AMP-activated protein kinase gamma-subunit isoforms and their role in AMP binding. *Biochem. J.* 346:659-669.
- Claret, M., M. A. Smith, R. L. Batterham, C. Selman, A. I. Choudhury, L. G. Fryer, M. Clements, H. Al-Qassab, H. Heffron, A. W. Xu, J. R. Speakman, G. S. Barsh, B. Viollet, S. Vaulont, M. L. Ashford, D. Carling and D. J. Withers. 2007. AMPK is essential for energy homeostasis regulation and glucose sensing by POMC and AgRP neurons. *J. Clin. Invest.* 117:2325-2336.
- Close, B., K. Banister, V. Baumans, E. M. Bernoth, N. Bromage, J. Bunyan, W. Erhardt, P. Flecknell, N. Gregory, H. Hackbarth, D. Morton and C. Warwick. 1997. Recommendations for euthanasia of experimental animals: Part 2. DGXT of the European Commission. *Lab. Anim.* 31:1-32.
- Corton, J. M., J. G. Gillespie and D. G. Hardie. 1994. Role of the AMP-activated protein kinase in the cellular stress response. *Curr. Biol.* 4:315-324.
- Cowley, M., J. L. Smart, M. Rubinstein, M. G. Cerdán, S. Diano, T. L. Horvath, R. D. Cone and M. L. Low. 2001. Leptin activates anorexigenic POMC neurons through a neural network in the arcuate nucleus. *Nature* 411:480-484.
- da Silva Xavier, G., I. Leclerc, I. P. Salt, B. Doiron, D. G. Hardie, A. Kahn and G. A. Rutter. 2000. Role of AMP-activated protein kinase in the regulation by glucose of islet beta cell gene expression. *Proc. Natl. Acad. Sci. USA.* 97:4023-4028.
- Date, Y., N. Murakami, K. Toshinai, S. Matsukura, A. Nijima, H. Matsuo, K. Kangawa and M. Nakazato. 2002. The role of the gastric afferent vagal nerve in ghrelin-induced feeding and growth hormone secretion in rats. *Gastroenterology* 123:1120-1128.
- Dridi, S., C. Ververken, F. B. Hillgartner, A. Lutgarde, E. Van der Gucht, L. Cnops, E. Decuypere and J. Buyse. 2006. FAS inhibitor cerulenin reduces food intake and melanocortin receptor gene expression without modulating the other (an)orexigenic neuropeptides in chickens. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 291:R138-R147.
- Dzambo, N., B. J. van Denderen, A. L. Hevener, S. B. Jørgensen, J. Honeyman, S. Galic, Z. P. Chen, M. J. Watt, D. J. Campbell, G. R. Steinberg and B. E. Kemp. 2010. AMPK beta1 deletion reduces appetite, preventing obesity and hepatic insulin resistance. *J. Biol. Chem.* 285:115-122.
- Foretz, M., N. Ancellin, F. Andreelli, Y. Saintillan, P. Grondin, A.

- Kahn, B. Thorens, S. Vaulont and B. Viollet. 2005. Short-term overexpression of a constitutively active form of AMP-activated protein kinase in the liver leads to mild hypoglycemia and fatty liver. *Diabetes* 54:1331-1339.
- Hahn, T., J. Breininger, D. Baskin and M. Schwartz. 1998. Coexpression of Agrp and NPY in fasting-activated hypothalamic neurons. *Nat. Neurosci.* 1:271-272.
- Hardie, D. and S. Hawley. 2001. AMP-activated protein kinase: the energy charge hypothesis revisited. *Bioessays* 23:1112-1119.
- Hardie, D. G., J. W. Scott, D. A. Pan and E. R. Hudson. 2003. Management of cellular energy by the AMP-activated protein kinase system. *FEBS Lett.* 546:113-120.
- Hawley, S. A., M. Davison, A. Woods, S. P. Davies, R. K. Beri, D. Carling and D. G. Hardie. 1996. Characterization of the AMP-activated protein kinase from rat liver and identification of threonine-172 as the major site at which it phosphorylates AMP-activated protein kinase. *J. Biol. Chem.* 271:27879-27887.
- Hawley, S. A., D. A. Pan, K. J. Mustard, L. Ross, J. Bain, A. M. Edelman, B. G. Frenguelli and D. G. Hardie. 2005. Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab.* 2:9-19.
- Hemminki, A., D. Markie, I. Tomlinson, E. Avizienyte, S. Roth, A. Loukola, G. Bignell, W. Warren, M. Aminoff, P. Höglund, H. Järvinen, P. Kristo, K. Pelin, M. Ridanpää, R. Salovaara, T. Toro, W. Bodmer, S. Olschwang, A. S. Olsen, M. R. Stratton, A. de la Chapelle and L. A. Aaltonen. 1998. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature* 391:184-187.
- He, W., T. Lam, S. Obici and L. Rossetti. 2006. Molecular disruption of hypothalamic nutrient sensing induces obesity. *Nat. Neurosci.* 9:227-233.
- Hong, S. P., F. C. Leiper, A. Woods, D. Carling and M. Carlson. 2003. Activation of yeast Snf1 and mammalian AMP-activated protein kinase by upstream kinases. *Proc. Natl. Acad. Sci. USA.* 100:8839-8843.
- Hurley, R., K. A. Anderson, J. M. Franzone, B. E. Kemp, A. R. Means and L. A. Witters. 2005. The Ca²⁺/calmodulin-dependent protein kinase kinases are AMP-activated protein kinase kinases. *J. Biol. Chem.* 280:29060-29066.
- Jenne, D. E., H. Reimann, J. Nezu, W. Friedel, S. Loff, R. Jeschke, O. Müller, W. Back and M. Zimmer. 1998. Peutz-Jeghers syndrome is caused by mutation in a novel serine threonine kinase. *Nat. Genet.* 18:38-43.
- Khan, M., K. Dodo, K. Yahata, S. Nishimoto, H. Ueda, T. Taneike, T. Kitazawa and Y. Hosaka. 2006. Intracerebroventricular administration of growth hormone releasing peptide-6 (GHRP-6) inhibits food intake, but not food retention of crop and stomach in neonatal chicks. *J. Poult. Sci.* 43:35-40.
- Kola, B. 2008. Role of AMP-activated protein kinase in the control of appetite. *J. Neuroendocrinol.* 20:942-951.
- Kola, B., M. Boscaro, G. A. Rutter, A. B. Grossman and M. Korbonits. 2006. Expanding role of AMPK in endocrinology. *Trends Endocrinol. Metab.* 17:205-215.
- Lam, T. K., G. J. Schwartz and L. Rossetti. 2005. Hypothalamic sensing of fatty acids. *Nat. Neurosci.* 8:579-584.
- Lane, M. D., Z. Hu, S. H. Cha, Y. Dai, M. Wolfgang and A. Sidhaye. 2005. Role of malonyl-CoA in the hypothalamic control of food intake and energy expenditure. *Biochem. Soc. Trans.* 33:1063-1067.
- López, M., R. Lage, A. K. Saha, D. Pérez-Tilve, M. J. Vázquez, L. Varela, S. Sangiao-Alvarellos, S. Tovar, K. Raghay, S. Rodríguez-Cuenca, R. M. Deoliveira, T. Castañeda, R. Datta, J. Z. Dong, M. Culler, M. W. Sleeman, C. V. Álvarez, R. Gallego, C. J. Lelliott, D. Carling, M. H. Tschöp, C. Diéguez and A. Vidal-Puig. 2008. Hypothalamic fatty acid metabolism mediates the orexigenic action of ghrelin. *Cell Metab.* 7:389-399.
- Lage, R., C. Diéguez, A. Vidal-Puig and M. López. 2008. AMPK: a metabolic gauge regulating whole-body energy homeostasis. *Trends Mol. Med.* 14:539-549.
- Lim, C. T., B. Kola and M. Korbonits. 2010. AMPK as a mediator of hormonal signalling. *J. Mol. Endocrinol.* 44:87-97.
- Livak, K. J. and T. D. Schmittgen. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $[\Delta\Delta]CT$ method. *Methods* 25:402-408.
- Marsin, A. S., L. Bertrand, M. H. Rider, J. Deprez, C. Beaufoye, M. F. Vincent, G. Van den Berghe, D. Carling and L. Hue. 2000. Phosphorylation and activation of heart PFK-2 by AMPK has a role in the stimulation of glycolysis during ischaemia. *Curr. Biol.* 10:1247-1255.
- Minokoshi, Y., T. Alquier, N. Furukawa, Y. B. Kim, A. Lee, B. Xue, J. Mu, F. Foufelle, P. Ferre, M. J. Birnbaum, B. J. Stuck and B. B. Kahn. 2004. AMP-kinase regulates food intake by responding to hormonal and nutrient signals in the hypothalamus. *Nature* 428:569-574.
- Mitchellhill, K. I., D. Stapleton, G. Gao, C. House, B. Michell, F. Katsis, L. A. Witters and B. E. Kemp. 1994. Mammalian AMP-activated protein kinase shares structural and functional homology with the catalytic domain of yeast Snf1 protein kinase. *J. Biol. Chem.* 269:2361-2364.
- Momcilovic, M., S. P. Hong and M. Carlson. 2006. Mammalian TAK1 activates Snf1 protein kinase in yeast and phosphorylates AMP-activated protein kinase *in vitro*. *J. Biol. Chem.* 281:25336-25343.
- Morton, G. J., D. E. Cummings, D. G. Baskin, G. S. Barsh and M. W. Schwartz. 2006. Central nervous system control of food intake and body weight. *Nature* 443:289-295.
- Mu, J., J. T. Brozinick, O. Valladares, M. Bucan and M. J. Birnbaum. 2001. A role for AMP-activated protein kinase in contraction- and hypoxia-regulated glucose transport in skeletal muscle. *Mol. Cell* 7:1085-1094.
- Musi, N., N. Fujii, M. F. Hirshman, I. Ekberg, S. Fröberg, O. Ljungqvist, A. Thorell and L. J. Goodyear. 2001. AMP-activated protein kinase (AMPK) is activated in muscle of subjects with type 2 diabetes during exercise. *Diabetes* 50:921-927.
- Nakazato, M., N. Murakami, Y. Date, M. Kojima, H. Matsuo, K. Kangawa and S. Matsukura. 2001. A role for ghrelin in the central regulation of feeding. *Nature* 409:194-198.
- Obici, S., Z. Feng, A. Arduini, R. Conti and L. Rossetti. 2003. Inhibition of hypothalamic carnitine palmitoyltransferase-1 decreases food intake and glucose production. *Nat. Med.* 9:756-761.
- Olfert, E. D., B. M. Cross and A. A. McWilliam. 1993. Guide to the care and use of experimental animals. Canadian Council on

- Animal Care, 1000-151 Slater Street, Ottawa, Ontario K1P 5H3.
- Ollmann, M. M., B. D. Wilson, Y. K. Yang, J. A. Kerns, Y. Chen, I. Gantz and G. S. Barsh. 1997. Antagonism of central melanocortin receptors *in vitro* and *in vivo* by agouti-related protein. *Science* 278:135-138.
- Proszkowiec-Weglarz, M., M. P. Richards, B. D. Humphrey, R. W. Rosebrough and J. P. McMurtry. 2009. AMP-activated protein kinase and carbohydrate response element binding protein: a study of two potential regulatory factors in the hepatic lipogenic program of broiler chickens. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol.* 154:68-79.
- Proszkowiec-Weglarz, M., M. P. Richards, R. Ramachandran and J. McMurtry. 2006. Characterization of the AMP-activated protein kinase pathway in chickens. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol.* 143:92-106.
- Proszkowiec-Weglarz, M. and M. P. Richards. 2009. Expression and activity of the 5'-adenosine monophosphate-activated protein kinase pathway in selected tissues during chicken embryonic development. *Poult. Sci.* 88:159-178.
- Richards, M. P., R. W. Rosebrough, C. N. Coon and J. P. McMurtry. 2010. Feed intake regulation for the female broiler breeder: In theory and in practice. *J. Appl. Poult. Res.* 19:182-193.
- Richards, M. P. and M. Proszkowiec-Weglarz. 2007. Mechanisms regulating feed intake, energy expenditure, and body weight in poultry. *Poult. Sci.* 86:1478-1490.
- Roseberry, A. G., H. Liu, A. C. Jackson, X. Cai and J. M. Friedman. 2004. Neuropeptide Y-mediated inhibition of proopiomelanocortin neurons in the arcuate nucleus shows enhanced desensitization in *ob/ob* mice. *Neuron* 41:711-722.
- Saito, E., H. Kaiya, T. Tachibana, S. Tomonaga, D. Denbow, K. Kangawa and M. Furuse. 2005. Inhibitory effect of ghrelin on food intake is mediated by the corticotropin-releasing factor system in neonatal chickens. *Regul. Pept.* 125:201-208.
- Scott, J. W., S. A. Hawley, K. A. Green, M. Anis, G. Stewart, G. A. Scullion, D. G. Norman and D. G. Hardie. 2004. CBS domains form energy-sensing modules whose binding of adenosine ligands is disrupted by disease mutations. *J. Clin. Invest.* 113: 274-284.
- Shaw, R. J., K. A. Lamia, D. Vasquez, S. H. Koo, N. Bardeesy, R. A. DePinho, M. Montminy and L. C. Cantley. 2005. The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. *Science* 310:1642-1646.
- Stapleton, D., K. I. Mitchelhill, G. Gao, J. Widmer, B. J. Michell, T. The, C. M. House, C. S. Fernandez, T. Cox, L. A. Witters and B. E. Kemp. 1996. Mammalian AMP-activated protein kinase subfamily. *J. Biol. Chem.* 271:611-614.
- Stapleton, D., E. Woollatt, K. I. Mitchelhill, J. K. Nicholl, C. S. Fernandez, B. J. Michell, L. A. Witters, D. A. Power, G. R. Sutherland and B. E. Kemp. 1997. AMP-activated protein kinase isoenzyme family: subunit structure and chromosomal location. *FEBS Lett.* 409:452-456.
- Takahashi, K. A. and R. D. Cone. 2005. Fasting induces a large, leptin-dependent increase in the intrinsic action potential frequency of orexigenic arcuate nucleus neuropeptide Y/Agouti-related protein neurons. *Endocrinology* 146:1043-1047.
- Toshinai, K., Y. Date, N. Murakami, M. Shimada, M. S. Mondal, T. Shimbara, J. L. Guan, Q. P. Wang, H. Funahashi, T. Sakurai, S. Shioda, S. Matsukura, K. Kangawa and M. Nakazato. 2003. Ghrelin-induced food intake is mediated via the orexin pathway. *Endocrinology* 144:1506-1512.
- Tsujii, S., and G. A. Bray. 1989. Acetylation alters the feeding response to MSH and beta-endorphin. *Brain Res. Bull.* 23: 165-169.
- Turnley, A. M., D. Stapleton, R. J. Mann, L. A. Witters, B. E. Kemp and P. F. Bartlett. 1999. Cellular distribution and developmental expression of AMP-activated protein kinase isoforms in mouse central nervous system. *J. Neurochem.* 72: 1707-1716.
- Xue, B. and B. Kahn. 2006. AMPK integrates nutrient and hormonal signals to regulate food intake and energy balance through effects in the hypothalamus and peripheral tissues. *J. Physiol.* 574:73-83.
- Xu, P., C. J. Denbow, N. Meiri and D. M. Denbow. 2011. Fasting of 3-day-old chickens leads to changes in histone H3 methylation status. *Physiol. Behav.* 105:276-282.
- Xu, P., P. B. Siegel and D. M. Denbow. 2011. Genetic selection for body weight in chickens has altered responses of the brain's AMPK system to food intake regulation effect of ghrelin, but not obestatin. *Behav. Brain Res.* 221:216-226.
- Yaswen, L., N. Diehl, M. Brennan and U. Hochgeschwender. 1999. Obesity in the mouse model of pro-opiomelanocortin deficiency responds to peripheral melanocortin. *Nat. Med.* 5: 1066-1070.