

## BOUNDED WEAK SOLUTION FOR THE HAMILTONIAN SYSTEM

Q-HEUNG CHOI AND TACKSUN JUNG\*

ABSTRACT. We investigate the bounded weak solutions for the Hamiltonian system with bounded nonlinearity decaying at the origin and periodic condition. We get a theorem which shows the existence of the bounded weak periodic solution for this system. We obtain this result by using variational method, critical point theory for indefinite functional.

### 1. introduction

Let  $G(t, z(t))$  be a  $C^2$  function defined on  $R^1 \times R^{2n}$  which is  $2\pi$ -periodic with respect to the first variable  $t$ . In this paper we investigate the number of  $2\pi$ -periodic solutions of the following Hamiltonian system

$$(1.1) \quad \begin{aligned} \dot{p}(t) &= -G_q(t, p(t), q(t)), \\ \dot{q}(t) &= G_p(t, p(t), q(t)), \end{aligned}$$

where  $p, q \in R^n$ ,  $z = (p, q)$ . Let  $J$  be the standard symplectic structure on  $R^{2n}$ , i. e.,

$$J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix},$$

---

Received February 19, 2013. Revised March 18, 2013. Accepted March 20, 2013.  
2010 Mathematics Subject Classification: 35Q72.

Key words and phrases: Hamiltonian system, bounded nonlinearity, variational method, critical point theory for indefinite functional,  $(P.S.)_c$  condition.

This work was supported by Inha University Research Grant.

\*Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2013.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

where  $I_n$  is the  $n \times n$  identity matrix. Then (1.1) can be rewritten as

$$(1.2) \quad -J\dot{z} = G_z(t, z(t)),$$

where  $\dot{z} = \frac{dz}{dt}$  and  $G_z$  is the gradient of  $G$ . We assume that  $G \in C^2(R^1 \times R^{2n}, R^1)$  satisfies the following conditions:

(G1)  $G \in C^2(R^1 \times R^{2n}, R^1)$ ,

(G2)  $G(t, z(t)) = O(|z|^2)$  as  $|z| \rightarrow 0$ ,  $G(t, \theta) = 0$ ,  $G_z(t, \theta) = \theta$ , where  $\theta = (0, \dots, 0)$ ,

(G3) there exists  $C > 0$  such that  $|G(t, \xi)| < C \forall t \in R, \xi \in R^3$ .

Several authors ([1], [3], [4], [5] etc.) studied the nonlinear hamiltonian system. Jung and Choi ([3], [4]) considered (1.1) with nonsingular potential nonlinearity or jumping nonlinearity crossing one eigenvalue, or two eigenvalues, or several eigenvalues. Chang ([1]) proved that (1.1) has at least two nontrivial  $2\pi$ -periodic weak solutions under some asymptotic nonlinearity. Jung and Choi ([3]) proved that (1.1) has at least  $m$  weak solutions, which are geometrically distinct and nonconstant under some jumping nonlinearity.

We are looking for the weak solutions of (1.1) with the conditions (G1)-(G3). The  $2\pi$ -periodic weak solution  $z = (p, q) \in E$  of (1.1) satisfies

$$\int_0^{2\pi} (\dot{z} - JG_z(t, z(t))) \cdot Jw dt = 0 \quad \text{for all } w \in E,$$

i. e.,

$$\int_0^{2\pi} [(\dot{p} + G_q(t, z(t))) \cdot \psi - (\dot{q} - G_p(t, z(t))) \cdot \phi] dt = 0$$

for all  $\zeta = (\phi, \psi) \in E$ ,

where  $E$  is introduced in section 2.

Our main result is as follows:

**THEOREM 1.1.** *Assume that  $G$  satisfies the conditions (G1) – (G3). Then system (1.1) has at least one bounded  $2\pi$ -periodic solution.*

For the proof of our main result we approach the variational method and apply the critical point theory to indefinite functional. The outline of the proof of Theorem 1.1 is as follows: In Section 2, we introduce the perturbed operator  $A_\epsilon$  such that  $A_\epsilon^{-1}$  is a compact operator, and the associated functional  $I(z)$  corresponding to the operator  $A_\epsilon$ , prove that

$I(z)$  satisfies *Fréchet* differentiability, and state the critical point theorem for indefinite functional. In section 3, we show that the associated functional  $I(z)$  satisfies the geometrical assumptions of the critical point theorem for indefinite functional, and prove Theorem 1.1.

## 2. Compact operator and variational approach

Let  $L^2([0, 2\pi], R^{2n})$  denote the set of  $2n$ -tuples of the square integrable  $2\pi$ -periodic functions and choose  $z \in L^2([0, 2\pi], R^{2n})$ . Then it has a Fourier expansion  $z(t) = \sum_{k=-\infty}^{k=+\infty} a_k e^{ikt}$ , with  $a_k = \frac{1}{2\pi} \int_0^{2\pi} z(t) e^{-ikt} dt \in C^{2n}$ ,  $a_{-k} = \bar{a}_k$  and  $\sum_{k \in \mathbb{Z}} |a_k|^2 < \infty$ . Let

$$A : z(t) \mapsto -J\dot{z}(t)$$

with domain

$$\begin{aligned} D(A) &= \{z(t) \in H^1([0, 2\pi], R^{2n}) \mid z(0) = z(2\pi)\} \\ &= \{z(t) \in L^2([0, 2\pi], R^{2n}) \mid \sum_{k \in \mathbb{Z}} (\epsilon + |k|)^2 |a_k|^2 < +\infty\}, \end{aligned}$$

where  $\epsilon$  is a positive small number. Then  $A$  is a self-adjoint operator. Let  $\{M_\lambda\}$  be the spectral resolution of  $A$ , and let  $\alpha$  be a positive number such that  $\alpha \notin \sigma(A)$  and  $[-\alpha, \alpha]$  contains only one element  $0$  of  $\sigma(A)$ . Let

$$P_0 = \int_{-\alpha}^{\alpha} dM_\lambda, \quad P_+ = \int_{\alpha}^{+\infty} dM_\lambda, \quad P_- = \int_{-\infty}^{-\alpha} dM_\lambda.$$

Let

$$\begin{aligned} L_0 &= P_0 L^2([0, 2\pi], R^{2n}), \\ L_+ &= P_+ L^2([0, 2\pi], R^{2n}), \\ L_- &= P_- L^2([0, 2\pi], R^{2n}). \end{aligned}$$

For each  $u \in L^2([0, 2\pi], R^{2n})$ , we have the decomposition

$$u = u_0 + u_+ + u_-,$$

where  $u_0 \in L_0$ ,  $u_+ \in L_+$ ,  $u_- \in L_-$ . According to  $A$ , there exists a small number  $\epsilon > 0$  such that  $-\epsilon \notin \sigma(A)$ . Let us define the space  $E$  as follows:

$$E = D(|A|^{\frac{1}{2}}) = \{z \in L^2([0, 2\pi], R^{2n}) \mid \sum_{k \in \mathbb{Z}} (\epsilon + |k|) |a_k|^2 < \infty\}$$

with the scalar product

$$(z, w)_E = \epsilon(z, w)_{L^2} + (|A|^{\frac{1}{2}}z, |A|^{\frac{1}{2}}w)_{L^2}$$

and the norm

$$\|z\| = (z, z)_E^{\frac{1}{2}} = \left( \sum_{k \in \mathbb{Z}} (\epsilon + |k|) |a_k|^2 \right)^{\frac{1}{2}}.$$

The space  $E$  endowed with this norm is a real Hilbert space continuously embedded in  $L^2([0, 2\pi], R^{2n})$ . The scalar product in  $L^2$  naturally extends as the duality pairing between  $E$  and  $E' = W^{-\frac{1}{2}, 2}([0, 2\pi], R^{2n})$ . We note that the operator  $(\epsilon + |A|)^{-1}$  is a compact linear operator from  $L^2([0, 2\pi], R^{2n})$  to  $E$  such that

$$((\epsilon + |A|)^{-1}w, z)_E = \int_0^{2\pi} (w(t), z(t)) dt.$$

Let

$$A_\epsilon = \epsilon I + A.$$

Let

$$E_0 = |A_\epsilon|^{-\frac{1}{2}}L_0, \quad E_+ = |A_\epsilon|^{-\frac{1}{2}}L_+, \quad E_- = |A_\epsilon|^{-\frac{1}{2}}L_-.$$

Then  $E = E_0 \oplus E_+ \oplus E_-$  and for  $z \in E$ ,  $z$  has the decomposition  $z = z_0 + z_+ + z_- \in E$ , where

$$z_0 = |A_\epsilon|^{-\frac{1}{2}}u_0, \quad z_+ = |A_\epsilon|^{-\frac{1}{2}}u_+, \quad z_- = |A_\epsilon|^{-\frac{1}{2}}u_-.$$

Thus we have

$$\|z_0\|_{E_0} = \|u_0\|_{L_0}, \quad \|z_+\|_{E_+} = \|u_+\|_{L_+}, \quad \|z_-\|_{E_-} = \|u_-\|_{L_-}$$

and that  $E_0, E_+, E_-$  are isomorphic to  $L_0, L_+, L_-$ , respectively. The associated functional of (1.2) on  $E$  is as follows:

$$\begin{aligned} (2.1) \quad I(z) &= \frac{1}{2} (\|A_\epsilon^{\frac{1}{2}}z_+\|_{L^2}^2 + \|A_\epsilon^{\frac{1}{2}}M_+z_0\|^2 - \|(-A_\epsilon)^{\frac{1}{2}}z_-\|_{L^2}^2 \\ &\quad - \|(-A_\epsilon)^{\frac{1}{2}}M_-z_0\|^2) - \psi_\epsilon(z), \\ &= \frac{1}{2} (\|u_+\|^2 + \|M_+u_0\|^2 - \|M_-u_0\|^2 - \|u_-\|^2) - \psi_\epsilon(z), \end{aligned}$$

where  $\psi_\epsilon(z) = \psi(z) + \frac{\epsilon}{2}\|z\|_{L^2}^2$ ,  $\psi(z) = \int_0^{2\pi} G(t, z(t)) dt$ . Let

$$F(z) = G_z(t, z(t)).$$

By  $G \in C^2$ ,  $\psi(z) = \int_0^{2\pi} G(t, z(t))dt \in C^2(S^1 \times D, R^1)$ . Let

$$F_\epsilon(z) = \epsilon I + F(z) = \epsilon I + G_z(t, z(t)).$$

Then (1.2) can be rewritten as

$$(2.2) \quad A_\epsilon(z) = F_\epsilon(z).$$

The Euler equation of the functional  $I(z)$  is the system

$$(2.3) \quad u_+ = |A_\epsilon|^{-\frac{1}{2}} P_+ F_\epsilon(z),$$

$$(2.4) \quad u_- = -|A_\epsilon|^{-\frac{1}{2}} P_- F_\epsilon(z),$$

$$(2.5) \quad \begin{aligned} M_+ u_0 &= |A_\epsilon|^{-\frac{1}{2}} M_+ P_0 F_\epsilon(z) \\ M_- u_0 &= -|A_\epsilon|^{-\frac{1}{2}} M_- P_0 F_\epsilon(z). \end{aligned}$$

Thus  $z = z_0 + z_+ + z_-$  is a solution of (2.2) if and only if  $u = u_0 + u_+ + u_-$  is a critical point of  $I$ . The system (2.3)-(2.5) is reduced to

$$(2.6) \quad A_\epsilon z_+ = P_+ F_\epsilon(z_0 + z_+ + z_-) \quad \text{or} \quad z_+ = (A_\epsilon)^{-1} P_+ F_\epsilon(z_0 + z_+ + z_-),$$

$$(2.7) \quad A_\epsilon z_- = P_- F_\epsilon(z_0 + z_+ + z_-) \quad \text{or} \quad z_- = (A_\epsilon)^{-1} P_- F_\epsilon(z_0 + z_+ + z_-),$$

$$(2.8) \quad \begin{aligned} A_\epsilon M_+ z_0 &= M_+ P_0 F_\epsilon(z_0 + z_+ + z_-), \\ A_\epsilon M_- z_0 &= M_- P_0 F_\epsilon(z_0 + z_+ + z_-). \end{aligned}$$

By the following Lemma 2.1, the weak solutions of (1.2) coincide with the critical points of the functional  $I(z)$ .

**LEMMA 2.1.** *Assume that  $G$  satisfies the conditions (G1) – (G3). Then  $I(z)$  is continuous and Fréchet differentiable in  $E$  with Fréchet derivative*

$$(2.9) \quad DI(z)w = \int_0^{2\pi} (A_\epsilon z - F_\epsilon(z)) \cdot w dt \quad \text{for all } w \in E,$$

Moreover  $DI \in C$ . That is,  $I \in C^1$ .

*Proof.* First we prove that  $I(z) = \int_0^{2\pi} [\frac{1}{2}A_\epsilon z - G(t, z(t)) - \frac{\epsilon}{2}z^2]dt$  is continuous and *Fréchet* differentiable in  $E$ . For  $z, w \in E$ ,

$$\begin{aligned} & |I(z+w) - I(z)| \\ &= \left| \int_0^{2\pi} \frac{1}{2}A_\epsilon(z+w) \cdot (z+w)dt - \int_0^{2\pi} [G(t, z+w) + \frac{\epsilon}{2}(z+w)^2]dt \right. \\ &\quad \left. - \int_0^{2\pi} \frac{1}{2}A_\epsilon(z) \cdot zdt + \int_0^{2\pi} [G(t, z) + \frac{\epsilon}{2}z^2]dt \right. \\ &= \left| \int_0^{2\pi} \frac{1}{2}[A_\epsilon(z) \cdot w + A_\epsilon(w) \cdot z + A_\epsilon(w) \cdot w]dt \right. \\ &\quad \left. - \int_0^{2\pi} [G(t, z+w) - G(t, z) + \frac{\epsilon}{2}(2z \cdot w + w^2)]dt \right|. \end{aligned}$$

We have

$$\begin{aligned} (2.10) \quad & \left| \int_0^{2\pi} [G(t, z+w) - G(t, z)]dt \right| \\ & \leq \left| \int_0^{2\pi} [G_z(t, z(t)) \cdot w + O(\|w\|_{R^{2n}})]dt \right| = O(\|w\|_{R^{2n}}). \end{aligned}$$

Thus we have

$$|I(z+w) - I(z)| = O(\|w\|_{R^{2n}}).$$

Next we shall prove that  $I(z)$  is *Fréchet* differentiable in  $E$ . For  $z, w \in E$ ,

$$\begin{aligned} & |I(z+w) - I(z) - DI(z)w| \\ &= \left| \int_0^{2\pi} \frac{1}{2}A_\epsilon(z+w) \cdot (z+w)dt - \int_0^{2\pi} [G(t, z+w) + \frac{\epsilon}{2}(z+w)^2]dt \right. \\ &\quad \left. - \int_0^{2\pi} \frac{1}{2}A_\epsilon(z) \cdot zdt + \int_0^{2\pi} [G(t, z) + \frac{\epsilon}{2}z^2]dt \right. \\ &\quad \left. - \int_0^{2\pi} A_\epsilon(z) \cdot wdt + \int_0^{2\pi} [G_z(t, z) + \epsilon z \cdot w]dt \right. \\ &= \left| \int_0^{2\pi} \frac{1}{2}[A_\epsilon(w) \cdot zdt + A_\epsilon(w) \cdot w]dt \right. \\ &\quad \left. - \int_0^{2\pi} [G(t, z+w) - G(t, z) - G_z(t, z) + \frac{\epsilon}{2}w^2]dt \right|. \end{aligned}$$

By (2.10), we have

$$\int_0^{2\pi} [G(t, z + w) - G(t, z) - G_z(t, z)w] dt = O(\|w\|_{R^{2n}}).$$

Thus

$$|I(z + w) - I(z) - DI(z)w| = O(\|w\|_{R^{2n}}).$$

□

Now, we recall the critical point theorem for the indefinite functional (cf. [2]).

Let

$$B_r = \{u \in E \mid \|u\| \leq r\},$$

$$\partial B_r = \{u \in E \mid \|u\| = r\}.$$

**THEOREM 2.1.** *Critical point theorem for the indefinite functional*  
Let  $E$  be a real Hilbert space with  $E = E_1 \oplus E_2$  and  $E_2 = E_1^\perp$ . Suppose that  $I \in C^1(E, R)$ , satisfies (PS), and

(I1)  $I(u) = \frac{1}{2}(Lu, u) + bu$ , where  $Lu = L_1P_1u + L_2P_2u$  and  $L_i : E_i \rightarrow E_i$  is bounded and self adjoint,  $i = 1, 2$ ,

(I2)  $b'$  is compact, and

(I3) there exists a subspace  $\tilde{E} \subset E$  and sets  $S \subset E$ ,  $Q \subset \tilde{E}$  and constants  $\alpha > \omega$  such that

(i)  $S \subset E_1$  and  $I|_S \geq \alpha$ ,

(ii)  $Q$  is bounded and  $I|_{\partial Q} \leq \omega$ ,

(iii)  $S$  and  $\partial Q$  link.

Then  $I$  possesses a critical value  $c \geq \alpha$ .

### 3. Proof of Theorem 1.1

We shall show that the functional  $I(z)$  satisfies the geometric assumptions of the critical point theorem for indefinite functional.

**LEMMA 3.1.** *Palais-Smale condition* Assume that  $G$  satisfies (G1) – (G3). Then  $I(z)$  satisfies the Palais-Smale condition: If for a sequence  $(z_k)$ ,  $I(z_k)$  is bounded from above and  $DI(z_k) \rightarrow 0$  as  $k \rightarrow \infty$ , then  $(z_k)$  has a convergent subsequence.

*Proof.* Let  $(z_k)$  be a sequence with

$$(3.1) \quad I(z_k) \leq M$$

and

$$(3.2) \quad DI(z_k) = z_k - A_\epsilon^{-1}(G_z(t, z_k) + \epsilon z_k) \rightarrow \theta \quad \text{as } m \rightarrow \infty,$$

where  $A_\epsilon^{-1}$  is compact operator and  $\theta = (0, \dots, 0)$ . We claim that  $\{z_k\}$  has a convergent subsequence. Since  $G_z(t, z(t)) + \epsilon z_k$  is bounded for a small constant  $\epsilon$  and  $A_\epsilon^{-1}$  is compact operator, by (3.2),  $\{z_k\}$  has a convergent subsequence. Thus we prove the lemma.  $\square$

Let

$$B_\rho = \{z \in E \mid \|z\| \leq \rho\},$$

$$\partial B_\rho = \{z \in E \mid \|z\| = \rho\},$$

$$Q = (\bar{B}_R \cap E_-) \oplus \{re \mid e \in \partial B_1 \cap E_+, 0 < r < R\}.$$

LEMMA 3.2. Assume that  $G$  satisfies the conditions (G1) – (G3). there exist a constant  $\rho > 0$  and sets  $S \subset E$ ,  $Q \subset E$  such that

- (i)  $\partial B_\rho \subset E_+$  and  $I|_{\partial B_\rho} > 0$ ,
- (ii)  $Q$  is bounded and  $I|_{\partial Q} < 0$ ,
- (iii)  $\partial B_\rho$  and  $\partial Q$  link.

*Proof.* (i) Let  $z \in E_+ \subset E$ . Since  $G(x, t, z)$  is bounded, there exists a constant  $C > 0$  such that

$$\begin{aligned} I(z) &= \frac{1}{2}(\|A_\epsilon^{\frac{1}{2}} z_+\|_{L^2}^2 + \|A_\epsilon^{\frac{1}{2}} M_+ z_0\|^2 - \|(-A_\epsilon)^{\frac{1}{2}} z_-\|_{L^2}^2 - \|(-A_\epsilon)^{\frac{1}{2}} M_- z_0\|^2) \\ &\quad - \int_0^{2\pi} G(t, z) dt - \frac{\epsilon}{2} \|z\|_{L^2}^2 \\ &\geq \frac{1}{2}(\|A_\epsilon^{\frac{1}{2}} z_+\|_{L^2}^2 - C - \frac{\epsilon}{2} \|z\|_{L^2}^2) \end{aligned}$$

for  $C > 0$ . Then there exist a constant  $\rho > 0$  such that if  $z \in \partial B_\rho \cap E_+$ , then  $I(z) > 0$ .

(ii) Let us choose  $e \in B_1 \cap E_+$ . Let  $z \in \bar{B}_r \cap E_- \oplus \{re \mid 0 < r\}$ . Then  $z = w + y$ ,  $w \in \bar{B}_r \cap E_-$ ,  $y = re$ . We note that

$$\text{If } w \in \bar{B}_r \cap E_-, \text{ then } \int_0^{2\pi} A_\epsilon(z) \cdot z dt = -\|(-A_\epsilon)^{\frac{1}{2}} z_-\|_{L^2}^2 \leq 0.$$



By (G3),  $G(t, w + re)$  is bounded from below. Thus, there exists a constant  $C_1 > 0$  such that if  $z = w + re$ , then we have

$$\begin{aligned} I(z) &= \frac{1}{2}r^2 - \|(-A_\epsilon)^{\frac{1}{2}}z_-\|_{L^2}^2 - \int_0^{2\pi} G(t, w + re)dt - \frac{\epsilon}{2}\|z\|_{L^2}^2 \\ &\leq \frac{1}{2}r^2 - \|(-A_\epsilon)^{\frac{1}{2}}z_-\|_{L^2}^2 + C_1 - \frac{\epsilon}{2}\|z\|_{L^2}^2. \end{aligned}$$

We can choose a constant  $R > r$  such that if  $z = w + re \in Q = (\bar{B}_r \cap E_-) \oplus \{re \mid e \in B_1 \cap E_+, 0 < r < R\}$ , then  $I(z) < 0$ . Thus we prove the lemma.  $\square$

#### PROOF OF THEOREM 1.1

By Lemma 2.1,  $I(z)$  is continuous and *Fréchet* differentiable in  $E$  and moreover  $DI \in C$ . By Lemma 3.1,  $I(z)$  satisfies the *(P.S.)* condition. By Lemma 3.2, there exist a constant  $\rho > 0$  and sets  $\partial B_\rho \subset E_+$  with radius  $\rho > 0$ ,  $Q \subset E$  such that  $I|_{\partial B_\rho} > 0$ ,  $Q$  is bounded and  $I|_{\partial Q} < 0$ , and  $\partial B_\rho$  and  $\partial Q$  link. By Theorem 2.1,  $I(z)$  possesses a critical value  $c > 0$ . Thus (1.1) has at least one nontrivial periodic weak solution. Thus we prove Theorem 1.1

#### References

- [1] K. C. Chang, *Infinite dimensional Morse theory and multiple solution problems*, Birkhäuser, 1993.
- [2] Benci, V., Rabinowitz, P.H, *Critical point theorems for indefinite functionals*, Invent. Math. **52** (1979), 241–273.
- [3] T. Jung and Q. H. Choi, *On the number of the periodic solutions of the nonlinear Hamiltonian system*, Nonlinear Anal. **71** (12) (2009), e1100-e1108.
- [4] T. Jung and Q. H. Choi, *Periodic solutions for the nonlinear Hamiltonian systems*, Korean J. Math. **17** (3) (2009), 331–340.
- [5] P. H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, CBMS. Regional Conf. Ser. Math. **65**, Amer. Math. Soc. Providence, Rhode Island, 1986.

Department of Mathematics Education  
 Inha University  
 Incheon 402-751, Korea  
*E-mail*: qheung@inha.ac.kr

Department of Mathematics  
Kunsan National University  
Kunsan 573-701, Korea  
*E-mail:* [tsjung@kunsan.ac.kr](mailto:tsjung@kunsan.ac.kr)