



Review Article

Sex-specific Profiles of Blood Metal Levels Associated with Metal–Iron Interactions

Byung-Kook Lee¹, Yangho Kim^{2,*}¹ Korean Industrial Health Association, Seoul, Korea² Department of Occupational and Environmental Medicine, Ulsan University Hospital, University of Ulsan College of Medicine, Ulsan, Korea

ARTICLE INFO

Article history:

Received 4 February 2014

Received in revised form

26 April 2014

Accepted 30 June 2014

Available online 5 July 2014

Keywords:

cadmium

ferritin

iron

lead

manganese

ABSTRACT

The mechanisms by which iron is absorbed are similar to those of divalent metals, particularly manganese, lead, and cadmium. These metals, however, show different toxicokinetics in relation to menarche or menopause, although their interaction with iron is the same. This review focuses on the kinetics of these three toxic metals (manganese, lead, and cadmium) in relation to menarche, pregnancy, and menopause. The iron–manganese interaction is the major factor determining sex-specific differences in blood manganese levels throughout the whole life cycle. The effects of estrogen overshadow the association between iron deficiency and increased blood lead concentrations, explaining why women, despite having lower ferritin concentrations, have lower blood lead concentrations than men. Iron deficiency is associated with elevated cadmium levels in premenopausal women, but not in postmenopausal women or men; these findings indicate that sex-specific differences in cadmium levels at older ages are not due to iron–cadmium interactions, and that further studies are required to identify the source of these differences. In summary, the potential causes of sex-specific differences in the blood levels of manganese, lead, and cadmium differ from each other, although all these three metals are associated with iron deficiency. Therefore, other factors such as estrogen effects, or absorption rate as well as iron deficiency, should be considered when addressing environmental exposure to toxic metals and sex-specific differences in the blood levels of these metals.

© 2014, Occupational Safety and Health Research Institute. Published by Elsevier. All rights reserved.

1. Introduction

Toxic metals have become ubiquitous in the built environment, and thus the general populations are increasingly becoming exposed to these metals. Most research in toxicology and environmental and occupational health has only involved men. However, increasing evidence suggests that the toxicity of environmental pollutants, including toxic metals, may manifest differently in women than in men. Although many epidemiological studies have reported data separately for men and women, differences between the genders have seldom been evaluated [1–3]. Moreover, little is known about toxicity in relation to specific periods in women's lives, such as menarche and menopause. Menstruation may cause iron deficiency, which, in turn, may be related to increased gastrointestinal absorption of toxic metals [4,5]. Mechanisms of iron absorption are similar to those of other divalent metals, particularly

manganese, lead, and cadmium [4,5] and a dietary deficiency of iron can lead to excess absorption of manganese [6–13], lead [14–23], and cadmium [24–29].

Metal ions show different toxicokinetics in relation to menarche, pregnancy, or menopause, despite similar metal–iron interactions. Thus, this review focuses on the kinetics of the toxic metals with a similar metal–iron interaction, such as manganese, lead, and cadmium, in relation to menarche, pregnancy, and menopause in the general population without occupational exposure.

2. Manganese

Manganese is a naturally occurring element that is abundantly present in the environment. It is an essential dietary nutrient needed for proper functioning of the human body at specific concentrations. It plays a role in bone formation, protein and energy

* Corresponding author. Department of Occupational and Environmental Medicine, Ulsan University Hospital, University of Ulsan College of Medicine, 290-3 Cheonha-Dong, Dong-Gu, Ulsan 682-060, Korea.

E-mail address: yanghokm@nuri.net (Y. Kim).

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

metabolism, and metabolic regulation, and functions as a cofactor in a number of enzymatic reactions [30]. Because manganese is an essential element [30–32], its absorption, disposition, and biliary excretion are actively controlled by homeostatic mechanisms [30,33]. These processes also play an important role in manganese toxicokinetics, which differ from those of nonessential toxic metals such as lead and cadmium. Heavy occupational exposure to manganese can cause a neurologic impairment clinically called “manganism,” a motor syndrome that is similar to but differentiated from idiopathic Parkinson’s disease [33–37]. The mechanisms involved in iron absorption are similar to those of divalent metals, particularly manganese [4,27], and a dietary deficiency of iron can lead to excess absorption of manganese [6–13]. Higher concentrations of blood manganese are reported in females of childbearing age than in males because the former have lower concentrations of ferritin [11,12,38]. Moreover, blood manganese levels are lower in menopausal than in premenopausal women because the former have higher concentrations of ferritin [39]. Significant increases in the mean whole blood manganese levels throughout the pregnancy period were reported previously [32,40,41]. This increase in manganese levels may be related to the enhanced absorption of manganese due to upregulation of iron absorption, particularly during the late pregnancy periods [42,43], because mechanisms of iron absorption are similar to those of other divalent metals, particularly manganese and cadmium [4,5]. However, age-related increase in blood manganese is not observed [11], because its absorption, disposition, and biliary excretion are actively controlled by homeostatic mechanisms as an essential element [30]. Thus, the iron–manganese interaction is the major factor determining sex-specific differences in blood manganese levels in relation to menstrual, reproductive, and menopausal factors (Table 1) [11,32,38–41].

3. Lead

Lead is a widespread environmental pollutant that can damage the central nervous, renal, cardiovascular, reproductive, and hematological systems. Lead accumulated in bones has a half-life ranging from years (trabecular bone) to decades (cortical bone) [44]. Approximately 90% of the lead in the body is stored in the skeleton [45].

Because iron is absorbed by mechanisms similar to those of other divalent metal ions, including lead and manganese, a dietary deficiency of iron can lead to excess absorption of lead [46–48]. Several studies have suggested an association between iron status and blood lead concentration in children [14–23] and in premenarchal females [49]. However, the relationship between iron deficiency and increased lead absorption has not been widely reported in adults [50,51]. Despite having lower ferritin levels, postmenarchal females have lower blood lead concentrations than males of similar ages [49,52–56]. Furthermore, blood lead concentrations are higher

in menopausal than in premenopausal women, despite the former having higher ferritin levels. Our previous study [49] suggested that these paradoxical findings are probably due to the potential confounding effect of estrogen on blood lead levels. Several studies have shown that blood lead concentrations are higher in menopausal than in menstruating women, because lead may be mobilized from the skeleton during periods of increased bone demineralization, such as in menopause [57–62]. In addition, lower blood lead levels were observed in postmenopausal women receiving estrogen replacement therapy than in past or never users [63–65]. The National Health and Nutrition Examination Survey data showed that blood lead levels were lower among girls who had attained sexual maturity [66]. Estrogen triggers rapid bone formation during pubertal development, inducing more rapid deposition of calcium, as well as lead, from blood into bone and possibly causing redistribution of lead throughout the body. These findings indicate that the effects of estrogen overshadow the association between iron deficiency and increased blood lead levels, and explain why women, despite having lower ferritin levels, have lower blood lead levels than men. These sex-specific differences in blood levels of lead between men and women were reported previously [53,67]. In addition, Wu et al [66] found a significant difference in blood lead concentration between Chinese male and female children aged 7–14 years (5.072 µg/dL vs. 4.389 µg/dL), but not in children aged <7 years. Our previous report also did not show a significant sex-specific difference in blood lead concentration between premenarchal girls and boys [49]. There was also a significant decrease in the mean whole blood lead levels throughout the pregnancy period [68–71] although some recovery is also seen in the late stages of pregnancy in some studies [70,72,73]. This decrease during pregnancy may be mainly related to physiological factors, such as increases in plasma estrogen concentrations and their effect on lead redistribution. Thus, the decreasing effect of estrogen on blood lead is a major determinant of sex-specific differences in blood lead levels as a common thread relating menstrual, reproductive, and menopausal factors (Table 2) [40,49,53,57,58,61,62,67–71,74,75]. However, age-related increase in blood lead is observed because of increase in exposure with age [67,74].

4. Cadmium

Cadmium is a ubiquitous environmental pollutant with a biological half-life in the body exceeding 10 years. Cadmium levels in the body accumulate with age, as only a minute part (0.01–0.02%) is excreted per day [76]. Cadmium has been reported to have cumulative effects on mortality and cardiovascular, neurologic, renal, and developmental diseases [76].

Blood cadmium is a valid biomarker of recent cadmium exposure [77], whereas urinary cadmium is a biomarker of lifetime exposure to cadmium [77]. Iron and cadmium have similar

Table 1
Behavior of blood manganese (Mn) concentrations according to age- and sex-related variables

| Variables | Refs (n) | Study populations and findings |
|-----------|---------------------------------------|---|
| Age | [11] (2,005) | Korean general population aged ≥20 y; KNHANES 2008/No significant change between population in the 20s and 40s |
| Sex | [11] (2,005) [38] (297) | Korean general population aged ≥20 y; KNHANES 2008/GM of blood Mn in females vs. males: 1.403 µg/dL vs. 1.192 µg/dL* Canadian general population/GM of blood Mn in females vs. males: 0.750 µg/dL vs. 0.675 µg/dL* |
| Menopause | [39] (1,826) | Korean general population KNHANES 2008–2009/GM of blood Mn in premenopausal vs. postmenopausal women: 1.443 µg/dL vs. 1.296 µg/dL* |
| Pregnancy | [32] (34) [40] (290) [41] (470) | Australian general population/maternal blood Mn during pregnancy from 10 wks to 20 wks vs. 34 wks: 0.375 µg/dL vs. 0.575 µg/dL Canadian general population/maternal blood GM of Mn during pregnancy at delivery vs. 1 st trimester and nonpregnant women: 1.56 µg/dL vs. 0.85 µg/dL and 0.746 µg/dL Canadian general population/maternal blood AM of Mn during pregnancy at delivery vs. nonpregnant 2.4 µg/dL vs. 0.8–1.2 µg/dL |

AM, arithmetic mean; GM, geometric mean; KNHANES, Korea National Health and Nutrition Examination Survey.

* Statistically significant.

Table 2
Behavior of blood lead concentrations according to age- and sex-related variables

| Variables | Refs (n) | Study populations and findings |
|-----------|--|--|
| Age | [67] (9,961) [74] (5,924) | U.S. general population; NHANES 1999–2002/significant increase with age Korean general population aged ≥ 20 y; KNHANES 2008–2010/significant increase with age |
| Sex | [67] (9,961) [53] (8,793) [74] (5,924) [75] (3,181) [49] (798) | U.S. general population; NHANES 1999–2002/GM of blood lead in men vs. women: 2.08 $\mu\text{g/dL}$ vs. 1.31 $\mu\text{g/dL}$ U.S. general population; NHANES 2009–2010/GM of blood lead in men vs. women: 1.31 $\mu\text{g/dL}$ vs. 0.966 $\mu\text{g/dL}$ Korean general population aged ≥ 20 y; KNHANES 2008–2010/GM of blood lead in men vs. women: 2.620 $\mu\text{g/dL}$ vs. 2.008 $\mu\text{g/dL}$ * Children aged 0–14 y from Beijing, China/AM of blood lead in male children vs. female children aged 7–14 y: 5.072 $\mu\text{g/dL}$ vs. 4.389 $\mu\text{g/dL}$, but no differences in children aged < 7 y Korean general adolescent population; KNHANES 2010–2011/GM of blood lead in premenarchal girls vs. boys: 1.403 $\mu\text{g/dL}$ * vs. 1.497 $\mu\text{g/dL}$ |
| Menarche | [40] (396) | Korean general adolescent population; KNHANES 2010–2011/GM of blood lead in premenarchal vs. postmenarchal girls: 1.403 $\mu\text{g/dL}$ vs. 1.136 $\mu\text{g/dL}$ * |
| Menopause | [61] (745) [57] (846) [58] (3,136) [62] (3,221) | Korean general population; KNHANES 2008–2009/blood GM of lead in premenopausal vs. postmenopausal women: 2.0 $\mu\text{g/dL}$ vs. 2.274 $\mu\text{g/dL}$ * U.S. general population; NHANES 1976–1980/blood GM of lead in premenopausal vs. postmenopausal women aged 40–60 y: 11.63 $\mu\text{g/dL}$ vs. 13.09 $\mu\text{g/dL}$ * U.S. general population; NHANES 1982–1984/blood GM of lead in premenopausal vs. postmenopausal women: 7.5 $\mu\text{g/dL}$ vs. 8.9 $\mu\text{g/dL}$ * U.S. general population; NHANES 1999–2010/blood GM of lead in premenopausal vs. postmenopausal women aged 45–55 y: 1.23 $\mu\text{g/dL}$ vs. 1.71 $\mu\text{g/dL}$ * |
| Pregnancy | [68] (26,570) [69] (128 vs. 120) [71] (165 vs. 27) [70] (12) | U.S. general population; NHANES 2003–2008/blood GM of lead among pregnant women aged 18–49 y vs. nonpregnant women: 0.64 $\mu\text{g/dL}$ vs. 0.85 $\mu\text{g/dL}$ * Chinese general population (Chengdu)/blood AM of 3 trimesters vs. healthy control group: 5.957 $\mu\text{g/dL}$, 5.517 $\mu\text{g/dL}$, and 5.577 $\mu\text{g/dL}$ vs. 6.87 $\mu\text{g/dL}$ * UK general population/blood AM of lead levels of pregnant vs. nonpregnant women: 12.22 $\mu\text{g/dL}$ vs. 13.25 $\mu\text{g/dL}$ (insignificant) Australian immigrants/changes in blood GM of lead during pregnancy and postpartum followed U-shaped patterns |

AM, arithmetic mean; GM, geometric mean; KNHANES, Korea National Health and Nutrition Examination Survey; NHANES, National Health and Nutrition Examination Survey.
* Statistically significant.

absorption mechanisms [4,27], and animal experiments have shown that metabolic interactions may take place between cadmium and iron [78,79]. Increased cadmium uptake in animals with low iron stores has been documented previously [24,25,28,29]. Moreover, cadmium concentrations were shown to be associated with decreasing iron stores in premenopausal women [1,12,52,80–85]. Thus, blood cadmium concentrations are higher in females of childbearing age than in males because the former have lower ferritin concentrations [53,74,86,87]. There was a significant increase in the whole blood cadmium levels during late pregnancy, in particular [80,88,89]. This increase in cadmium levels may be related to enhanced cadmium absorption due to an upregulated iron absorption, particularly during the late pregnancy period [43,80], because mechanisms of iron absorption are similar to those of other divalent metals, particularly manganese and cadmium [4,5]. However, blood cadmium concentrations have rarely been reported to decrease after menopause. Blood cadmium levels were

found to be higher in menopausal than in premenopausal women, and higher in postmenopausal women than in men of corresponding age. By contrast, one study found no statistically significant differences in blood cadmium levels in postmenopausal and premenopausal women [90]; however, urinary cadmium concentrations were higher in women than in men aged over 50 years [87], with between-gender differences greater in individuals aged > 50 years than those aged < 50 years [91]. In addition, cadmium levels were significantly higher in the hair of elderly females than males [92]. Similarly, our recent study [93] found that blood cadmium concentrations were higher in women than in men aged > 50 years, and there was no association between iron deficiency and elevated cadmium levels in postmenopausal women [94,95] or in men [85,96]. Thus, sex-specific differences in cadmium levels at older ages are not only due to iron–cadmium interaction, but also due to a higher absorption rate in females than in males that could be also influenced by other confounding factors (iron stores; Table 3)

Table 3
Behavior of blood/urine cadmium concentrations according to age- and sex-related variables

| Variables | Refs (n) | Study populations and findings |
|-----------|--|---|
| Age | [74] (5,924) [53] (8,793) [86] (2,257) [87] (1,055) | Korean general population aged ≥ 20 y; KNHANES 2008–2010/GM of blood cadmium had a significant increase with age U.S. general population NHANES 2009–2010/GM of urinary cadmium had a significant increase with age U.S. general population aged ≥ 6 y; NHANES 2003–2004/GM of urinary cadmium had a significant increase with age Bangladesh general population aged ≥ 8 y/median urinary cadmium levels had a significant increase with age |
| Sex | [53] (8,793) [74] (5,924) [86] (2,257) [87] (1,055) | U.S. general population; NHANES 2009–2010/GM of blood cadmium in men vs. women: 0.279 $\mu\text{g/L}$ vs. 0.326 $\mu\text{g/L}$ * Korean general population aged ≥ 20 y; KNHANES 2008–2010/GM of blood cadmium in men vs. women: 0.780 $\mu\text{g/L}$ vs. 1.194 $\mu\text{g/L}$ * U.S. general population aged ≥ 6 y; NHANES 2003–2004/GM of urinary cadmium in men aged ≥ 12 y was lower than in women; but no difference in children aged 6–11 y Bangladesh general population aged ≥ 8 y/median urinary cadmium level in men aged 30–50 y, 51–88 y vs. women; 0.66 $\mu\text{g/L}$ vs. 0.81 $\mu\text{g/L}$, 0.88 $\mu\text{g/L}$ vs. 1.1 $\mu\text{g/L}$ |
| Menopause | [93] (3,700) [87] (149) [91] (1,670) | Korean general population/blood GM of cadmium in premenopausal vs. postmenopausal women: 0.995 $\mu\text{g/L}$ vs. 1.165 $\mu\text{g/L}$ * Bangladesh general population aged > 51 y/median urinary cadmium in women vs. men: 1.1 $\mu\text{g/L}$ vs. 0.88 $\mu\text{g/L}$ German general population aged ≥ 25 y/between-gender differences in blood GM of cadmium greater in individuals aged > 50 y than aged < 50 y |
| Pregnancy | [97] (120) [88] (2,882) [89] (281) [80] (216) | Spanish general population/no significant changes in urinary GM of cadmium during pregnancy and postpartum: 0.44 $\mu\text{g/L}$ vs. 0.64 $\mu\text{g/L}$ Chinese general population/significant changes in the blood median cadmium between during late pregnancy periods and nonpregnant women; 0.75 $\mu\text{g/L}$ vs. 0.5 $\mu\text{g/L}$ Bangladesh general population/median blood cadmium increased 15% from early pregnancy (0.5 $\mu\text{g/L}$) to 6 mo postpartum Swedish general population/median blood cadmium increased 13% from early pregnancy (0.16 $\mu\text{g/L}$) to 3 mo postpartum |

GM, geometric mean; KNHANES, Korea National Health and Nutrition Examination Survey; NHANES, National Health and Nutrition Examination Survey.
* Statistically significant.

[53,74,80,86–89,91,93,97,98]. Further study is required to clarify the source of these differences. However, age-related increase in blood cadmium is remarkable because of increase in oral exposure with age [53,74,86].

5. Summary

Exposure is the major determinant of blood/urine metal concentrations in humans. This review focused on the general population without occupational exposure. We assumed a similar exposure between the two sexes, which is a limitation of this review, and discussed the kinetics of the toxic metals with a similar metal–iron interaction, such as manganese, lead, and cadmium, in relation to menstrual, reproductive, and menopausal factors.

The potential causes of sex-specific differences in the blood levels of manganese, lead, and cadmium differ from each other, although all three are associated with iron deficiency. Therefore, other factors such as estrogen effects, or absorption rate as well as iron deficiency, should be considered when addressing environmental exposure to toxic metals and sex-specific differences in the blood levels of these metals.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Acknowledgments

This study was supported by a grant of the Korean Health Technology R&D Project, Ministry of Health and Welfare, Republic of Korea (Grant No. HI13C0713).

References

- Vahter M, Berglund M, Akesson A, Lidén C. Metals and women's health. *Environ Res* 2002;88:145–55.
- Vahter M, Berglund M, Akesson A. Toxic metals and the menopause. *J Br Menopause Soc* 2004;10:60–4.
- Vahter M, Akesson A, Lidén C, Ceccatelli S, Berglund M. Gender differences in the disposition and toxicity of metals. *Environ Res* 2007;104:85–95.
- Hurley LS, Keen CL. Manganese. In: Underwood E, Mertz W, editors. Trace elements in human health and animal nutrition. New York: Academic Press; 1987. p. 185–223.
- Mackenzie B, Garrick MD. Iron imports. II. Iron uptake at the apical membrane in the intestine. *Am J Physiol Gastrointest Liver Physiol* 2005;289:G981–6.
- Davis CD, Wolf TL, Greger JL. Varying levels of manganese and iron affect absorption and gut endogenous losses of manganese by rats. *J Nutr* 1992;122:1300–8.
- Mena I, Horiuchi K, Burke K, Cotzias GC. Chronic manganese poisoning. Individual susceptibility and absorption of iron. *Neurology* 1969;19:1000–6.
- Finley JW, Johnson PE, Johnson LK. Sex affects manganese absorption and retention by humans from a diet adequate in manganese. *Am J Clin Nutr* 1994;60:949–55.
- Finley JW. Manganese absorption and retention by young women is associated with serum ferritin concentration. *Am J Clin Nutr* 1999;70:37–43.
- Kim Y, Park JK, Choi Y, Yoo CI, Lee CR, Lee H, Lee JH, Kim SR, Jeong TH, Yoon CS, Park JH. Blood manganese concentration is elevated in iron deficiency anemia patients, whereas globus pallidus signal intensity is minimally affected. *Neurotoxicology* 2005;26:107–11.
- Kim Y, Lee BK. Iron deficiency increases blood manganese level in the Korean general population according to KNHANES 2008. *Neurotoxicology* 2011;32:247–54.
- Meltzer HM, Brantsaeter AL, Borch-Johnsen B, Ellingsen DG, Alexander J, Thomassen Y, Stigum H, Ydersbond TA. Low iron stores are related to higher blood concentrations of manganese, cobalt and cadmium in non-smoking, Norwegian women in the HUNT 2 study. *Environ Res* 2010;110:497–504.
- Park S, Sim CS, Lee H, Kim Y. Blood manganese concentration is elevated in infants with iron deficiency. *Biol Trace Elem Res* 2013;155:184–9.
- Bradman A, Eskenazi B, Sutton P, Athanasoulis M, Goldman LR. Iron deficiency associated with higher blood lead in children living in contaminated environments. *Environ Health Perspect* 2001;109:1079–84.
- Choi JW, Kim SK. Association between blood lead concentrations and body iron status in children. *Arch Dis Child* 2003;88:791–2.
- Hammad TA, Sexton M, Langenberg P. Relationship between blood lead and dietary iron intake in preschool children. A cross-sectional study. *Ann Epidemiol* 1996;6:30–3.
- Park S, Sim CS, Lee H, Kim Y. Effects of iron therapy on blood lead concentrations in infants. *J Trace Elem Med Biol* 2014;28:56–9.
- Turgut S, Polat A, Inan M, Turgut G, Emmungil G, Bican M, Karakus TY, Genc O. Interaction between anemia and blood levels of iron, zinc, copper, cadmium and lead in children. *Indian J Pediatr* 2007;74:827–30.
- Watson WS, Hume R, Moore MR. Oral absorption of lead and iron. *Lancet* 1980;2:236–7.
- Watson WS, Morrison J, Bethel MI, Baldwin NM, Lyon DT, Dobson H, Moore MR, Hume R. Food iron and lead absorption in humans. *Am J Clin Nutr* 1986;44:248–56.
- Wright RO, Shannon MW, Wright RJ, Hu H. Association between iron deficiency and low-level lead poisoning in an urban primary care clinic. *Am J Public Health* 1999;189:1049–53.
- Yip R, Norris TN, Anderson AS. Iron status of children with elevated blood lead concentrations. *J Pediatr* 1981;98:922–5.
- Yip R, Dallman PR. Developmental changes in erythrocyte protoporphyrin: roles of iron deficiency and lead toxicity. *J Pediatr* 1984;104:710–3.
- Park JD, Cherrington NJ, Klaassen CD. Intestinal absorption of cadmium is associated with divalent metal transporter 1 in rats. *Toxicol Sci* 2002;68:288–94.
- Ryu DY, Lee SJ, Park DW, Choi BS, Klaassen CD, Park JD. Dietary iron regulates intestinal cadmium absorption through iron transporters in rats. *Toxicol Lett* 2004;152:19–25.
- Tallkvist J, Bowlus CL, Lönnnerdal B. DMT1 gene expression and cadmium absorption in human absorptive enterocytes. *Toxicol Lett* 2001;122:171–7.
- Zoller H, Koch RO, Theurl I, Obrist P, Pietrangolo A, Montosi G, Haile DJ, Vogel W, Weiss G. Expression of the duodenal iron transporters divalent-metal transporter 1 and ferroportin 1 in iron deficiency and iron overload. *Gastroenterology* 2001;120:1412–9.
- Flanagan PR, McLellan JS, Haist J, Cherian G, Chamberlain MJ, Valberg LS. Increased dietary cadmium absorption in mice and human subjects with iron deficiency. *Gastroenterology* 1978;74:841–6.
- Piasek M, Blanus M, Kostial K, Laskey JW. Low iron diet and parenteral cadmium exposure in pregnant rats: the effects on trace elements and fetal viability. *Biometals* 2004;17:1–14.
- Karki P, Lee E, Aschner M. Manganese neurotoxicity: a focus on glutamate transporters. *Ann Occup Environ Med* 2013;25:4.
- Bellinger DC. Prenatal exposures to environmental chemicals and children's neurodevelopment: an update. *Saf Health Work* 2013;4:1–11.
- Spencer A. Whole blood manganese levels in pregnancy and the neonate. *Nutrition* 1999;15:731–4.
- Lucchini RG, Kim Y. Health effects of manganese. In: Vojtisek M, Prakash R, editors. Metals and neurotoxicity. Jalgaon, Maharashtra (India): Society for Science and Environment; 2009. p. 119–47.
- Barbeau A, Inoué N, Cloutier T. Role of manganese in dystonia. *Adv Neurol* 1976;14:339–52.
- Calne DB, Chu NS, Huang CC, Lu CS, Olanow W. Manganese and idiopathic parkinsonism: similarities and differences. *Neurology* 1994;44:1583–6.
- Olanow CW. Manganese-induced parkinsonism and Parkinson's disease. *Ann N Y Acad Sci* 2004;1012:209–23.
- Kim Y, Kim JW. Toxic encephalopathy. *Saf Health Work* 2012;3:243–56.
- Baldwin M, Mergler D, Larribe F, Bélanger S, Tardif R, Bilodeau L, Hudnell K. Bioindicator and exposure data for a population based study of manganese. *Neurotoxicology* 1999;20:343–53.
- Lee BK, Kim Y. Effects of menopause on blood manganese levels in women: analysis of 2008–2009 Korean National Health and Nutrition Examination Survey data. *Neurotoxicology* 2012;33:401–5.
- Takser L, Lafond J, Bouchard M, St-Amour G, Mergler D. Manganese levels during pregnancy and at birth: relation to environmental factors and smoking in a Southwest Quebec population. *Environ Res* 2004;95:119–25.
- Zota AR, Ettinger AS, Bouchard M, Amarasiwardena CJ, Schwartz J, Hu H, Wright RO. Maternal blood manganese levels and infant birth weight. *Epidemiology* 2009;20:367–73.
- Kirchgesner M, Sherif YS, Schwarz FJ. Changes in absorption of manganese during gravidity and lactation. *Ann Nutr Metab* 1982;26:83–9 [in German].
- Bothwell TH. Iron requirements in pregnancy and strategies to meet them. *Am J Clin Nutr* 2000;72:257S–64S.
- World Health Organization (WHO). Environmental health criteria 165. Inorganic lead. Geneva (Switzerland): WHO; 1995.
- Pounds JG, Long GJ, Rosen JF. Cellular and molecular toxicity of lead in bone. *Environ Health Perspect* 1991;91:17–32.
- Conrad ME, Barton JC. Factors affecting the absorption and excretion of lead in the rat. *Gastroenterology* 1978;74:731–40.
- Ragan HA. Effects of iron deficiency on the absorption and distribution of lead and cadmium in rats. *J Lab Clin Med* 1977;90:700–6.
- Six KM, Goyer RA. The influence of iron deficiency on tissue content and toxicity of ingested lead in the rat. *J Lab Clin Med* 1977;74:128–36.
- Sim CS, Kim Y, Lee H, Park CY, Ham JO, Lee BK. Iron deficiency increases blood lead levels in boys and pre-menarche girls surveyed in KNHANES 2010–2011. *Environ Res* 2014;130C:1–6.
- Flanagan PR, Chamberlain MJ, Valberg LS. The relationship between iron and lead absorption in humans. *Am J Clin Nutr* 1982;36:823–9.

- [51] Lee BK, Kim Y. Iron deficiency is associated with increased levels of blood cadmium in the Korean general population: analysis of 2008–2009 Korean National Health and Nutrition Examination Survey data. *Environ Res* 2012;112:155–63.
- [52] Bărăny E, Bergdahl IA, Bratteby LE, Lundh T, Samuelson G, Skerfving S, Oskarsson A. Iron status influences trace element levels in human blood and serum. *Environ Res* 2005;98:215–23.
- [53] Centers for Disease Control and Prevention (CDC). Fourth National Report on Human Exposure to Environmental (Updated Tables). Atlanta (GA): CDC; 2013. p. 164.
- [54] Kim NS, Lee BK. National estimates of blood lead, cadmium, and mercury levels in the Korean general adult population. *Int Arch Occup Environ Health* 2011;84:53–63.
- [55] Kristiansen J, Christensen JM, Iversen BS, Sabbioni E. Toxic trace element reference levels in blood and urine: influence of gender and lifestyle factors. *Sci Total Environ* 1997;204:147–60.
- [56] Watanabe T, Fujita H, Koizumi A, Chiba K, Miyasaka M, Ikeda M. Baseline level of blood lead concentration among Japanese farmers. *Arch Environ Health* 1985;40:170–6.
- [57] Silbergeld EK, Schwartz J, Mahaffey K. Lead and osteoporosis: mobilization of lead from bone in postmenopausal women. *Environ Res* 1988;47:79–94.
- [58] Symanski E, Hertz-Picciotto I. Blood lead levels in relation to menopause, smoking, and pregnancy history. *Am J Epidemiol* 1995;141:1047–58.
- [59] Muldoon SB, Salamone LM, Cauley JA, Allen L. Effect of bone mineral density changes on blood lead levels in peri-menopausal women. (Abstract). *Am J Epidemiol* 1997;145:S21.
- [60] Grandjean P, Nielsen GD, Jørgensen PJ, Hørdler M. Reference intervals for trace elements in blood: significance of risk factors. *Scand J Clin Lab Invest* 1992;52:321–37.
- [61] Lee BK, Kim Y. Association between bone mineral density and blood lead level in menopausal women: analysis of 2008–2009 Korean National Health and Nutrition Examination Survey data. *Environ Res* 2012;115:59–65.
- [62] Mendola P, Brett K, Dibari JN, Pollack AZ, Tandon R, Shenassa ED. Menopause and lead body burden among US women aged 45–55. *NHANES 1999–2010*. *Environ Res* 2013;121:110–3.
- [63] Berkowitz GS, Wolff MS, Lapinski RH, Todd AC. Prospective study of blood and tibia lead in women undergoing surgical menopause. *Environ Health Perspect* 2004;112:1673–8.
- [64] Jurczak A, Brodowski J, Grochans E, Karakiewicz B, Szkup-Jabłońska M, Wiederszała S, Mroczek B, Włoszczak-Szubska A, Grzywacz A. Effect of menopausal hormone therapy on the levels of magnesium, zinc, lead and cadmium in postmenopausal women. *Ann Agric Environ Med* 2013;20:147–51.
- [65] Nash D, Silbergeld E, Mager L, Stolley P. Menopause, hormone replacement therapy (HRT), and blood lead levels among adult women from NHANES III, 1988–1994. (Abstract). *Am J Epidemiol* 1998;147:S93.
- [66] Wu T, Buck GM, Mendola P. Blood lead levels and sexual maturation in U.S. girls: the Third National Health and Nutrition Examination Survey, 1988–1994. *Environ Health Perspect* 2003;111:737–41.
- [67] Muntner P, Menke A, DeSalvo KB, Rabito FA, Batuman V. Continued decline in blood lead levels among adults in the United States: the National Health and Nutrition Examination Surveys. *Arch Intern Med* 2005;165:2155–61.
- [68] Jones L, Parker JD, Mendola P. Blood lead and mercury levels in pregnant women in the United States, 2003–2008. *NCHS Data Brief* 2010:1–8.
- [69] Jiang Y, Wang H, Chen J, Zhang G, Chen L, Dai W, Zhou W, Yang H, Shi H. Blood lead levels during different trimesters of pregnancy and the possible influencing factors in Chengdu, China. *Biol Trace Elem Res* 2011;144:27–35.
- [70] Gulson BL, Mizon KJ, Palmer JM, Korsch MJ, Taylor AJ, Mahaffey KR. Blood lead changes during pregnancy and postpartum with calcium supplementation. *Environ Health Perspect* 2004;112:1499–507.
- [71] Alexander FW, Delves HT. Blood lead levels during pregnancy. *Int Arch Occup Environ Health* 1981;48:35–9.
- [72] Hertz-Picciotto I, Schramm M, Watt-Morse M, Chantala K, Anderson J, Osterloh J. Patterns and determinants of blood lead during pregnancy. *Am J Epidemiol* 2000;152:829–37.
- [73] Lagerkvist BJ, Ekesrydh S, Englyst V, Nordberg GF, Soderberg HA, Wiklund DE. Increased blood lead and decreased calcium levels during pregnancy: a prospective study of Swedish women living near a smelter. *Am J Public Health* 1996;86:1247–52.
- [74] Kim Y, Lee BK. Associations of blood lead, cadmium, and mercury with estimated glomerular filtration rate in the Korean general population: analysis of 2008–2010 Korean National Health and Nutrition Examination Survey data. *Environ Res* 2012;118:124–9.
- [75] Wu Y, Yang X, Ge J, Zhang J. Blood lead level and its relationship to certain essential elements in the children aged 0 to 14 years from Beijing, China. *Sci Total Environ* 2011;409:3016–20.
- [76] Agency for Toxic Substances and Disease Registry. Toxicological profile for cadmium. Atlanta (GA): Agency for Toxic Substances and Disease Registry; 2008.
- [77] Järup L, Åkesson A. Current status of cadmium as an environmental health problem. *Toxicol Appl Pharmacol* 2009;238:201–8.
- [78] Goyer RA. Nutrition and metal toxicity. *Am J Clin Nutr* 1995;61:646S–50S.
- [79] Goyer RA. Toxic and essential metal interactions. *Annu Rev Nutr* 1997;17:37–50.
- [80] Åkesson A, Berglund M, Schütz A, Bjellerup P, Bremme K, Vahter M. Cadmium exposure in pregnancy and lactation in relation to iron status. *Am J Public Health* 2002;92:284–7.
- [81] Berglund M, Åkesson A, Nermell B, Vahter M. Intestinal absorption of dietary cadmium in women depends on body iron stores and fiber intake. *Environ Health Perspect* 1994;102:1058–66.
- [82] Gallagher CM, Chen JJ, Kovach JS. The relationship between body iron stores and urine cadmium concentrations in US never-smoking, non-pregnant women aged 20–49 years. *Environ Res* 2011;111:702–7.
- [83] Kippler M, Ekström EC, Lönnerdal B, Goessler W, Åkesson A, El Arifeen S, Persson LA, Vahter M. Influence of iron and zinc status on cadmium accumulation in Bangladeshi women. *Toxicol Appl Pharmacol* 2007;222:221–6.
- [84] Mijal RS, Holzman CB. Blood cadmium levels in women of childbearing age vary by race/ethnicity. *Environ Res* 2010;110:505–12.
- [85] Satarug S, Ujjin P, Vanavanitkun Y, Baker JR, Moore MR. Influence of body iron store status and cigarette smoking on cadmium body burden of healthy Thai women and men. *Toxicol Lett* 2004;148:177–85.
- [86] Ruiz P, Mumtaz M, Osterloh J, Fisher J, Fowler BA. Interpreting NHANES biomonitoring data, cadmium. *Toxicol Lett* 2010;198:44–8.
- [87] Berglund M, Lindberg AL, Rahman M, Yunus M, Grandér M, Lönnerdal B, Vahter M. Gender and age differences in mixed metal exposure and urinary excretion. *Environ Res* 2011;111:1271–9.
- [88] Zhang Z, Yuan E, Liu J, Lou X, Jia L, Li X, Zhang L. Gestational age-specific reference intervals for blood copper, zinc, calcium, magnesium, iron, lead, and cadmium during normal pregnancy. *Clin Biochem* 2013;46:777–80.
- [89] Kippler M, Goessler W, Nermell B, Ekström EC, Lönnerdal B, El Arifeen S, Vahter M. Factors influencing intestinal cadmium uptake in pregnant Bangladeshi women—a prospective cohort study. *Environ Res* 2009;109:914–21.
- [90] González-Estecha M, Trasobares E, Fuentes M, Martínez MJ, Cano S, Vergara N, Gaspar MJ, González-Revaldería J, Barciela MC, Bugarín Z, Fernández MD, Badía P, Pintos C, González M, Guillén JJ, Bermejo P, Fernández C, Arroyo M., PESA Group. Blood lead and cadmium levels in a six hospital employee population. PESA study, 2009. *J Trace Elem Med Biol* 2011;25:S22–9.
- [91] Hoffmann K, Krause C, Seifert B. The German Environmental Survey 1990/92 (GerES II): primary predictors of blood cadmium levels in adults. *Arch Environ Health* 2001;56:374–9.
- [92] Yasuda H, Yoshida K, Yasuda Y, Tsutsui T. Two age-related accumulation profiles of toxic metals. *Curr Aging Sci* 2012;5:105–11.
- [93] Kim SH, Kim Y, Kim NS, Lee BK. Gender difference in blood cadmium concentration in the general population: can it be explained by iron deficiency? *J Trace Elem Med Biol* 2014;28:322–7.
- [94] Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, Kayama F. Comprehensive study of the effects of age, iron deficiency, diabetes mellitus, and cadmium burden on dietary cadmium absorption in cadmium-exposed female Japanese farmers. *Toxicol Appl Pharmacol* 2004;196:114–23.
- [95] Tsukahara T, Ezaki T, Moriguchi J, Furuki K, Fukui Y, Ukai H, Okamoto S, Sakurai H, Ikeda M. No significant effect of iron deficiency on cadmium body burden or kidney dysfunction among women in the general population in Japan. *Int Arch Occup Environ Health* 2003;76:275–81.
- [96] Olsson IM, Bensryd I, Lundh T, Ottosson H, Skerfving S, Oskarsson A. Cadmium in blood and urine—impact of sex, age, dietary intake, iron status, and former smoking—association of renal effects. *Environ Health Perspect* 2002;110:1185–90.
- [97] Hernandez M, Schuhmacher M, Fernandez JD, Domingo JL, Llobet JM. Urinary cadmium levels during pregnancy and postpartum. A longitudinal study. *Biol Trace Elem Res* 1996;53:205–12.
- [98] Choudhury H, Harvey T, Thayer WC, Lockwood TF, Stiteler WM, Goodrum PE, Hassett JM, Diamond GL. Urinary cadmium elimination as a biomarker of exposure for evaluating a cadmium dietary exposure—biokinetics model. *J Toxicol Environ Health A* 2001;63:321–50.