



Kidney size estimation in Korean children with Technesium-99m dimercaptosuccinic acid scintigraphy

Min Jung Lee, MD, Mi Kyung Son, MD, Byung Ok Kwak, MD, Hye Won Park, MD, Sochung Chung, MD, PhD, Kyo Sun Kim, MD, PhD
Department of Pediatrics, Konkuk University Medical Center, Konkuk University School of Medicine, Seoul, Korea

Purpose: Renal size is an important indicator to determine adequate organ growth in children. The aim of this study was to estimate renal size with Technesium-99m dimercaptosuccinic acid (DMSA) scan and propose a simple formula for predicting renal length in normal Korean children.

Methods: This study included 346 children (148 boys and 198 girls; age range, 1 month to 17 years) in whom renal length was measured using the DMSA scan. Patients with anatomical renal abnormalities or acute pyelonephritis were excluded. Children were divided into two groups: 214 children (61.8%) were less than a year old (group 1) and 132 (38.2%) were ≥ 1 year (group 2).

Results: Renal length was larger on the left side than the right side, and there was no significant gender-related difference in renal length. We propose the following formula for renal length based on the analysis of the 346 children in our study: the formula was as follows: $4.682 \times \text{age (month)}^{0.137}$, $R^2=0.780$. In group 1, the formula was renal length (cm) = $0.127 \times \text{age (month)} + 5.144$, $R^2=0.354$, and in group 2, the formula was $0.334 \times \text{age (year)} + 6.477$, $R^2=0.829$.

Conclusion: It is difficult to establish simple formulae in infants ($R^2=0.354$). Therefore, further studies including relevant variables are needed for this age group. We proposed formulae to estimate renal length in Korean children over 1 year of age by using the DMSA scan.

Key words: Kidney, Length, Child, Technesium-99m dimercaptosuccinic acid scan

Corresponding author: Kyo Sun Kim, MD, PhD
Department of Pediatrics, Konkuk University Medical Center, 120 Neungdong-ro, Gwangjin-gu, Seoul 143-729, Korea

Tel: +82-2-2030-7370

Fax: +82-2-2030-5009

E-mail: 19890009@kuh.ac.kr

Received: 17 August, 2013

Revised: 13 September, 2013

Accepted: 10 October, 2013

Introduction

Accurate measurement of organ size is important in estimating adequate growth in children. Renal size is an important indicator not only to evaluate the children with renal disease, but also to determine adequate organ growth¹⁻⁴. Renal size can be estimated by measuring renal length, renal volume, cortical volume or thickness. Despite the measurement of renal volume is the most accurate method to estimate renal size, the measurement of renal length is widely accepted because of its simplicity and reproducibility^{5,6}.

There are several published reports about the renal length measurement by renal ultrasonography (US), but little is known about the renal length measurement using Technesium-99m dimercaptosuccinic acid (DMSA) scan⁷. Although renal US has been widely used to measure renal length in children, inter- and intra-observer variability exist^{6,8,9}. DMSA scan is known more reproducible than renal US, and can also detect cortical abnormalities and assess renal function¹⁰.

In this study, we measured renal length with DMSA scan and established a simple equation for predicting renal length in Korean children.

Copyright © 2014 by The Korean Pediatric Society

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Materials and methods

1. Patients and methods

This study included 700 hospitalized children who were clinically suspected with urinary tract infection and underwent DMSA scan at Konkuk University Medical Center from May 2005 to August 2011. DMSA scan were performed to the patients who had pyuria in urinalysis collected by urine bag, urine catheterization, midstream urine sample or took antibiotics prior to admission, after the fever subsided. The children with clinical pyelonephritis and presented renal scarring or culture-proven urinary tract infection were excluded. The patients were excluded if they had renal abnormalities on US or DMSA scan such as altered echogenicity, solitary kidney, multicystic kidney disease, hydronephrosis, vesicourethral reflux, acute nephritis, photon defect of kidney or renal scar. Patients with organ transplantation, renal surgery, failure to thrive were also excluded. Medical records of 346 children who met inclusion criteria were reviewed retrospectively.

There were 148 boys and 198 girls. All those examined were divided into two age groups: less than 1 year old and 1 year of age or older. Because kidney growth were highest and most variable during the first year of life, but leveled off afterward.

Renal lengths were measured in both kidneys by assessing the maximum pixel length on DMSA scan. And the mean renal lengths was defined as half of the sum of the standard deviation scores in the right and left kidney. The mean renal length was used for further analysis, being referred to as 'Renal length' unless stated otherwise.

2. Statistical analysis

Data were analyzed by PASW ver. 18.0 (SPSS Inc., Chicago, IL, USA). Differences in variables between the groups were compared using independent t-test, and the correlations between age and renal length were obtained using regression analysis and Pearson correlation analysis. The relationship between whole age group and renal length was assessed by power regression model. Using R^2 as the criterion for comparing and choosing linear and nonlinear fashion, renal length less than 1 year old were statistically analyzed using a nonlinear model described; The group of 1 year of age or older was analyzed using a linear regression model of the general form $y=ax+b$. Values with $P<0.05$ were considered statistically significant.

Results

A total of 346 Korean children, 148 boys (42.8%) and 198 girls (57.2%), were included in this study. Their median age was 1.35

years with age distribution between 1 month and 17 years.

Children were divided into two age groups: 214 children (61.8%) were less than 1 year old (group 1) and 132 (38.2%) were 1 year of age or older (group 2). Mean renal length was 5.86 ± 0.62 cm in group 1, and 7.65 ± 1.36 cm in group 2 ($P<0.001$) (Table 1). Renal length was larger in left side than right side (6.63 ± 1.36 cm vs. 6.46 ± 1.28 cm, $P<0.001$), and no statistically significant difference in size between the gender in group 2. Renal length increased with age (Fig. 1). Growth rates of renal length were highest during the first year of life, and then stabilized in a linear fashion after 1 year of age. From the data of the 346 children, the regression equation for renal length was produced as follows:

Table 1. Clinical characteristics and renal size of the subjects

Characteristic	Group 1 (n=214)	Group 2 (n=132)
Age	5.70 ± 2.95 (mo)	3.54 ± 3.72 (yr)
Sex (male/female)	107/107	41/91
Renal length (cm)		
Total	$5.86\pm 0.62^*$	$7.65\pm 1.36^*$
Left side	$5.92\pm 0.68^\dagger$	$7.79\pm 1.41^\dagger$
Right side	$5.80\pm 0.63^\dagger$	$7.53\pm 1.35^\dagger$
Boys	$5.72\pm 0.59^\ddagger$	7.69 ± 1.27
Girls	$6.01\pm 0.63^\ddagger$	7.65 ± 1.41

Values are presented as mean±standard deviation.

*Statistical differences between group 1 and 2 ($P<0.001$). † Statistical differences between left and right sides ($P<0.001$). ‡ Statistical differences between genders ($P<0.001$).




Fig. 1. Scattered diagram of mean renal length measured using the Technesium-99m dimercaptosuccinic acid scan in all the study subjects. Renal length (cm) = $4.682 \times \text{age (mo)}^{0.137}$, $R^2 = 0.780$.




Fig. 2. Scatterplot and linear regression results of mean renal length obtained using Technesium-99m dimercaptosuccinic acid scan in children less than 1 year (A) and ≥1 year (B). (A) Renal length in children less than 1 year of age. Renal length (cm)=0.127×age (mo)+5.144, $R^2=0.354$. (B) Renal length in children ≥1 year. Renal length (cm)=0.334 age (yr)+6.477, $R^2=0.829$.

Table 2. Coefficients of determination (R^2) of formulas for determining renal length

Variable	Age<1 yr	Age≥1 yr
	Length (cm)= 0.127×age (mo)+5.144	Length (cm)= 0.334×age (yr)+6.477
Total	0.354	0.829
Left side	0.346	0.787
Right side	0.301	0.829
Boys	0.376	0.805
Girls	0.275	0.838

4.682×age (month)^{0.137}, $R^2=0.780$. In group 1, the formula was renal length (cm)=0.127×age (month)+5.144, $R^2=0.354$ (Fig. 2A), and 0.334×age (year)+6.477, $R^2=0.829$ in group 2 (Fig. 2B). Coefficients of determination (R^2) were calculated overall group and separately for left kidneys alone, right kidneys alone, boys alone and girls alone. R^2 values of subsets were presented in Table 2.

Discussion

In this study, we measured renal length with age using DMSA scan and produced a simple formula for predicting renal length in children ($R^2=0.780$). The formula for the renal length were presented separately according to the age (1 year) based on the growth velocity of the kidney.

Accurate measurement of organ size is important in children to estimate the adequate growth. Renal size measurement is one of the important indicator to determine adequate growth and renal function^{1,2,11}. Renal length can be measured by several modalities such as renal US, computed tomography (CT) scan, and DMSA scan^{5,12}. CT scan has several disadvantages for children such as radiation exposure and potentially nephrotoxic contrast agent administration, therefore it is recommended as secondary diagnostic tools for further evaluation of renal lesions that are detected with other imaging modalities^{5,13}. Renal US is one of the most common imaging method for assessment of visceral organ without any risk of radiation. There are many previous studies about measurement of renal length by US^{5,14-17}. Although renal US has been widely used in children, inter- and intra-observer variability exists in measuring kidney size^{6,8,9,18,19}. DMSA scan is known more reproducible and accurate modality than US in measuring renal length, as the maximum longitudinal renal axis can be more reliably obtained²⁰. It can also detect cortical abnormalities and assess renal function²¹, and has less radiation exposure compared to CT scan^{7,10,20,22}. Although DMSA scan is a useful tool for estimating renal size in children, a few studies has been published^{7,10,20,22}. Several published studies have been suggested simple equations to estimate the renal size by US in normal Korean children^{23,24}. But there is no reports with DMSA scan.

This study showed that left kidneys are larger than right kidney which was consistent with prior studies^{23,25}.

Akhavan et al.²⁵ proposed age-based formula for estimating renal length by US in children over 1 year old; renal length (cm)=0.278×age (year)+6.103 ($R^2=0.812$). It is similar to our age-based formula that examined by DMSA scan. Lobo Sotomayor et al.⁷ proposed similar formula using DMSA scan; renal length (cm)=0.2644×age (year)+6.319 for children over 1 year old and 6.1727×age (year)^{0.1535} in whole age group. They are similar to

our age-based formula in Korean children examined by DMSA scan. Further studies are required to establish age-based formula for estimating renal length and reference values using various modalities.

Our study showed that renal length increased with age, and the relationship between renal length and age was biphasic which consisted with previous report²⁵. It is well known that most rapid renal growth occurs during the first year of life and then the renal growth rate finally stabilizes in linear fashion^{16,25,26}. So the children were divided into two age groups: children less than 1 year old and children 1 year or older.

In this study, the renal length could be predicted with this linear equation in children older than 1 year of age ($R^2=0.829$). Meanwhile, the ability of explanation of formula for renal length in infants was much lower ($R^2=0.354$). Renal length is also influenced by intrauterine factors and gestational size during the first year of life^{16,25,27-29}. Therefore, it is difficult to establish simple formula in infants. Further studies including large population and associated factors are needed in this age group.

This study had some limitations. The data was obtained retrospectively from the patients who were clinically suspected with urinary tract infection and performed DMSA scan. However, for minimizing the factors that affect this study, we excluded the patients with a fever, a positive urine culture result, acute pyelonephritis and with genitourinary anomalies. A prospective study will be necessary in normal Korean children.

In addition, we did not consider relevant variables such as height, weight, nutritional status and genetic factors^{3,16,30}. Following studies would be required considering these relevant variables.

In conclusion, we established the simple age-based equation predicting renal length by DMSA scan in children over 1 year of age. It is suggested that this formula would be a useful tool to estimate adequate renal size and growth in Korean children in clinical practice.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

References

1. Lane PH, Belsha CW, Plummer J, Steinhardt GF, Lynch RE, Wood EG. Relationship of renal size, body size, and blood pressure in children. *Pediatr Nephrol* 1998;12:35-9.
2. Koff SA, Peller PA. Diagnostic criteria for assessing obstruction in the newborn with unilateral hydronephrosis using the renal

- growth-renal function chart. *J Urol* 1995;154(2 Pt 2):662-6.
3. Chen JJ, Pugach J, Patel M, Luisiri A, Steinhardt GF. The renal length nomogram: multivariable approach. *J Urol* 2002;168:2149-52.
4. Gavela T, Sanchez Bayle M, Gomez Mardones G, Gallego S, Martinez-Perez J, Moya MT. Ecographic study of kidney size in children. *Nefrologia* 2006;26:325-9.
5. Kadioglu A. Renal measurements, including length, parenchymal thickness, and medullary pyramid thickness, in healthy children: what are the normative ultrasound values? *AJR Am J Roentgenol* 2010;194:509-15.
6. Zerlin JM, Blane CE. Sonographic assessment of renal length in children: a reappraisal. *Pediatr Radiol* 1994;24:101-6.
7. Lobo Sotomayor G, Ladron de Guevara Hernandez D, Arnello Viveros F, Perez Rivera A, Jimenez Jorquera C. Renal size in children with normal renal 99mTc-DMSA scan: building a nomogram with scintigraphic length. *Rev Esp Med Nucl* 2004;23:102-5.
8. Schlesinger AE, Hernandez RJ, Zerlin JM, Marks TI, Kelsch RC. Interobserver and intraobserver variations in sonographic renal length measurements in children. *AJR Am J Roentgenol* 1991;156:1029-32.
9. Sargent MA, Wilson BP. Observer variability in the sonographic measurement of renal length in childhood. *Clin Radiol* 1992;46:344-7.
10. Kim SS, Bang WJ, Seo JW, Cho KS, Han SW. Discrepancy of measured renal length between ultrasonography and dimercaptosuccinic acid (DMSA) scintigraphy. *Korean J Urol* 2007;48:77-81.
11. Kim BW, Song MK, Chung S, Kim KS. Evaluation of kidney size in children: a pilot study of renal length as a surrogate of organ growth. *Korean J Pediatr* 2012;55:54-7.
12. Moorthy HK, Venugopal P. Measurement of renal dimensions in vivo: a critical appraisal. *Indian J Urol* 2011;27:169-75.
13. Prassopoulos P, Cavouras D. Renal parenchymal thickness in children measured by computed tomography. *Eur Urol* 1994;25:51-4.
14. Dinkel E, Ertel M, Dittrich M, Peters H, Berres M, Schulte-Wissermann H. Kidney size in childhood. Sonographical growth charts for kidney length and volume. *Pediatr Radiol* 1985;15:38-43.
15. Rosenbaum DM, Korngold E, Teele RL. Sonographic assessment of renal length in normal children. *AJR Am J Roentgenol* 1984;142:467-9.
16. Vujic A, Kosutic J, Bogdanovic R, Prijic S, Milicic B, Igrutinovic Z. Sonographic assessment of normal kidney dimensions in the first year of life: a study of 992 healthy infants. *Pediatr Nephrol* 2007;22:1143-50.
17. Otiv A, Mehta K, Ali U, Nadkarni M. Sonographic measurement of renal size in normal Indian children. *Indian Pediatr* 2012;49:533-6.
18. Sargent MA, Long G, Karmali M, Cheng SM. Interobserver variation in the sonographic estimation of renal volume in children. *Pediatr Radiol* 1997;27:663-6.
19. Ferrer FA, McKenna PH, Bauer MB, Miller SF. Accuracy of renal ultrasound measurements for predicting actual kidney size. *J Urol* 1997;157:2278-81.
20. Rossleigh MA, Farnsworth RH, Leighton DM, Yong JL, Rose M, Christian CL. Technetium-99m dimercaptosuccinic acid scintigraphy studies of renal cortical scarring and renal length. *J Nucl Med* 1998;39:1280-5.
21. Larsson SH, Aperia A. Renal growth in infancy and childhood: experimental studies of regulatory mechanisms. *Pediatr Nephrol* 1991;5:439-42.
22. Wallin L, Thörne J, Palmer J, Bajc M. Kidney size estimation in

- piglets using dimercaptosuccinic acid (DMSA) scintigraphy. *Clin Physiol* 1997;17:591-7.
23. Kim KS, Park JH. Sonographic assessment of renal size in normal children. *Korean J Nephrol* 1989;8:384-9.
 24. Kim IO, Cheon JE, Lee YS, Lee SW, Kim OH, Kim JH, et al. Kidney length in normal Korean children. *J Korean Soc Ultrasound Med* 2010;29:181-8.
 25. Akhavan A, Brajtford JS, McLeod DJ, Kabarriti AE, Rosenberg HK, Stock JA. Simple, age-based formula for predicting renal length in children. *Urology* 2011;78:405-10.
 26. Zerlin JM, Meyer RD. Sonographic assessment of renal length in the first year of life: the problem of "spurious nephromegaly". *Pediatr Radiol* 2000;30:52-7.
 27. Drougia A, Giapros V, Hotoura E, Papadopoulou F, Argyropoulou M, Andronikou S. The effects of gestational age and growth restriction on compensatory kidney growth. *Nephrol Dial Transplant* 2009;24:142-8.
 28. Mesrobian HG, Laud PW, Todd E, Gregg DC. The normal kidney growth rate during year 1 of life is variable and age dependent. *J Urol* 1998;160(3 Pt 2):989-93.
 29. Schmidt IM, Main KM, Damgaard IN, Mau C, Haavisto AM, Chellakooty M, et al. Kidney growth in 717 healthy children aged 0-18 months: a longitudinal cohort study. *Pediatr Nephrol* 2004;19:992-1003.
 30. Ganesh R, Vasanthi T, Lalitha J, Rajkumar J, Muralinath S. Correlation of renal length with somatic variables in Indian children. *Indian J Pediatr* 2010;77:326-8.