

RESEARCH ARTICLE

Clinical Study on Carboplatin for Treating Pediatric Patients with Wilms Tumors

Yong Zhang¹, Ling-Li Sun¹, Tao Li^{1*}, Hui Sun¹, Guo-Jia Mao²

Abstract

This analysis was conducted to evaluate the efficacy and safety of carboplatin based chemotherapy in treating pediatric patients with Wilms tumors. **Methods:** Clinical studies evaluating the efficacy and safety of carboplatin based regimens on response and safety for pediatric patients with Wilms tumors were identified using a predefined search strategy. Pooled response rates (RRs) of treatment were calculated. **Results:** In carboplatin based regimens, 4 clinical studies which including 127 patients with advanced Wilms tumors were considered eligible for inclusion. With this carboplatin based chemotherapy, 2 clinical studies included carboplatin, ifosfamide and etoposide. Systemic analysis suggested that, in all patients, the pooled PR was 64.5% (82/127) in carboplatin based regimens. Thrombocytopenia and leukocytopenia were the main side effects. No grade III or IV renal or liver toxicity was observed. No treatment related death occurred with carboplatin based treatment. **Conclusion:** This systemic analysis suggests that carboplatine based regimens are associated with a reasonable response rate and accepted toxicities for treating pediatric patients with Wilms tumors.

Keywords: Carboplatine - Wilms tumor - toxicity

Asian Pac J Cancer Prev, **15** (17), 7277-7280

Introduction

Combined treatment for patients with Wilms' tumor (WT) is an important factor contributing to the improvement of prognosis (Pritchard-Jones et al., 2003; Kalapurakal et al., 2004). It was reported that the response rate of children with relapsed or resistant WT to a combination of carboplatin, ifosfamide, and etoposide, at the maximal tolerated dose of 635 mg/m² for carboplatin, is associated with a CR rate of 32% and a total response rate of 63% (Kung et al., 1993; Kung et al., 1995). In the field of cancer chemotherapy including in the treatment of WT, carboplatin is a common choice (Akkuzu et al., 2013; Bag et al., 2013; Butthongkomvong et al., 2013; Fu et al., 2013; 2013; Khemapech et al., 2013; Kucukoztas et al., 2013; Natukula et al., 2013; Oranratanaphan et al., 2013; Ozdemir et al., 2013; Zhang et al., 2013; Pitakkarnkul et al., 2013). It was reported that carboplatin, at a dose of 550 mg/m² every 3 weeks, as a single drug resulted in a response rate of 53% in children with relapsed WT (de Camargo et al., 1994). Other agents, e.g., etoposide and ifosfamide also demonstrated activity (Tournade et al., 1988; Pein et al., 1993; de Camargo et al., 1994), a response rate of 50% to ifosfamide (3 gm/m² over 2 days, every 2 weeks) in children with relapsed WT was also reported (Tournade et al., 1988). The use of etoposide (200 mg/m²/day for 5 days) as a single agent in relapsed WT was previous reported in a response rate of 42% (Pein et al., 1993). However, the role of carboplatin in treating patients with WT has not been systemically investigated.

According to this background, we hypothesize that carboplatin originated regimen could be established as an optimal schedule for treating pediatric patients with WT.

Materials and Methods

Search strategy

We searched PUBMED, by using the following search term: (carboplatin) and (Wilms tumor). All clinical studies evaluating the impact of carboplatin on the response or survival and side effects for WT published in English prior to Jun 1st of 2014 were identified. If samples of two studies overlap, only the latest one was included. Additional articles were obtained from references within the articles identified by the electronic search. We did not consider meeting abstracts or unpublished reports.

Inclusion and exclusion criteria

We reviewed abstracts of all citations and retrieved studies. The following criteria were used to include published studies: (1) clinical studies, combined with cyclophosphamide or etoposide; (2) The study was performed in accordance with the Helsinki Declaration (1964, amended in 1975 and 1983) of the World Medical Association. Eligibility criteria included histologically or cytologically verified metastatic and/or locally advanced WT, the presence of at least one bidimensionally measurable lesion, a performance status (WHO) 2, age < 18 years. Studies were excluded if one of the following existed: (1) duplicate data; (2) no sufficient data were reported.

¹Shiyuan Taihe Hospital of Hubei Province Department of Pediatrics, The Affiliated Hospital of Hubei Medical College, Shiyuan, ²Department of Surgery, Jiujiang Third People's Hospital of Jiangxi Province, Jiujiang, China *For correspondence: zhyong885900@sohu.com

Data collection and analysis

Selection of trials and data extraction: The titles and abstracts of publications identified according to the above search strategy were assessed independently for inclusion by two authors; the full text was selected for further assessment if the abstract suggests relevance. Disagreement was resolved by discussion. Data was extracted by independent authors. The following recorded data were extracted: author, publication data, and country of the first or corresponding author, the number of patients.

Results

There were 68 papers relevant to the search words by the end of June, 2014. Via steps of screening the title and reading the abstract, 4 studies were identified (de Camargo et al., 1994; Zoubek et al., 1995; Abu-Ghosh et al., 2002; Kung et al., 1995) when carboplatin was used as single or in combination of chemotherapy. These studies had been carried out in Brazil, European countries, and the United States. The following outcomes were presented in at least all studies and extracted for combined analysis: response rate, including the rate of complete or partial response (CR or PR) and toxicities.

Characteristics of carboplatin as a single agent of chemotherapy, studies included are presented as short-term outcomes: the response rate of de Camargo et al. was 53.3% (8/15), of Zoubek et al. (1995) was 77.8% (7/9). When carboplatin was used in combination with ifosfamide and etoposide, 2 studies included in this study are presented and the short-term outcomes suggested that the response rate of Abu-Ghosh et al. was 81.8%, of Kung et al. was 63%. Totally, 127 patients were enrolled and 82 patients achieved CR or PR, the pooled response rate thus was 64.5% (82/127) in carboplatin based regimens.

Thrombocytopenia and leukocytopenia were the main side effects. No grade III or IV renal or liver toxicity were observed. No treatment related death occurred in carboplatin based treatment.

Discussion

In previous studies, investigators have reported good response rates of carboplatin in treating patients with relapsed WT (Marina et al., 1994; Marina et al., 1994). But, it remains to be determined whether the higher doses of chemotherapy containing carboplatin could be compared with that of lower doses and what combination should be considered. Especially, consideration should be focused on the minimal instead of maximal dose needed to cure WT (O'Dwyer et al., 1985; Grundy et al., 1989). The use of two agents combination regimen containing carboplatin and vp-16 in treating patients with refractory or relapsed WT for two courses achieved a response rate of 73% (Pein et al., 1994). However, only 31% of patients remain disease-free and in CR at a median of 40 months of follow-up. Further, three-agent regimens containing carboplatin, e.g. the use of ICE for patients in this setting is associated with a response rate of 71% (Kung et al., 1993; Kung et al., 1995). The use of high-dose chemotherapy followed by autologous

stem cell transplantation may be effective in salvaging patients with relapsed WT and poor prognostic features. A 3-year disease-free survival and overall survival of $50 \pm 17\%$ and $60 \pm 18\%$, respectively, were reported after high-dose chemotherapy and autologous stem cell transplantation in a group of patients with high-risk relapsed WT transplanted in post-relapse CR or PR (Pein et al., 1998). It was reported that carboplatin, at a dose of 550 mg/m^2 every 3 weeks, as a single drug resulted in a response rate of 53% in children with relapsed WT (de Camargo et al., 1994). When the dose of chemotherapy is increased toxicities are invariably increased. Usually, hematopoietic toxicity is the major factor limiting the use of more dose-intensive chemotherapy regimens for children with recurrent/relapsed WT. The incidence of grade III/IV neutropenia was 81% for patients treated with ifosfamide/etoposide (ifosfamide at $2000 \text{ mg/m}^2/\text{day}$ for 3 days, etoposide at $100 \text{ mg/m}^2/\text{day}$ for 3 days) and the incidence of grade III/IV thrombocytopenia was 25.5% (Kung et al., 1993). Similarly, Miser et al. reported a high incidence of grade III/IV neutropenia (96% of cycles) following the ifosfamide/etoposide combination and a lower incidence of thrombocytopenia (20-30%) (Miser et al., 1987; Miser et al., 1993). The carboplatin/etoposide combination could result in a 92% incidence of grade III/IV neutropenia, and a higher incidence of grade IV thrombocytopenia when compared with ifosfamide/etoposide regimens (Pein et al., 1994). The combination of ICE therapy in pediatric patients is reported to bear a similar grade III/IV neutropenia (83-84%) as reported by both POG and CCG (Kung et al., 1993; Kung et al., 1995; Cairo et al., 2001). Grade III/IV thrombocytopenia was 59 and 84% in both POG and CCG study groups, respectively. Thus, consideration should be fully paid to the condition of patients.

In previous, it was identified that subgroups of relapsed WT patients with poor prognostic had features, e.g., unfavorable histology, abdominal recurrence with previous radiation therapy, advanced disease stage, early recurrence (<12 months) and previous three-drug therapy (Grundy et al., 1989). Recently Pein et al. reported the results of a multivariate analysis of eight adverse prognostic factors in children with relapsed WT and showed a significant effect on 3-year survival of unfavorable histology, early recurrence ≤ 6 months, multiple organ recurrence and lymph node involvement at recurrence (Pein et al., 1999). The use of ICE chemotherapy in previous study resulted in a high response rate of 82% with long-term overall and progression-free survival ($63.6 \pm 14.5\%$) of all the patients who attained a CR initially or subsequently, either following further chemotherapy, radiotherapy, or high-dose chemotherapy and PBSC transplantation. The dose-intensive therapy allows patients with high-risk disease and poor prognostic features to enter into a CR or a PR with the probability of further therapy with newer therapeutic agents and/or novel therapeutic approaches (i.e. high-dose chemotherapy followed by PBSC transplantation). Although both patients with unfavorable histology had a poor outcome a larger number of patients with unfavorable histology are needed to better evaluate the efficacy of ICE in this subgroup. In

comparison, Tannous et al. recently reported the results of an intergroup CCG and POG trial of a retrieval study in children with relapsed WT (Tannous et al., 2000). High-risk patients were treated with chemotherapy regimens containing carboplatin (1 gm/m²)/etoposide (300 mg/m²) and cyclophosphamide (2200 mg/m²)/etoposide (500 mg/m²) during induction and maintenance in addition to local radiotherapy and surgery. CR was seen in 42% of patients and PR in 36% following two cycles of induction therapy (overall response rate 78%). Patients in CR after surgery were treated with maintenance chemotherapy cycles of alternating cyclophosphamide/etoposide and carboplatin/etoposide. The reported event-free survival and survival in this group of patients was 59 and 64%, respectively. These results suggested that patients with different risk factors should be treated with different regimens.

Our study suggested that when carboplatin was used in combination with ifosfamide and etoposide, 2 studies included in this study are presented and the short-term outcomes suggested that the response rate of Abu-Ghosh et al. was 81.8%, of Kung et al. was 63%. Totally, 127 patients were enrolled and 82 patients achieved CR or PR, the pooled response rate thus was 64.5% (82/127) in carboplatin based regimens. Thrombocytopenia and leukocytopenia were the main side effects. No grade III or IV renal or liver toxicity were observed. No treatment related death occurred in carboplatin based treatment. Thus, we recommend that patients with low risk factors should be treated with single carboplatin or carboplatin combined with another agent, e.g., etoposide or ifosfamide; patients with high risk factors and good tolerability could be treated with carboplatin combined with other two agents.

In conclusion, this systemic analysis suggests that carboplatine based regimens are associated with reasonable response rate and accepted toxicities for treating pediatric patients with Wilms tumor.

References

- Abu-Ghosh AM, Krailo MD, Goldman SC, et al (2002). Ifosfamide, carboplatin and etoposide in children with poor-risk relapsed Wilms' tumor: a Children's Cancer Group report. *Ann Oncol*, **13**, 460-9.
- Akkuzu G, Ayhan A (2013). Sexual functions of Turkish women with gynecologic cancer during the chemotherapy process. *Asian Pac J Cancer Prev*, **14**, 3561-4.
- Anderson B, Krailo M, Davenport V et al (1999). Ifosfamide, carboplatin and etoposide (ICE) is a superb retrieval regimen in a large cohort (n = 107) of children with recurrent/refractory soft tissue sarcomas (STS): the Children's Cancer Group (CCG) experience. *J Pediatr Hematol Oncol*, **21**, 337 (Abstr 505).
- Bag A, Pant NK, Jeena LM, et al (2013). GSTT1 null and MPO -463G polymorphisms and carboplatin toxicity in an Indian population. *Asian Pac J Cancer Prev*, **14**, 4739-42.
- Butthongkomvong K, Sirachainan E, Jhankumpha S, et al (2013). Treatment outcome of palliative chemotherapy in inoperable cholangiocarcinoma in Thailand. *Asian Pac J Cancer Prev*, **14**, 3565-8.
- Cairo MS, Krailo M, Miser J et al (1996). Excellent results of the use of ifosfamide, carboplatin, etoposide (ICE) in children with recurrent or refractory solid tumors. *Proc Am Soc Clin Oncol*, **15**, 468 (Abstr 1474).
- Cairo MS, Krailo MD, Weinthal JA et al (1998). A phase I study of granulocyte-macrophage-colony stimulating factor/interleukin-3 fusion protein (PIXY321) following ifosfamide, carboplatin, and etoposide therapy for children with recurrent or refractory solid tumors: a report of the Children's Cancer Group. *Cancer*, **83**, 1449-60.
- Cairo MS, Shen V, Krailo MD et al (2001). Prospective randomized trial between two doses of granulocyte colony-stimulating factor after ifosfamide, carboplatin, and etoposide in children with recurrent or refractory solid tumors: a Children's Cancer Group Report. *J Pediatr Hematol Oncol*, **23**, 30-8.
- Castello MA, Clerico A, Jenkner A, Dominici C. A pilot study of high-dose carboplatin and pulsed etoposide in the treatment of childhood solid tumors (1990). *Pediatr Hematol Oncol*, **7**, 129-35.
- Curtis BM, Williams DE, Broxmeyer HE et al (1991). Enhanced hematopoietic activity of a human granulocyte/macrophage colony-stimulating factor-interleukin 3 fusion protein. *Proc Natl Acad Sci USA*, **88**, 5809-13.
- D'Angio GJ, Breslow N, Beckwith JB et al (1989). Treatment of Wilms' tumor. Results of the Third National Wilms' Tumor Study. *Cancer*, **64**, 349-60.
- Daw N, Jones D, Gregornik D et al (1999). Renal function after ifosfamide, carboplatin and etoposide (ICE), nephrectomy, and radiation in children with high risk Wilms' tumor. *Proc Am Soc Clin Oncol*, **18**, 560 (Abstr 2159).
- De Camargo B, Melaragno R, Saba e Silva N et al (1994). Phase II study of carboplatin as a single drug for relapsed Wilms' tumor: experience of the Brazilian Wilms' Tumor Study Group. *Med Pediatr Oncol*, **22**, 258-60.
- Ettlinger LJ, Gaynon PS, Krailo MD et al (1994). A phase II study of carboplatin in children with recurrent or progressive solid tumors. A report from the Children's Cancer Group. *Cancer*, **73**, 1297-301.
- Frappaz D, Michon J, Hartmann O et al (1992). Etoposide and carboplatin in neuroblastoma: a French Society of Pediatric Oncology phase II study. *J Clin Oncol*, **10**, 1592-601.
- Fu JH, Gao Z, Ren CC, et al (2013). Comparison of clinical efficacy of three different neoadjuvant approaches (chemotherapy combined vaginal intracavitary irradiation, neoadjuvant chemotherapy alone or radiotherapy) combined with surgery for patients with stage Ib2 and IIa2 cervical cancer. *Asian Pac J Cancer Prev*, **14**, 2377-81.
- Genet JC, Doz F, Bouffet E et al (1994). Carboplatin and VP 16 in medulloblastoma: a Phase II Study of the French Society of Pediatric Oncology (SFOP). *Med Pediatr Oncol*, **23**, 422-7.
- Green DM, Beckwith JB, Breslow NE et al (1994). Treatment of children with stages II to IV anaplastic Wilms' tumor: a report from the National Wilms' Tumor Study Group. *J Clin Oncol*, **12**, 2126-31.
- Green DM, Breslow NE, Evans I et al (1996). Treatment of children with stage IV favorable histology Wilms' tumor: a report from the National Wilms' Tumor Study Group. *Med Pediatr Oncol*, **26**, 147-52.
- Grundt P, Breslow N, Green DM et al (1989). Prognostic factors for children with recurrent Wilms' tumor: results from the Second and Third National Wilms' Tumor Study. *J Clin Oncol*, **7**, 638-47.
- Harrap KR (1985). Preclinical studies identifying carboplatin as a viable cisplatin alternative. *Cancer Treat Rev*, **12**, 21-33.
- Kalapurakal JA, Dome JS, Perlman EJ, et al (2004). Management of Wilms' tumour: current practice and future goals. *Lancet Oncol*, **5**, 37-46.

- Kaplan EL, Meier P (1958). Non-parametric estimation from incomplete observations. *J Am Stat Assoc*, **53**, 457-81.
- Khemapech N, Oranratanaphan S, Termrungruanglert W, et al (2013). Salvage chemotherapy in recurrent platinum-resistant or refractory epithelial ovarian cancer with Carboplatin and distearoylphosphatidylcholine pegylated liposomal Doxorubicin (lipo-dox®). *Asian Pac J Cancer Prev*, **14**, 2131-5.
- Kucukoztas N, Dizdar O, Rahatli S, et al (2013). Impact of treatment strategies on local control and survival in uterine carcinosarcomas in Turkey. *Asian Pac J Cancer Prev*, **14**, 2959-62.
- Kung F, Hayes FA, Krischer J et al (1988). Clinical trial of etoposide (VP-16) in children with recurrent malignant solid tumors. A phase II study from the Pediatric Oncology Group. *Invest New Drugs*, **6**, 31-6.
- Kung FH, Bernstein ML, Camitta BM et al (1999). Ifosfamide/carboplatin/etoposide (ICE) in the treatment of advanced, recurrent Wilms' tumor. *Proc Am Soc Clin Oncol*, **18**, 559 (Abstr 2156).
- Kung FH, Desai SJ, Dickerman JD et al (1995). Ifosfamide/carboplatin/etoposide (ICE) for recurrent malignant solid tumors of childhood: a Pediatric Oncology Group phase I/II study. *J Pediatr Hematol Oncol*, **17**, 265-9.
- Kung FH, Pratt CB, Vega RA et al (1993). Ifosfamide/etoposide combination in the treatment of recurrent malignant solid tumors of childhood. A Pediatric Oncology Group phase II study. *Cancer*, **71**, 1898-903.
- Lebwohl D, Canetta R (1998). Clinical development of platinum complexes in cancer therapy: an historical perspective and an update. *Eur J Cancer*, **34**, 1522-34.
- Loehrer PJ Sr (1992). The history of ifosfamide. *Semin Oncol*, **19**, 2-6.
- Marina NM, Poquette CA, Cain AM et al (2000). Comparative renal tubular toxicity of chemotherapy regimens including ifosfamide in patients with newly diagnosed sarcomas. *J Pediatr Hematol Oncol*, **22**, 112-8.
- Marina NM, Rodman JH, Murry DJ et al (1994). Phase I study of escalating targeted doses of carboplatin combined with ifosfamide and etoposide in treatment of newly diagnosed pediatric solid tumors. *J Natl Cancer Inst*, **86**, 544-8.
- Marina NM, Wilimas JA, Meyer WH et al (1994). Refining therapeutic strategies for patients with resistant Wilms' tumor. *Am J Pediatr Hematol Oncol*, **16**, 296-300.
- Miser J, Krailo M, Hammond GD (1993). The combination of ifosfamide (IFOS), etoposide (VP16), and mesna (M): a very active regimen in the treatment of recurrent Wilms' tumor (WT). *Proc Am Soc Clin Oncol*, **12**, 417 (Abstr 1432).
- Miser JS, Kinsella TJ, Triche TJ et al (1987). Ifosfamide with mesna uroprotection and etoposide: an effective regimen in the treatment of recurrent sarcomas and other tumors of children and young adults. *J Clin Oncol*, **5**, 1191-8.
- Natukula K, Jamil K, Pingali UR, et al (2013). Survival analysis in advanced non small cell lung cancer treated with platinum based chemotherapy in combination with paclitaxel, gemcitabine and etoposide. *Asian Pac J Cancer Prev*, **14**, 4661-6.
- O'Dwyer PJ, Leyland-Jones B, Alonso MT et al (1985). Etoposide (VP-16-213). Current status of an active anticancer drug. *N Engl J Med*, **312**, 692-700.
- Oranratanaphan S, Khemapech N (2013). Characteristics and treatment outcomes of patients with malignant transformation arising from mature cystic teratoma of the ovary: experience at a single institution. *Asian Pac J Cancer Prev*, **14**, 4693-7.
- Ozdemir O, Ozdemir P, Veral A, et al (2013). ERCC1 expression does not predict survival and treatment response in advanced stage non-small cell lung cancer cases treated with platinum based chemotherapy. *Asian Pac J Cancer Prev*, **14**, 4679-83.
- Pappo AS, Etcubanas E, Santana VM et al (1993). A phase II trial of ifosfamide in previously untreated children and adolescents with unresectable rhabdomyosarcoma. *Cancer*, **71**, 2119-25.
- Pein F, Michon J, Valteau-Couanet D et al (1998). High-dose melphalan, etoposide, and carboplatin followed by autologous stem-cell rescue in pediatric high-risk recurrent Wilms' tumor: a French Society of Pediatric Oncology study. *J Clin Oncol*, **16**, 3295-301. Abstract
- Pein F, Pinkerton R, Tournade MF et al (1993). Etoposide in relapsed or refractory Wilms' tumor: a phase II study by the French Society of Pediatric Oncology and the United Kingdom Children's Cancer Study Group. *J Clin Oncol*, **11**, 1478-81.
- Pein F, Rey A, de Kraker J et al (1999). Multivariate analysis of adverse prognostic factors (APF) in children with recurrent (Rec) Wilms' tumor (WT) after initial treatment according to SIOP-6 or SIOP-9 strategies. *Med Pediatr Oncol*, **33**, 170 (Abstr 0111).
- Pein F, Tournade MF, Zucker JM et al (1994). Etoposide and carboplatin: a highly effective combination in relapsed or refractory Wilms' tumor-a phase II study by the French Society of Pediatric Oncology. *J Clin Oncol*, **12**, 931-36.
- Pinkerton CR, Rogers H, James C et al (1985). A phase II study of ifosfamide in children with recurrent solid tumours. *Cancer Chemother Pharmacol*, **15**, 258-62.
- Pitakkarnkul S, Tangjitgamol S, Srijaipracharoen S, et al (2013). Treatment outcomes of paclitaxel for refractory or recurrent epithelial ovarian cancer patients in Thailand. *Asian Pac J Cancer Prev*, **14**, 2421-7.
- Pritchard-Jones K, Kelsey A, Vujanic G, Imeson J, et al (2003). : Older age is an adverse prognostic factor in stage I, favorable histology Wilms' tumor treated with vincristine monotherapy: a study by the United Kingdom Children's Cancer Study Group, Wilm's Tumor Working Group. *J Clin Oncol*, **21**, 3269-75.
- Rose WC, Schurig JE (1985). Preclinical antitumor and toxicologic profile of carboplatin. *Cancer Treat Rev*, **12**, 1-19.
- Shen V, Morris E, Davenport V et al (1999). Ifosfamide, carboplatin and etoposide (ICE) is an excellent retrieval regimen in children with recurrent/refractory CNS malignancies: the Children's Cancer Group (CCG) experience. *Proc Am Soc Clin Oncol*, **18**, 558 (Abstr 2151).
- Tannous R, Giller R, Holmes E et al (2000). Intensive therapy for high risk (HR) relapsed Wilms' tumor (WT). A CCG-4921/POG-9445 study report. *Proc Am Soc Clin Oncol*, **19**, 2315 (Abstr 2315).
- Tournade MF, Lemerle J, Brunat-Mentigny M et al (1988). Ifosfamide is an active drug in Wilms' tumor: a phase II study conducted by the French Society of Pediatric Oncology. *J Clin Oncol*, **6**, 793-96.
- Zhang YY, Gu KS (2013). DNA repair capacity in peripheral blood lymphocytes predicts efficacy of platinum-based chemotherapy in patients with gastric cancer. *Asian Pac J Cancer Prev*, **14**, 5507-12.