

# Seroprevalence and Potential Risk Factors Associated with *Neospora* spp. Infection among Asymptomatic Horses in Jordan

Abdelsalam Q. Talafha<sup>1,\*</sup>, Sameeh M. Abutarbush<sup>1</sup>, David L. Rutley<sup>2</sup>

<sup>1</sup>Department of Veterinary Clinical Sciences, Faculty of Veterinary Medicine, Jordan University of Science and Technology, Irbid 22110, Jordan;

<sup>2</sup>School of Animal and Veterinary Science, The University of Adelaide, South Australia, Australia

**Abstract:** This study aimed to determine the seroprevalence and to identify risk factors associated with *Neospora* spp. infection in horses in Jordan. Management related data were collected from each farm and individual horses. Sera from 227 horses from 5 of 6 climatic regions in Jordan were analyzed for the presence of antibodies to *Neospora* spp. by ELISA kit. The study was performed during spring of 2010. The association between seropositivity and risk factors was analyzed. A total of 7 (3%) of 227 sera had antibodies for *Neospora* spp. There was a significant regional difference ( $P=0.018$ ) between the 5 climatic regions. Positive cases were located in Amman and Irbid, while the other regions (Zarqa, Jordan Valley, and Wadi Mousa) had zero prevalence. The use of anthelmintics at least once a year resulted in a significant reduction of the seroprevalence to *Neospora* spp. (1.6% vs 9.8%). However, this might be a phenomenon by chance and a better hygiene since owners can invest in anthelmintics. Other risk factors such as age, gender, breed, usage, body condition score, grazing, presence of other animals mixed with the horses in the same property, and a history of previous diseases were not significantly associated with the seroprevalence to *Neospora* spp. infection. This is the first study to report on the presence of *Neospora* seropositive horses in Jordan. Further studies are warranted to better understand the role of certain risk factors in the transmission of *Neospora* spp. among horse population and to determine which *Neospora* spp. are responsible for the infection.

**Key words:** *Neospora* spp., horse, risk factor, seroprevalence, Jordan

## INTRODUCTION

Equine neosporosis is caused by an obligatory intracellular apicomplexan protozoan parasite, *Neospora caninum*. *N. caninum* belongs to the family Sarcocystidae and closely resembles *Toxoplasma gondii* [1]. *Neospora hughesi* is a second species of *Neospora* described in the horse [2]. The parasite infects a broad range of animals including cattle, sheep, goats, deer, horses, and dogs. Although *N. caninum* is an important cause of abortion in cows, as well as various congenital abnormalities in dogs, little is known regarding its pathogenicity and transmission in horses [3]. Exposure to *Neospora* spp. in horses is not uncommon, clinical disease associated with natural infections in adult horses, however, have been reported in only few cases. Moreover, it is uncertain whether *N. caninum*, *N.*

*hughesi*, or both are responsible for the disease in positive cases since both species cross-react serologically [4]. In horses, infection with *Neospora* spp. has been associated with neurological disorders, neonatal diseases, and abortion [5]. Various laboratory techniques have been used to diagnose *N. caninum*-infected animals. Immunohistochemistry is used to demonstrate *N. caninum* in the placenta or fetal tissues [6]. Serology tests such as indirect fluorescent antibody test (IFAT), immunoblotting (IB), direct agglutination tests, and a wide variety of ELISAs are used to detect specific antibodies in sera of infected animals. In addition, DNA biotechnology can be used to distinguish *N. hughesi* from *N. caninum* [7].

According to the Jordan Ministry of Agriculture Yearly Statistical Data (2007), the total population of horses is 2,182 heads. In Jordan, the prevalence and risk factors associated with *N. caninum* infection has been reported in dairy herds [8]. Information regarding the prevalence of *Neospora* spp. in horses is lacking, therefore, the purpose of the study reported here was to determine the seroprevalence of *Neospora* spp. in horses and to identify potential risk factors for neosporosis in horses in Jordan.

•Received 26 November 2014, revised 19 January 2015, accepted 25 January 2015.

\*Corresponding author (talafha@just.edu.jo)

© 2015, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## MATERIALS AND METHODS

### Study animals and design

Total 227 clinically normal horses were enrolled in the study. The horse farms were selected randomly using the records of the Jordanian Ministry of Agriculture. Horses from each farm were selected randomly using a table of random digits. Horses sampled for the study were from 5 of 6 climatic regions in Jordan; Amman, Irbid, Zarqa, Jordan Valley, and Wadi Mousa (Fig. 1). The 6th region was not sampled due to the lack of horse population. About 40-50 horses were sampled from each region. The study was performed during spring of 2010. Each farmer was interviewed to gather information about each horse, farm characteristics, and herd management. Table 1 shows a list of specific data that was collected on each horse and the farm included in the study.

### Blood samples

Whole blood (5 to 10 ml) was collected from the jugular vein of each horse using vacuum plain tubes (Aysset tube®, Adana, Turkey) and transported on ice to the laboratory within 2

hr. Blood was centrifuged at 5,000 g for 10 min, and sera were harvested and stored in microtubes at -20°C until analyzed.

### Laboratory analysis


For determination of *Neospora* spp. seroprevalence, antibodies in sera were detected by an indirect ELISA kit (The Herd-Chek Anti-*Neospora caninum* Antibody Test Kit®, IDEXX Laboratories, Westbrook, Maine, USA) according to the manufacturer's instructions. The test has a known sensitivity and specificity of 98.6% and 98.9%, respectively. The resulted prevalence was adjusted to the test sensitivity and specificity using the formula published previously [9].

### Statistical analysis

The seroprevalence and risk factor analysis of *Neospora* spp. infection were determined using the chi square tests of association [10]. Variables with a *P*-value of < 0.05 were considered statistically significant.

## RESULTS

Table 1 shows the distribution of *Neospora* spp. positive and negative horses (*n* = 227) from Jordan with various risk factors that might be associated with the seropositivity to equine neosporosis. The overall seroprevalence of *Neospora* spp. was 3% (7/227 horses). There was a significant regional difference (*P* = 0.018) with the positive cases belonging to Amman and Irbid. The following risk factors were not significantly (*P* > 0.05) associated with seroprevalence of *Neospora* spp.: horse age, gender, breed, usage, body condition score on a scale of 1-9 (a score of 1 represents emaciation whereas a score of 9 is a profound obesity), grazing, presence of other animals mixed with the horses in the same property, and a history of previous diseases such as gastrointestinal (colic and diarrhea), respiratory (fever, coughing, naso-ocular discharge), integumentary (dermatitis), and musculoskeletal diseases (lameness) since 1 year before the commencement of the study. Of 227 horses, 186 (82%) received anthelmintic medication at least once or more a year. Ivermectin and/or piperazine were the most commonly used anthelmintics. Greater than or equal to 1 deworming per year was significantly (*P* = 0.0017) associated with a reduction in the seroprevalence of *Neospora* spp. (1.6% vs 9.8%). This finding was supported by an effect of any drench reducing the seroprevalence of *Neospora* spp. (*P* = 0.0063). Of 186 horses drenched, only 3 cases (1.6%) of *Neospora* infection were iden-



**Fig. 1.** Map of Jordan divided into 6 climate regions according to the model of Koppen (anon., 1984). It also indicates the location of 5 sites sampled in this study. 1. Warm temperature rainy (Irbid), 2. Cool steppe (Amman), 3. Cool desert (Zarqa), 4. Warm desert (Jordan Valley), 5. Cool temperate rainy (Wadi Mousa), and 6. Warm steppe (not sampled).

**Table 1.** Regional distribution of *Neospora* spp. positive and negative horses (n=227) from Jordan and the risk factors analysis associated with the occurrence of equine neosporosis

Variable	Description		<i>Neospora</i> spp.(%)	
	N	%	Positive	Negative
Location (climatic region)*				
Amman	46	20.3	3 (6.5)	43
Irbid	41	18	4 (9.8)	37
Zarqa	45	19.8	0 (0)	45
Jordan Valley	49	21.6	0 (0)	49
Wadi Mousa	46	20.3	0 (0)	46
Age (year)				
Juvenile <3	21	9.3	1 (4.7)	20
Adult 3-10	142	62.6	5 (3.5)	137
old > 10	64	28.1	1 (1.6)	63
Gender				
Female	96	42.3	1 (1)	95
Male and gelding	131	57.7	6 (4.6)	125
Breed				
Mixed local	171	75.3	5 (3)	166
Thoroughbred	56	24.7	2 (3.6)	54
Horse usage				
Competitive (racing, polo, showing, and jumping)	121	53.3	2 (1.6)	119
Non-competitive (breeding, working)	106	46.7	5 (4.7)	101
Body condition score (scale 1-9)				
Under condition <4	18	8	1 (5.5)	17
Normal condition 4-5.5	180	79.3	6 (3.3)	174
Over condition > =6	29	12.7	0 (0)	29
Grazing				
Yes	62	27.3	2 (3.2)	60
No	165	72.7	5 (3)	160
Presence of other animals (ruminants, dogs, and donkeys) mixed-in with horses				
Yes	121	53.3	5 (4.1)	116
No	106	46.7	2 (1.9)	104
History of previous body systems diseases				
Yes	88	38.8	3 (3.4)	85
No	139	61.2	4 (2.9)	135
Use of any anthelmintic medication*				
Yes	186	82	3 (1.6)	183
No	41	18	4 (9.8)	37
Number of dewormings since 1 year before start of study (nil to 9)				
Nil	41	18.1	4 (9.8)	37
1	47	20.7	1 (2.1)	46
>1	139	61.2	2 (1.4)	137

\*P-value &lt;0.05.

tified. In contrast, the other 4 cases of *Neospora* infection were identified from 41 horses (9.8%) that had not been drenched for worms. Testing the association of specific anthelmintic drugs with the seroprevalence of *Neospora* spp. was not possible due to the low seroprevalence of *Neospora*.

## DISCUSSION

Neosporosis in horses has been reported from different

parts of the world. To the authors' knowledge, this is the first study in Jordan to determine the prevalence of antibodies to *Neospora* spp. and to evaluate risk factors associated with the occurrence of neosporosis in horses. Unfortunately, we were unable to determine which *Neospora* spp. were infected the seropositive horses since both species cross-react serologically [4]. The results of this study indicated that horses in Jordan are exposed to *Neospora* spp. with an overall seroprevalence of 3%. This percentage is considerably lower than the seroprevalence

of *Neospora* infection that have been reported in horses from the United States [11] and Czech Republic [12]. In the Middle East, seroprevalence of *Neospora* spp. was reported from Israel [13] and Saudi Arabia [14]. The seroprevalence of *Neospora* spp. infection in horses is considerably different among and within countries. These variations might be due to differences in the serological tests used in each study, limitations of the testing methods, and non-standardized controls, and cut-off values applied [15]. In addition, the study design, criteria for sample collection, different levels of exposure to the various risk, and protective factors for infection or disease might influence the seroprevalence of *Neospora* spp. which make the comparison between results from different studies difficult [16].

Determining various risk factors and understanding their role in disease transmission and epidemiology is critical for the development and implementation of proper measures to control equine neosporosis. The seroprevalence of *Neospora* spp. was higher in Amman and Irbid compared to other climatic regions of the country. It was noticed that these 2 regions have the highest density of horse population in Jordan. Significant differences in the seroprevalence of *Neospora* spp. among horse groups from various geographical regions were reported from USA [15]. However, no significant difference was found in the seropositivity rates from 4 districts in Niğde province of Turkey [17].

The present study showed a non-significant decrease in the seroprevalence of *Neospora* spp. with increased horse age. This result is contrary to the previous report from Israel [13] which found a significantly higher seroprevalence to *Neospora* spp. as horses get older than 10 years of age (because of an increased likelihood of exposure through horizontal transmission over time). In a Turkish study, a non-significant increase of seroprevalence to *Neospora* spp. (22.2% and 27.2%) was found between 2 age groups; 1-10 and 11-20 years old, respectively [17]. The absence of significant effect for the gender on the seroprevalence of *Neospora* spp. in this study is in agreement with findings of Villalobos et al. [18]. The horse breed was also not significantly associated with the seroprevalence to *Neospora* spp. which agrees with the study of Kligler et al. [13] in Israel. The presence of other animals mixed-in with horses in the same property was also not significantly associated with seroprevalence to *Neospora* spp. In cattle, it is generally accepted that the presence of farm dogs, the definitive host for *N. caninum*, increases the chance of *N. caninum* infection [19]. The present study showed that the use of anthelmintic medication at least

once a year resulted in a significant reduction in the seroprevalence of *Neospora* spp. However, this might be a phenomenon by chance and a better hygiene since owners can invest in anthelmintics. In cattle, various antimicrobial agents have been tested against *N. caninum* in vitro but there is still no safe and effective chemotherapy that clears infection completely [20]. However, in vitro and in vivo experimental studies have shown a promising effect for toltrazuril and ponazuril on tachyzoites of *N. caninum* in calves [21]. Further studies are necessary to determine which species, *N. caninum* or *N. hughesi*, infect the horses. Furthermore, detailed studies to better understand various risk factors that affect the transmission of the parasite among horses are warranted.

## ACKNOWLEDGMENTS

We are highly thankful to the horse owners for their willingness to participate and to provide information about their animals. The study was financially supported by the Deanship of Research, Jordan University of Science and Technology, Irbid, Jordan.

## CONFLICT OF INTEREST

We declare that we have no conflict of interest.

## REFERENCES

1. Dubey JP, Barr BC, Barta JR, Bjerkås I, Björkman C, Blagburn BL, Bowman DD, Buxton D, Ellis JT, Gottstein B, Hemphill A, Hill DE, Howe DK, Jenkins MC, Kobayashi Y, Koudela B, Marsh AE, Mattsson JG, McAllister MM, Modrý D, Omata Y, Sibley LD, Speer CA, Trees AJ, Uggla A, Upton SJ, Williams DJ, Lindsay DS. Redescription of *Neospora caninum* and its differentiation from related coccidia. *Int J Parasitol* 2002; 32: 929-946.
2. Marsh AE, Barr BC, Packham AE, Conrad PA. Description of a new *Neospora* species (Protozoa: Apicomplexa: Sarcocystidae). *J Parasitol* 1998; 84: 983-991.
3. Pitel PH, Lindsay DS, Caure S, Romand S, Pronost S, Gargala G, Mitchell SM, Hary C, Thulliez P, Fortier G, Ballet JJ. Reactivity against *Sarcocystis neurona* and *Neospora* by serum antibodies in healthy French horses from two farms with previous equine protozoal meningoencephalitis-like cases. *Vet Parasitol* 2003; 111: 1-7.
4. Al-Qassab S, Reichel MP, Ivens A, Ellis JT. Genetic diversity amongst isolates of *Neospora caninum*, and the development of a multiplex assay for the detection of distinct strains. *Mol Cell*

- Probes 2009; 23: 132-139.
5. Lindsay DS. Neosporosis: an emerging protozoal disease of horses. *Equine Vet J* 2001; 33: 116-118.
  6. Lindsay DS, Dubey JP. Immunohistochemical diagnosis of *Neospora caninum* in tissue sections. *Am J Vet Res* 1989; 50: 1981-1983.
  7. de Waal T. Advances in diagnosis of protozoan diseases. *Vet Parasitol* 2012; 189: 65-74.
  8. Talafha AQ, Al-Majali AM. Prevalence and risk factors associated with *Neospora caninum* infection in dairy herds in Jordan. *Trop Anim Health Prod* 2013; 45: 479-485.
  9. Noordhuizen JP, Frankena K, van der Hoofd C, Graat EA. Application of Quantitative Methods in Veterinary Epidemiology. The Netherlands. Wageningen Co. Ltd Press. 1997.
  10. Kaps M, Lamberson W. Biostatistics for animal science. Wallingford, U.K. CAB International. 2004.
  11. Dubey JP, Romand S, Thulliez P, Kwok OC, Shen SK, Gamble HR. Prevalence of antibodies to *Neospora caninum* in horses in North America. *J Parasitol* 1999; 85: 968-969.
  12. Bártová E, Sedláčková K, Syrová M, Literák I. *Neospora* spp. and *Toxoplasma gondii* antibodies in horses in the Czech Republic. *Parasitol Res* 2010; 107: 783-785.
  13. Kligler EB, Shkap V, Baneth G, Mildenberg Z, Steinman A. Seroprevalence of *Neospora* spp. among asymptomatic horses, aborted mares and horses demonstrating neurological signs in Israel. *Vet Parasitol* 2007; 148: 109-113.
  14. Al Anazi AD, Mohamed AS. Seroprevalence of *Neospora* spp. in horses from Central Province of Saudi Arabia. *Afr J Biotechnol* 2013; 12: 982-985.
  15. Vardeleon D, Marsh AE, Thorne JG, Loch W, Young R, Johnson PJ. Prevalence of *Neospora hughesi* and *Sarcocystis neurona* antibodies in horses from various geographical locations. *Vet Parasitol* 2001; 95: 273-282.
  16. Jakubek EB, Lundén A, Uggla A. Seroprevalences of *Toxoplasma gondii* and *Neospora* sp. infections in Swedish horses. *Vet Parasitol* 2006; 138: 194-199.
  17. Karatepe M, Karatepe B. Investigation of seroprevalence of *Neospora* spp. in Horses in Niğde Province (Turkey). *Kafkas Univ Vet Fak Derg* 2012; 18: A39-A42.
  18. Villalobos EM, Ueno TE, de Souza SL, Cunha EM, do Carmo Custódio de Souza Hunold Lara M, Gennari SM, Soares RM. Association between the presence of serum antibodies against *Neospora* spp. and fetal loss in equines. *Vet Parasitol* 2006; 142: 372-375.
  19. Vanleeuwen JA, Haddad JP, Dohoo IR, Keefe GP, Tiwari A, Scott HM. Risk factors associated with *Neospora caninum* seropositivity in randomly sampled Canadian dairy cows and herds. *Prev Vet Med* 2010; 93: 129-138.
  20. Dubey JP, Schares G. Neosporosis in animals-the last five years. *Vet Parasitol* 2011; 180: 90-108.
  21. Strohbusch M, Müller N, Hemphill A, Krebber R, Greif G, Gottstein B. Toltrazuril treatment of congenitally acquired *Neospora caninum* infection in newborn mice. *Parasitol Res* 2009; 104: 1335-1343.