

Effect of Colchicine on the Induction of *Prunella vulgaris* for. *albiflora* Nakai

Soo-Jeong Kwon*, Swapan Kumar Roy**, Kab-Yeon Cho*, Young-Ja Moon*, Sun-Hee Woo**, and Hag-Hyun Kim*[†]

*Department of Food Nutrition and Cookery, Woosong College, Daejeon 300-715, Korea

**Department of Crop Science, Chungbuk National University, Cheong-ju 361-763, Korea

ABSTRACT This study was conducted to find out the effective induction method of tetraploid plants to obtain potential data for cultivating superior varieties by colchicine treatment. The seed germination were decreased by the higher concentration of colchicine treatment and longer soaking time. A total of 907 individuals were germinated in 16 treated plots except control (untreated plot) and 28 tetraploids were induced which was about 3.1% of the number of seed germinated. The plant regeneration rate by colchicine treatment on explant of *Prunella vulgaris* for. *albiflora* Nakai under in vitro culture was decreased with the higher concentration of colchicine. While a total of 312 individuals were regenerated in all treatments, the explant was soaked in more than 0.05% for over 1 hour, tetraploid could be obtained. In particular, for the soaking treatment in 0.05% for 6 hours and 12 hours, 37 tetraploids were induced, which was about 57.8% of the number of plant regenerated. In accordance with the observation on doubling of DNA contents in leaf in order to identify polyploid, the peak DNA content of G1 phase was 101.3 for diploid and 197.2 for tetraploid. The result confirmed the doubling of DNA content. Furthermore, the number of chloroplasts per guard cell depending on polyploid was around 10 in diploid and 19.3 in tetraploid, which was around 1.9 times as much as diploid.

Keywords : tetraploid, colchicine, DNA content, number of chloroplasts, morphological characteristic

As one of allied species of *Prunella vulgaris* var. *lilacina*, '*Prunella vulgaris* for. *albiflora* Nakai' belonging to *Prunella vulgaris* var. *lilacina* family is a perennial plant which bear the white colored flower. Therefore, its scarcity is recognized compared to *Prunella vulgaris* var. *lilacina* that producing purple flowers. In Korea, it is mainly used for medicine and nectar source but supply is limited compared to demand and the method of cultivation is very primitive and there is little

full-fledged research for the improvement of varieties and cultivation methods.

The artificial induction of tetraploid plants is generally being made a lot by varying concentration, treatment time and treatment method etc. Colchicine is known to be the inhibition of the formation of spindle fibers by combining tubulin in the process of somatic cell division of plants and induces polyploidization of chromosome by interfering with the formation of microtubules and anodal movement of chromosome in the middle stage of cell division (Hadlaczky *et al.*, 1983).

In general, organs or stems of tetraploid plants become larger (Cockerham and Galletta, 1976; Lapins, 1975) and stems get thicker and longer and leaves and flowers get larger as well. In addition to polyploidization, particularly, the component content of secondary metabolite such as sugars in candy cane, vitamin C in fruit of tomatoes and apples, nicotine in tobacco leaves etc. are changed and characteristics such as virus-free resistance (Hahn, 1969), freezing resistance of mulberry (Park, 1994) etc. are improved in some cases.

Generally, methods to identify polyploidy were regarded as the observation of pollen grains (Bamberg and Hanneman, 1991), number of chloroplast (Chaudhari and Barrow, 1975; Dudley, 1958; Bae *et al.*, 2001), stomatal cells (Borrino and Powell, 1987), marker genes such as 'seed maker' or 'embryo maker' (Bingham, 1969; Verdenius, 1973; Nonda and Chase, 1966), characteristics of the plant (Gaines and Aase, 1926; Hougas and Peloquin, 1957) but the analysis method by the amount of intracellular DNA and chromosome observation known as the most accurate methods were mainly used (Galbraith *et al.*, 1983; Miyoshi and Asakura, 1996; Sari *et al.*, 1999). However, the disadvantages of the method by chromosome when usually used that some technical problems and efforts were accompanied and a large amount of materials cannot be

[†]Corresponding author: (Phone) +82-10-8919-3911 (E-mail) hkyushu@hanmail.net

<Received 23 November, 2014; Accepted 25 January, 2015>

examined. Currently, therefore, polyploidy is determined by simply measuring DNA content with flow cytometry that can analyze a large amount of materials accurately.

In order to obtain basic data for cultivating superior varieties of '*Prunella vulgaris* for. *albiflora* Nakai', this study were conducted to examine the appropriate materials and appropriate concentrations, soaking time, etc. and to find out the effective induction method of tetraploid plants by colchicine treatment.

MATERIALS AND METHODS

Tetraploid cultivation by colchicine treatment

Seeds of '*Prunella vulgaris* for. *albiflora* Nakai' were grown in 2013 and explants of cultured plants were grown after in vitro sterilization that were used as experimental materials. For treatment, 20 ml of 0.01, 0.05, 0.1 and 0.5% colchicine solution were applied in Petridish on which 50 seeds were soaked and left for 1, 3, 6 and 12 hours under the low temperature condition (5°C). Each treatment was repeated 3 times. After soaking, each seed was washed 3~4 times with sterile water and then, it was sown in bed soil for gardening and sprouted in the constant temperature room of 25°C. Germination was investigated when cotyledons emerged and leaves were collected when more than 6 foliage leaves emerged to check the presence or absence of polyploidy.

Also, in order to treat explants of cultured plants, 40 ml of 0.01, 0.05 and 0.1% colchicine solution was put in a beaker of 10 ml and then the nodes of '*Prunella vulgaris* for. *albiflora* Nakai' were cut to a length of about 1cm and soaked for 1, 6 and 12 hours. After soaking, each explant was washed with sterile water 3~4 times and then 5 explants were placed in MS medium and each treatment was repeated 9 times. Culture was made at 25±1°C and illuminated for 16 hours by luminosity of 40 μmol·m⁻²·s⁻¹.

DNA content analysis using cytometry

The plant leaf of each treatment plot was cut to a size of about 0.5 × 0.5 cm and then, HR-A solution (Patec Inc., Germany) was added and then the tissue was crushed to extract DNA. HR-B solution (Patec, Germany) was added to this solution for staining and then, doubling of DNA content was checked by using Flow cytometry (Patec PA-1, Germany) and polyploidy was determined through this result.

Investigation of the number of chloroplasts in guard cells

In order to observe chloroplasts, it was stained by separating the back side of leaves collected in the middle part of the plant to cut the back side epidermis and then, immersed it in iodine-potassium solution (1%(w/v) iodine, 2%(w/v) potassium iodide) for 2~3 hours and stained back side epidermis was examined through a microscope. To count the number of chloroplasts in guard cells, 30 cells were investigated in leaves of three objects per repetition and then the average was calculated.

RESULTS AND DISCUSSION

Seed soaking treatment of colchicine

Seed germination rate and polyploid induction rate were investigated by different concentration and soaking duration of colchicine in '*Prunella vulgaris* for. *albiflora* Nakai' seeds

Table 1. Effect of colchicine on chromosome doubling and germination in '*Prunella vulgaris* for. *albiflora* Nakai'.

Conc. (%)	Soaking time (hrs)	No. of seeds treated	No. of seeds germinated	% of germination	No. of tetraploids
Control		50	43.6	87.3a ^z	-
0.01	1	50	35	70.0ab	-
0.05		50	30.3	60.7bc	-
0.1		50	28	56.0bcd	-
0.5		50	20.3	40.7def	1
0.01	3	50	31.6	63.3bc	-
0.05		50	26.6	53.3bcd	-
0.1		50	20.6	41.3def	2
0.5		50	15.6	31.3efg	2
0.01	6	50	23.3	46.7cde	1
0.05		50	19.6	39.3def	7
0.1		50	11.6	23.3fgh	5
0.5		50	9.3	18.7gh	2
0.01	12	50	15	30.0efg	3
0.05		50	7.3	14.7gh	2
0.1		50	4.6	9.3h	2
0.5		50	3	6.0h	1

^zMean separation within columns by Duncan's multiple range test (p=0.05).




Fig. 1. Comparison of DNA contents in diploid and tetraploid of *Prunella vulgaris* for *albiflora* Nakai. Flow histograms showing DNA measurements of nuclei from leaves.

Table 2. Comparison of growth characteristics in diploid and tetraploid of *Prunella vulgaris* for *albiflora* Nakai.

Ploidy	Plant height (cm)	Leaf length (cm)	Leaf width (cm)	No. of leaves	Petiole length (cm)
Diploid	3.0±0.4b ^z	3.1±0.2a	2.2±0.2a	13.5±1.0b	2.7±0.1b
Tetraploid	4.3±0.6a	2.3±0.3b	2.2±0.3a	17.3±1.3a	2.7±0.5b

^zMean separation within columns by Duncan's multiple range test ($p=0.05$).

and the results were shown in Table 1.

The germination rate of seeds were tended to decrease when the higher the concentration of colchicine and the longer the soaking duration. In particular, in the case of 12-hour soaking treatment, the germination rate was found to be significantly suppressed to the range of 6~14.7 % in the concentration plot of more than 0.05%. Total 907 individuals showed 37.8% of germination rate that were germinated in 16 colchicine treated plots except for control (untreated plot). In order to check polyploidy, GAIN value of flow cytometry was fixed to 520.0 and then the DNA content of leaves was investigated. The result revealed that DNA content peak of G₁ phase was doubled to 101.3 in diploid (control plot), 197.2 in tetraploid. M₁ plants of tetraploid in which DNA content was observed doubled and as a result, 28 tetraploid plants with 3.1% content could be obtained (Fig. 1).

In order to increase the efficiency of tetraploid induction, seeds were treated with colchicine. Results showed that the induction efficiency of tetraploid plants could be observed higher in the longer duration of soaking time compare to

soaking concentration. Also, when colchicine was treated for 6 hours or 12 hours regardless of the high and low concentrations of colchicine, there were differences in the tetraploid induction efficiency but tetraploid plants were induced in all concentration plots. In particular, 5 and 7 tetraploid plants were obtained when treated for 6 hours at concentrations of 1% and 0.05, respectively, showing the highest induction of tetraploid plants.

From the above results, colchicine concentrations (ranges from 0.05~1%) and treatment duration 6 hours were considered to be suitable concentration and soaking time of colchicine seed treatment for tetraploid plant induction of *Prunella vulgaris* for. *albiflora* Nakai.

Prunella vulgaris for. *albiflora* Nakai identified as tetraploid was transplanted in pots filled with bed soil for gardening and cultivated for 2 months and then, the growth of the aerial part was investigated. The results are shown in Table 2 and Fig. 2. It was prevailed that the plant length of tetraploid plants was longer about 1.3 cm than that of diploid plants while the leaf length of diploid plants was longer. There were no differences




Fig. 2. Comparison of morphologic characteristics in diploid and tetraploid of *Prunella vulgaris* for *albiflora* Nakai.




Fig. 3. Comparison of number of chloroplasts per guard cell in diploid and tetraploid of *Prunella vulgaris* for *albiflora* Nakai.

in leaf width and petiole length between diploid and tetraploid plants while leaf number of tetraploid was found to be more by about 4 leaves (Table 2). Generally, in the case of tetraploid plants, stems are known to get thicker and leaves and flowers are also larger due to the expansion of organs or tissues (Cockerham & Galletta, 1976; Kim *et al.*, 2003) but *Prunella vulgaris* for. *albiflora* Nakai showed no significant difference except for significantly long plant length and leaf length. These results are considered to result from morphological differences in the early stage of growth.

Looking at the morphological characteristics of tetraploid

plants shown after colchicine treatment, several forms of tetraploid plants could be observed such as small or large plants compared to diploid plants. In particular, a lot of egg-shaped tetraploid objects could be seen compared to diploid with lanceolate leaves. In addition, diploid plants have the soft and thin leaf tissue and leaf color is also green while tetraploid plants showed the characteristics of thicker mesophyll tissue and dark green leaf color compared to diploid plants (Fig. 2).

The number of chloroplasts of guard cells of diploid and tetraploid of *Prunella vulgaris* for. *albiflora* Nakai is shown in Fig. 3. As for the number of chloroplasts per guard cell,

Table 3. Effect of colchicine on chromosome doubling and plant regeneration in *Hypericum patulum* Thunberg.

Conc. (%)	Soaking time (hrs)	No. of explants treated	No. of explants regenerated	% of regeneration	No. of tetraploids
0.01		45	44	97.8a ^z	0
0.05	1	45	42	93.3a	2
0.1		45	37	82.2ab	3
0.01		45	41	91.1a	6
0.05	6	45	36	80.0ab	18
0.1		45	28	62.2bc	11
0.01		45	39	86.7ab	5
0.05	12	45	28	62.2bc	19
0.1		45	17	37.8c	4

^zMean separation within columns by Duncan's multiple range test ($p=0.05$).




Fig. 4. Comparison of DNA contents in diploid and tetraploid of *Prunella vulgaris* for *albiflora* Nakai. Flow histograms showing DNA measurements of nuclei from leaves.

diploid plants have 10 chloroplasts and tetraploid plants 19.3 and tetraploid was found to have about two times more chloroplasts than diploid. These results were very similar to the results that the number of chloroplasts in guard cells according to polyploidy of potatoes significantly increases, 12.2 in haploid, 18.4 in triploid and 20.2 in tetraploid as polyploidy is increased (Cho *et al.*, 1994) and even for tobacco, the number of chloroplasts in guard cells also increases as polyploidy increases (Bae *et al.*, 2001).

In vitro cultured explant soaking treatment of colchicine

The effects of soaking treatment concentration and time duration of colchicine on polyploid induction of *Prunella*

vulgaris for. *albiflora* Nakai explants being cultured in vitro and plant regeneration are shown in Table 3 and Fig. 4.

405 explants of *Prunella vulgaris* for. *albiflora* Nakai were treated with colchicine and as a result, 312 individuals were regenerated at all concentrations, showing 77.0% of regeneration rate and the number of regenerated plants showed a tendency of lowering at the high concentration of colchicine and the longer period of time. In order to check polyploidy, the DNA content of regenerated 312 individuals was investigated after fixing GAIN value of flow cytometry to 490.0 and as a result, DNA content peak of G₁ phase was found to be 100.4 in diploid and 201.1 in tetraploid, showing that DNA content is doubled (Fig. 4). The DNA content of regenerated 312



Fig. 5. Comparison of morphologic characteristics in diploid and tetraploid of *Prunella vulgaris* for *albiflora* Nakai. cultured in vitro.

individuals was analyzed and as a result, tetraploid plants were obtained from a total of 68 objects, showing the high induction rate of 21.8%. Tetraploid plants were induced in all treatment plots except for colchicine 0.01% at 1 hour soaking treatment plot. However, in the case of plots treated for 6 and 12 hours at the concentration of 0.05%, and the induced 18, 19 tetraploid plants showed the highest results respectively. These results were very similar to the results that the tetraploid plants induction using in vitro culture of *Hypericum perforatum*, the highest induction rate was shown when immersed with colchicine 0.05% for 12 hours (Kwon *et al.*, 2013).

The above results showed that induction of tetraploid plants using in vitro culture was found to be about 7.0 times higher than the tetraploid induction rate of 3.1% obtained during seed treatment. So, it is more effective to treat colchicine in explants of in vitro culture showing vigorous cell division than directly treat it in seeds and may be used as a way of efficient induction of tetraploid plants.

The characteristics of tetraploid plants after colchicine treatment observed that leaf area gets wider and the form of the petiole gets thicker and shorter compared to diploid. It was observed that diploid carried the weak and thin leaf tissue and leaves were also light green while tetraploid plants showed the characteristics of thicker mesophyll tissue than diploid and leaf color of dark green (Fig. 5).

REFERENCES

- Bae, C. H., Y. H. Lee, D. C. Yang, K. S. Min, H. I. Kim, and H. Y. Lee. 2001. Changes of chloroplast number per guard cell pairs of leaves by ploidy level in *Nicotiana tabacum* L. cv. BY-4. Korean J. Plant Tissue Culture. 28(4) : 179-184.
- Bamberg, J. B. and R. E. Hanneman Jr. 1991. Rapid ploidy screening of tuberbearing *Solanum* (potato) species through pollen diameter measurement. Am. Potato J. 68 : 279-285.
- Bingham, E. T. 1969. Haploids from cultivated alfalfa *Medicago sativa* L. Nature 221 : 865-866.
- Borrino, E. M. and W. Powell. 1987. Stomatal guard cell length as an indicator of ploidy in microspore derived plants of barley. Genome 30 : 158-160.
- Chaudhari, H. K. and J. R. Barrow. 1975. Identification of cotton haploids by stomatal chloroplast-count technique. Crop Sci. 15 : 760-763.
- Cho, H. M., H. Y. Kim, and I. J. Mok. 1994. Stomatal cell characters as an indicator for haploid identification from progeny plants derived by 4xX2x interspecific crosses in potatoes. Korean J. Breed. 26(1) : 66-73.
- Cockerham, L. E. and G. J. Galletta. 1976. A survey of pollen characteristics in certain *Vaccinium* species. J. Amer. Soc. Hort. Sci. 101 : 671-676.
- Dudley, J. W. 1958. Number of chloroplasts in guard cells of inbred lines of tetraploid and diploid sugar beets. Agron. J. 50(3) : 169-170.
- Gaines, E. F. and H. C. Aase. 1926. A haploid wheat plant. Amer. J. Bot. 12 : 373-385.
- Galbraith, D. W., K. R. Harkins, J. M. Maddox, N. M. Ayres, D. P. Sharma, and E. Firoozabady. 1983. A rapid flow cytometric analysis of the cell cycle in intact plant tissues. Science 220 : 1049-1051.
- Hadlaczy, G., G. Bistray, T. Parznovszky, and D. Dudits. 1983. Mass isolation of plant chromosomes and nuclei. Planta. 157 : 278-385.
- Hahn, S. G. 1969. Studies on the Tetraploid Radish (*Raphanus Sativus* L.) Induced by Colchicine. I. On the Resistance to Virus Diseases. J. Kor. Soc. Hort. Sci. 5 : 48-56.
- Hougas, R. W. and S. J. Peloquin. 1957. A haploid plant of the potato variety Katahdin. Nature 180 : 1209-1210.
- Kim, I. H., H. H. Kim, E. Y. Hong, J. S. Yoon, J. K. Hwang, and

- C. H. Lee. 2003. Breeding of tetraploid in *Platycodon grandiflorum* (Jacq.) A. DC. by colchicine treatment. Korean J. Plant Res. 6 : 188-194.
- Kwon, S. J., K. Y. Cho, and H. H. Kim. 2013. A Tetraploid Induction in *Hypericum patulum* Thunberg by Colchicine Soaking Treatment. Korean J. Plant Res. 26 : 284-288.
- Lapins, K. O. 1975. Polyploidy and mutations induced in apricot by colchicine treatment Can. J. Genet. Cytol. 17 : 591-599.
- Miyoshi, K. and N. Asakura. 1996. Callus induction, regeneration of haploid plants and chromosome doubling in ovule cultures of pot gerbera (*Gevera jamesonii*). Plant Cell Rep. 16 : 1-5.
- Nonda, D. K. and S. S. Chase. 1966. An embryo marker for detecting monoploid of maize (*Zea mays* L.). Crop Sci. 6 : 213-215.
- Park, K. J. 1994 Cold-hardiness Tetraploid Induced by Colchicine Treatment in Mulberry Seedlings (*Morus alba* L. Yongchonppong/Kaeryangppong). Kor. J. Seric. Sci. 36 : 1-7.
- Sari, N., K. Abak, and M. Pitrat. 1999. Comparison of ploid level screening methods in watermelon: *Citrullus lanatus* (Thunb.) Matsum. and Nakai. Sci. Hort. 82 : 265-277.
- Verdenius, J. 1973. Selection from *Solanum tuberosum* group Phureja of genotypes combining high-frequency haploid induction with homozygosity for embryo-spot. Euphytica. 22 : 244-259.