



Infection of *Taenia asiatica* in a Bai Person in Dali, China

Li Wang^{1,2,*}, Xuenong Luo², Junling Hou², Aijiang Guo², Shaohua Zhang², Hailong Li³, Xuepeng Cai^{1,2,*}

¹College of Veterinary Medicine, Jilin University, Changchun, Jilin, P. R. China; ²State Key Laboratory of Veterinary Etiological Biology, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu, P. R. China; ³Department of Human Parasitology, School of Basic Medicine, Dali University, Dali, P. R. China

Abstract: We report here a human case of *Taenia asiatica* infection which was confirmed by genetic analyses in Dali, China. A patient was found to have symptoms of taeniasis with discharge of tapeworm proglottids. By sequencing of the mitochondrial cytochrome *c* oxidase subunit 1 (*cox1*) gene, we observed nucleotide sequence identity of 99% with *T. asiatica* and 96% with *T. saginata*. Using the cytochrome *b* (*cytb*) gene, 99% identity with *T. asiatica* and 96% identity with *T. saginata* were found. Our findings suggest that taeniasis of people in Dali, China may be mainly caused by *T. asiatica*.

Key words: *Taenia asiatica*, case report, *cox1* gene, *cytb* gene

INTRODUCTION

Human taeniasis is caused by 3 species of taeniid tapeworms, *Taenia solium*, *Taenia saginata*, and *Taenia asiatica*. Among these 3 species, *T. asiatica* was recorded as a distinct one from *T. saginata* in 1993 both of which are morphologically similar to each other [1-3]. *T. asiatica* is more restrictedly distributed in Asian countries, i.e., Korea, China, Taiwan, Thailand, Indonesia, Vietnam, Japan, and the Philippines [1].

Because of the morphological similarity, *T. asiatica* had previously been considered as *T. saginata* [2,3]. Thus, molecular methods became an efficient tool which can provide a clearer phylogenetic resolution. The methods of PCR-RFLP, single-strand conformation polymorphism (SSCP), a loop-mediated isothermal amplification (LAMP), multiplex PCR, and co-proDNA test have been reported [4-9]. These methods have their own advantages accordingly. Their mitochondrial cytochrome *c* oxidase 1 (*cox1*) and cytochrome *b* (*cytb*) gene sequences have been determined completely, and they have been widely used to study the population structure and genetic differentiation of several tapeworm species [10-12].

A previous study indicated that short sequences of the *cox1* gene can give a bias to analysis, and rather a complete se-

quence data would provide more reliable results [13,14]. Phylogenetic trees can show the relationships among different taxa. In the present study, we report a human case of *T. asiatica* infection by genetic analyses based on *cox1* and *cytb* genes in Dali, China.

CASE RECORD

The patient (Bai nationality, Dali, China) is a healthy 30-year-old man who visited Dali People's Hospital after finding a whitish yellow tapeworm segment in his feces on 26 May 2014. He had not experienced any abdominal discomfort or pain, and had not visited any foreign countries recently. He had a history of eating raw pork and raw pig liver mixed with sour sauce and salted garlic. His blood test results were normal.

The man was treated with traditional Chinese medicine (the combination of pumpkin seeds and areca nut extract) and 30% hydrated magnesium sulfate ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$) solution. An almost complete cestode (about 1.8 m long) without a scolex was expelled after about 3-6 hr of treatment. This tapeworm specimen was kept in a 50 ml centrifuge tube filled with 0.9% normal saline and forwarded to National Key Laboratory of Veterinary Etiological Biology (Lanzhou, China) for examinations.

Genomic DNA (gDNA) was extracted from the proglottid of the tapeworm using an AxyPrep small genomic DNA kit (Axygen, Beijing, China). Purified gDNA was used as a template for PCR for *cox1*. The PCR primers were up primer (5'-ATGAGTGT-TAAATTTTGTTAAGTT-3') and down primer (5'-TTAAACTA-AAAAACCACGAGC-3'). PCR was performed in a 50 μl reac-

•Received 15 October 2015, revised 5 December 2015, accepted 3 January 2016.

*Corresponding authors (sdhywangli@163.com; caixp@vip.163.com)

© 2016, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

tion mixture containing 25 µl of PrimeSTAR® HS Premix (Takara, Kyoto, Japan), 1.5 µl of 10 µM of each primers, 20 µl of distilled water, and 2 µl purified gDNA from tapeworm. The PCR reaction was carried out for 35 cycles of 94°C for 40 sec, 55°C

for 40 sec, 72°C for 60 sec. The reaction initial denaturing step at 95°C for 5 min and terminated with a final extension step at 72°C for 10 min. The PCR amplification of *cytb* sequence was performed using a primer pair of *cytb*: up primer (5'-ATGATTA-GATTAT TTCGACG-3') and down primer (5'-TTAATAAATCTTA-AAAAGAAACATAAGC-3'). PCR conditions for *cytb* gene were the same as those for *cox1*.

The amplification products were fractionated using 1% (w/v) agarose gel, and then purified (Axygen). The purified products were then ligated into PMD-18T vector (Takara), followed by transformation into *Escherichia coli* DH5a and sequencing. The sequences for *cox1* and *cytb* were aligned with the corresponding sequences of tapeworms obtained from GenBank database. Molecular identification of the tapeworm specimens was based on a comparison with the nucleotide sequences of *cox1* and *cytb* genes of *T. solium*, *T. Saginata*, and *T. asiatica*. Phylogenetic analyses were performed using the neighbor-joining (NJ) method by MEGA (Version 5) with the Kimura-2 parameter model [15]. Bootstrap analysis was performed with 1,000 replications.

The whitish yellow tapeworm from the patient was shown in Fig. 1. Approximately, 1,620 bp nucleotide sequence from the mitochondrial *cox1* gene of the tapeworm in the present




Fig. 1. An almost complete strobila (about 1.8 m in length), without the scolex, of *Taenia asiatica* recovered from our patient.




Fig. 2. Phylogenetic tree of 3 *Taenia* spp. using the mitochondrial *cox1* gene sequences. Our tapeworm specimen showed 99% identity (1 different base) with *T. asiatica* (AB533175.1) and 96% identity (65 different bases) with *T. saginata* (AB465239.1).




Fig. 3. Phylogenetic tree of 3 *Taenia* spp. using the mitochondrial *cytb* gene sequences. Our tapeworm specimen showed 99% identity (2 bp differences) with *T. asiatica* (no. AB066580.1) and 96% identity (43 bp different bases) with *T. saginata*.

study was obtained, which showed an identity of 99% (1 different base) with *T. asiatica* (AB533175.1) and 96% (65 different bases) with *T. saginata* (AB465239.1), respectively. The phylogenetic analysis revealed that our tapeworm was closely related to *T. asiatica* (Fig. 2). Furthermore, approximately 1,068 bp sequence of the mitochondrial *cytb* gene had an identity of 99% (2 bp differences) with *T. asiatica* (no. AB066580.1) and 96% (43 bp different bases) with *T. saginata*, respectively. Phylogenetic analysis showed that the cestode was closely related to *T. asiatica* (Fig. 3). Sequences of all *Taenia* specimens derived from different Asian countries formed a monophyletic group.

DISCUSSION

To date, taeniasis still occurs in the majority of regions in Southwest China including Yunnan, Sichuan, Guizhou, Qinghai, and many other provinces where a minority of people like to eat raw pork, undercooked beef, and raw pig liver mixed with sour sauce and salted garlic, and the prevalence and incidence of human taeniasis remains unknown in southwest China [1,16-18]. Eating raw liver which contains cysticerci is the main risk factor of tapeworm infection. In the present study, since infection of the patient was a result of consumption of undercooked pig liver, it is reasonable to guess that the taeniasis is due to *T. asiatica* caused in Dali of Yunnan Province. Neither *T. saginata* nor *T. solium* were detected during the same period, so *T. asiatica* was considered to be the dominant species causing human taeniasis in Dali.

As the living standards of people improve, the incidence of taeniasis became markedly lower in Dali, but the local Bai people still have the habit of eating raw pork and raw pig liver. So, it is necessary for them to know the dangers of taeniasis and to understand the routes of tapeworm transmission through continuous public health educations. There are 2 strategies for cutting off the domestic life of *T. asiatica*. First, local people should pay attention to prevent human taeniasis avoiding intake of uncooked or undercooked pig liver. Second, the patient who is infected with the tapeworm must be treated early with anthelmintics and keep pigs without contacting human feces. We believe that the taeniasis will be eliminated someday in the future if native people take above these 2 advices in Dali.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (no. 31372433), the Science Fund for Creative Research Groups of Gansu Province, China (no. 1210RJIA006) and the opening projects of National Key Laboratory of Veterinary Etiological Biology at Lanzhou Veterinary Research Institute of Chinese Academy of Agricultural Sciences, China (grant no. 201001).

CONFLICT OF INTEREST

We have no conflict of interest related to this study.

REFERENCES

1. Eom KS, Jeon HK, Rim HJ. Geographical distribution of *Taenia asiatica* and related species. *Korean J Parasitol* 2009; 47: 115-124.
2. Galan-Puchades MT, Fuentes MV. Diagnosis of human cysticercosis and *Taenia asiatica*. *Am J Trop Med Hyg* 2009; 81: 1165.
3. Eom KS, Rim HJ. Morphologic descriptions of *Taenia asiatica* sp. n. *Korean J Parasitol* 1993; 31: 1-6.
4. Bowles J, McManus DP. Genetic characterization of the Asian *Taenia*, a newly described taeniid cestode of humans. *Am J Trop Med Hyg* 1994; 50: 33-44.
5. Gass RB, Chilton NB. Characterisation of taeniid cestode species by PCR-RFLP of ITS2 ribosomal DNA. *Acta Trop* 1995; 59: 31-40.
6. Gass RB, Zhu X, Woods W. Genotyping *Taenia* tapeworms by single-strand conformation polymorphism of mitochondrial DNA. *Electrophoresis* 1999; 20: 2834-2847.
7. Gonzalez LM, Montero E, Harrison LJ, Parkhouse RM, Garate T. Differential diagnosis of *Taenia saginata* and *Taenia solium* infection by PCR. *J Clin Microbiol* 2000; 38: 737-744.
8. Nkouawa A, Sako Y, Nakao M, Nakaya K, Ito A. Loop-mediated isothermal amplification method for differentiation and rapid detection of *Taenia* species. *J Clin Microbiol* 2009; 47: 168-174.
9. Yamasaki H, Allan JC, Sato MO, Nakao M, Sako Y, Nakaya K, Qiu D, Mamuti W, Craig PS, Ito A. DNA differential diagnosis of taeniasis and cysticercosis by multiplex PCR. *J Clin Microbiol* 2004; 42: 548-553.
10. Gasser RB, Zhu X, Woods W. Genotyping *Taenia* tapeworms by single-strand conformation polymorphism of mitochondrial DNA. *Electrophoresis* 1999; 20: 2834-2837.
11. Nakao M, Okamoto M, Sako Y, Yamasaki H, Nakaya K, Ito A. A phylogenetic hypothesis for the distribution of two genotypes of the pig tapeworm *Taenia solium* worldwide. *Parasitology* 2002; 124: 657-662.
12. Martinez-Hernandez E, Jimenez-Gonzalez DE, Chenillo P, Alonso-Fernandez C, Maravilla P, Flisser A. Geographical widespread of two lineages of *Taenia solium* due to human migrations: can population genetic analysis strengthen this hypothesis? *Infect Genet Evol* 2009; 9: 1108-1114.
13. Jeon HK, Lee KH, Kim KH, Hwang UW, Eom KS. Complete sequence and structure of the mitochondrial genome of the human tapeworm, *Taenia asiatica* (Platyhelminthes; Cestoda). *Parasitology* 2005; 130: 717-726.
14. Jeon HK, Eom KS. *Taenia asiatica* and *Taenia saginata*: genetic divergence estimated from their mitochondrial genomes. *Exp Parasitol* 2006; 113: 58-61.
15. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011; 28: 2731-2739.
16. Zhang L, Tao H, Zhang B, Wang H, Wang Y, Li Z, Yang J, Yang B, Li Y, Pang Y, Zhang H, Wu Y. First discovery of *Taenia saginata asiatica* infection in Yunnan province. *Chin J Parasitol Parasit Dis* 1999; 17: 95-96 (in Chinese).
17. Wang ZR, Bao HE. Identification of *Taenia saginata* by mtCO I in four areas of Yunnan and Guizhou provinces. *Chin J Parasitol Parasit Dis* 2003; 21: 20-23 (in Chinese).
18. Li T, Craig PS, Ito A, Chen X, Qiu D, Qiu J, Sato MO, Wandra T, Bradshaw H, Li L, Yang Y, Wang Q. Taeniasis/cysticercosis in a Tibetan population in Sichuan Province, China. *Acta Trop* 2006; 100: 223-231.