

Structures of Pseudo Ideal and Pseudo Atom in a Pseudo Q -Algebra

YOUNG BAE JUN

Department of Mathematics Education, Gyeongsang National University, Jinju 52828, Korea

e-mail: skywine@gmail.com

HEE SIK KIM

Department of Mathematics, Hanyang University, Seoul 04763, Korea

e-mail: heekim@hanyang.ac.kr

SUN SHIN AHN*

Department of Mathematics Education, Dongguk University, Seoul 04620, Korea

e-mail: sunshine@dongguk.edu

ABSTRACT. As a generalization of Q -algebra, the notion of pseudo Q -algebra is introduced, and some of their properties are investigated. The notions of pseudo subalgebra, pseudo ideal, and pseudo atom in a pseudo Q -algebra are introduced. Characterizations of their properties are provided.

1. Introduction

Y. Imai and K. Iséki introduced two classes of abstract algebras: a BCK -algebra and a BCI -algebra ([7,8]). It is known that the class of BCK -algebras is a proper subclass of the class of BCI -algebras. Q. P. Hu and X. Li [5,6] introduced a wide class of abstract algebra: a BCH -algebra. They have shown that the class of BCI -algebra is a proper subclass of the class of BCH -algebra. BCK -algebras have several connections with other areas of investigation, such as: lattice ordered groups, MV -algebras, Wajsberg algebras, and implicative commutative semigroups. J. M. Font et al. [3] have discussed Wajsberg algebras which are term-equivalent to MV -algebras. D. Mundici [13] proved MV -algebras are categorically equivalent to bounded commutative BCK -algebra, and J. Meng [11] proved that implicative com-

* Corresponding Author.

Received December 19, 2014; accepted February 5, 2016.

2010 Mathematics Subject Classification: 06F35, 03G25.

Key words and phrases: pseudo atom, pseudo subalgebra, pseudo ideal, pseudo Q -algebra.

mutative semigroups are equivalent to a class of *BCK*-algebras. G. Georgescu and A. Iorgulescu [4] introduced the notion of a pseudo *BCK*-algebra. Y. B. Jun characterized pseudo *BCK*-algebras. He found conditions for a pseudo *BCK*-algebra to be \wedge -semi-lattice ordered. Y. B. Jun, H.S. Kim, J. Neggers [9] introduced the notion of a pseudo *d*-algebra as a generalization of the idea of a *d*-algebra. J. Neggers, S. S. Ahn and H. S. Kim ([14]) introduced a new notion, called a *Q*-algebra, which is a generalization of *BCH/BCI/BCK*-algebra, and generalized some theorems discussed in a *BCI*-algebra.

In this paper, we introduce the notion of pseudo *Q*-algebra as a generalization of *Q*-algebra and investigate some of their properties. We also define the notions of pseudo subalgebra, pseudo ideal, and pseudo atom in a pseudo *Q*-algebra and provide characterizations of their properties in a pseudo *Q*-algebra.

2. Preliminaries

A *Q*-algebra ([14]) is a non-empty set X with a constant 0 and a binary operation “ $*$ ” satisfying axioms:

- (I) $x * x = 0$,
- (II) $x * 0 = x$,
- (III) $(x * y) * z = (x * z) * y$

for all $x, y, z \in X$.

For brevity we also call X a *Q*-algebra. In X we can define a binary relation “ \leq ” by $x \leq y$ if and only if $x * y = 0$.

In a *Q*-algebra X the following property holds:

- (IV) $(x * (x * y)) * y = 0$, for any $x, y \in X$.

A *BCK*-algebra is a *Q*-algebra X satisfying the additional axioms:

- (V) $((x * y) * (x * z)) * (z * y) = 0$,
- (VI) $x * y = 0$ and $y * x = 0$ imply $x = y$,
- (VII) $0 * x = 0$,

for all $x, y, z \in X$.

Definition 2.1.([14]) Let $(X; *, 0)$ be a *Q*-algebra and $\emptyset \neq I \subset X$. I is called a *subalgebra* of X if

- (S) $x * y \in I$ whenever $x \in I$ and $y \in I$.

I is called an *ideal* of X if it satisfies:

- (Q_0) $0 \in I$,

(Q_1) $x * y \in I$ and $y \in I$ imply $x \in I$.

A Q -algebra X is called a QS -algebra ([1]) if it satisfies the following identity:

$$(x * y) * (x * z) = z * y, \quad \text{for any } x, y, z \in X.$$

Example 2.2.([1]) Let \mathbb{Z} be the set of all integers and let $n\mathbb{Z} := \{nz \mid z \in \mathbb{Z}\}$, where $n \in \mathbb{Z}$. Then $(\mathbb{Z}; -, 0)$ and $(n\mathbb{Z}; -, 0)$ are both Q -algebras and QS -algebras, where “ $-$ ” is the usual subtraction of integers. Also, $(\mathbb{R}; -, 0)$ and $(\mathbb{C}; -, 0)$ are Q -algebras and QS -algebras where \mathbb{R} is the set of all real numbers, \mathbb{C} is the set of all complex numbers.

Example 2.3. (1) Let $X = \{0, 1, 2\}$ be a set with the table as follows:

$*$	0	1	2
0	0	0	0
1	1	0	0
2	2	0	0

Then X is a Q -algebra, but not a QS/BCI -algebra, since $(2 * 0) * (2 * 1) = 2 \neq 1 = 1 * 0$.

(2) Let $X = \{0, 1, 2\}$ be a set with the table as follows:

$*$	0	1	2
0	0	2	1
1	1	0	2
2	2	1	0

Then X is both a Q -algebra and QS -algebra.

(3) Let $X = \{0, 1, 2\}$ be a set with the table as follows:

$*$	0	1	2
0	0	0	0
1	1	0	0
2	2	1	0

Then X is both a Q -algebra and BCI -algebra, but not a QS -algebra, since $(0 * 1) * (0 * 2) = 0 \neq 1 = 2 * 1$.

3. Pseudo Ideal

In the following, let X denote a pseudo Q -algebra unless otherwise specified.

Definition 3.1. A *pseudo Q -algebra* is a non-empty set X with a constant 0 and two binary operations “ $*$ ” and “ \diamond ” satisfying the following axioms: for any $x, y, z \in X$,

(P1) $x * x = x \diamond x = 0$;

(P2) $x * 0 = x = x \diamond 0$;

$$(P3) \quad (x * y) \diamond z = (x \diamond z) * y.$$

For brevity, we also call X a pseudo *BCH*-algebra. In X we can define a binary operation “ \preceq ” by $x \preceq y$ if and only if $x * y = 0$ if and only if $x \diamond y = 0$. Note that if $(X; *, 0)$ is a Q -algebra, then letting $x \diamond y := x * y$, produces a pseudo Q -algebra $(X; *, \diamond, 0)$. Hence every Q -algebra is a pseudo Q -algebra in a natural way.

Definition 3.2. Let $(X; *, \diamond, 0)$ be a pseudo Q -algebra and let $\emptyset \neq I \subseteq X$. I is called a *pseudo subalgebra* of X if $x * y, x \diamond y \in I$ whenever $x, y \in I$. I is called a *pseudo ideal* of X if it satisfies

$$(PI1) \quad 0 \in I;$$

$$(PI2) \quad x * y, x \diamond y \in I \text{ and } y \in I \text{ imply } x \in I \text{ for all } x, y \in X.$$

Example 3.3. Let $X := \{0, a, b, c\}$ be a set with the following Cayley tables:

$*$	0	a	b	c	\diamond	0	a	b	c
0	0	0	0	0	0	0	0	0	0
a	a	0	0	0	a	a	0	0	0
b	b	b	0	a	b	b	c	0	c
c	c	c	0	0	c	c	c	0	0

Then $(X; *, 0)$ and $(X; \diamond, 0)$ are not Q -algebras, since $(b * a) * c = a \neq 0 = (b * c) * a$ and $(b \diamond a) \diamond c = 0 \neq c = (b \diamond c) \diamond a$. It is easy to check that $(X; *, \diamond, 0)$ is a pseudo Q -algebra. Let $I := \{0, a\}$. Then I is both a pseudo subalgebra of X and a pseudo ideal of X . Let $J := \{0, a, c\}$. Then J is a pseudo subalgebra of X , but it is not a pseudo ideal of X since $b \diamond c = c \in J$ and $b * c = a \in J$, but $b \notin J$.

Proposition 3.4. Let I be a pseudo ideal of a pseudo Q -algebra X . If $x \in I$ and $y \preceq x$, then $y \in I$.

Proof. Assume that $x \in I$ and $y \preceq x$. Then $y * x = 0$ and $y \diamond x = 0$. By (PI1) and (PI2), we have $y \in I$. \square

Proposition 3.5. If X is a pseudo Q -algebra satisfying $a * b = a * c$ and $a \diamond b = a \diamond c$ for all $a, b, c \in X$, then $0 * b = 0 * c$ and $0 \diamond b = 0 \diamond c$.

Proof. For any $a, b, c \in X$, we have

$$0 * b = (a \diamond a) * b = (a * b) \diamond a = (a * c) \diamond a = (a \diamond a) * c = 0 * c$$

and

$$0 \diamond b = (a * a) \diamond b = (a \diamond b) * a = (a \diamond c) * a = (a * a) \diamond c = 0 \diamond c.$$

This concludes the proof. \square

Proposition 3.6. Let $(X; *, \diamond, 0)$ be a pseudo Q -algebra. Then the following hold: for all $x, y, z \in X$.

$$(i) \quad x * (x \diamond y) \preceq y, x \diamond (x * y) \preceq y,$$

- (ii) $x * y \preceq z \Leftrightarrow x \diamond z \preceq y$,
- (iii) $0 * (x * y) = (0 \diamond x) \diamond (0 * y)$,
- (iv) $0 \diamond (x \diamond y) = (0 * x) * (0 \diamond y)$,
- (v) $0 * x = 0 \diamond x$.

Proof. (i) By (P1) and (P3), we obtain $[x * (x \diamond y)] \diamond y = (x \diamond y) * (x \diamond y) = 0$ and $[x \diamond (x * y)] * y = (x * y) \diamond (x * y) = 0$. Hence $x * (x \diamond y) \preceq y$ and $x * (x \diamond y) \preceq y$.

(ii) $x * y \preceq z \Leftrightarrow (x * y) \diamond z = 0 \Leftrightarrow (x \diamond z) * y = 0 \Leftrightarrow x \diamond z \preceq y$.

(iii) and (iv) For any $x, y \in X$, by (P1) and (P3) we have

$$\begin{aligned}
 (0 \diamond x) \diamond (0 * y) &= [((x * y) * (x * y)) \diamond x] \diamond (0 * y) \\
 &= [((x * y) \diamond x) * (x * y)] \diamond (0 * y) \\
 &= [((x \diamond x) * y) * (x * y)] \diamond (0 * y) \\
 &= [(0 * y) \diamond (0 * y)] * (x * y) \\
 &= 0 * (x * y)
 \end{aligned}$$

and

$$\begin{aligned}
 (0 * x) * (0 \diamond y) &= [((x \diamond y) * (x \diamond y)) * x] * (0 \diamond y) \\
 &= [((x \diamond y) * x) * (x \diamond y)] * (0 \diamond y) \\
 &= [((x * x) \diamond y) \diamond (x \diamond y)] * (0 \diamond y) \\
 &= [(0 \diamond y) * (0 \diamond y)] \diamond (x \diamond y) \\
 &= 0 \diamond (x \diamond y).
 \end{aligned}$$

(v) For any $x \in X$, by (P1) and (P3) we obtain $0 * x = (x \diamond x) * x = (x * x) \diamond x = 0 \diamond x$. \square

Theorem 3.7. For any pseudo Q -algebra X , the set

$$K(X) := \{x \in X \mid 0 \preceq x\}$$

is a pseudo subalgebra of X .

Proof. Let $x, y \in K(X)$. Then $0 \preceq x$ and $0 \preceq y$. Hence $0 * x = 0 \diamond x = 0$ and $0 * y = 0 \diamond y = 0$. Since $0 * (x * y) = (0 \diamond x) \diamond (0 * y) = 0 \diamond 0 = 0$ and $0 \diamond (x \diamond y) = (0 * x) * (0 \diamond y) = 0 * 0 = 0$, we have $x * y, x \diamond y \in K(X)$. Thus $K(X)$ is a pseudo subalgebra of X . \square

Theorem 3.8. If I is a pseudo ideal of a pseudo Q -algebra X , then

- (i) $\forall x, y, z \in X, x, y \in I, z * y \preceq x \Rightarrow z \in I$,
- (ii) $\forall a, b, c \in X, a, b \in I, c \diamond b \preceq a \Rightarrow c \in I$.

Proof. (i) Suppose that I is a pseudo ideal of X and let $x, y, z \in X$ be such that $x, y \in I$ and $z * y \preceq x$. Then $(z * y) \diamond x = 0 \in I$. Since $x \in I$ and I is a pseudo ideal

of X , we have $z * y \in I$. Since $y \in I$ and I is a pseudo ideal of X , we obtain $z \in I$. Thus (i) is valid.

(ii) Let $a, b, c \in X$ be such that $a, b \in I$ and $c \diamond b \preceq a$. Then $(c \diamond b) * a = 0 \in I$ and so $c \diamond b \in I$. Since $b \in I$ and I is a pseudo ideal of X , we have $c \in I$. Thus (ii) is true. \square

Theorem 3.9. *Let I be a pseudo subalgebra of a pseudo Q -algebra X . Then I is a pseudo ideal of X if and only if $\forall x, y \in X, x \in I, y \in X - I \Rightarrow y * x \in X - I$ and $y \diamond x \in X - I$.*

Proof. Assume that I is a pseudo ideal of X and let $x, y \in X$ be such that $x \in I$ and $y \in X - I$. If $y * x \notin X - I$, then $y * x \in I$. Since I is a pseudo ideal of X , we have $y \in I$. This is a contradiction. Hence $y * x \in X - I$. Now if $y \diamond x \notin X - I$, then $y \diamond x \in I$ and so $y \in I$. This is a contradiction, and therefore $y \diamond x \in X - I$.

Conversely, assume that $\forall x, y \in X, x \in I, y \in X - I \Rightarrow y * x \in X - I$ and $y \diamond x \in X - I$. Since I is a pseudo subalgebra, we have $0 \in I$. Let $x \in I, y \in X$ such that $y * x, y \diamond x \in I$. If $y \notin I$, then $y * x, y \diamond x \in X - I$ by assumption. This is a contradiction. Hence $y \in I$. Thus I is a pseudo ideal of X . \square

Proposition 3.10. *Let A be a pseudo ideal of a pseudo Q -algebra X . If B is a pseudo ideal of A , then it is a pseudo ideal of X .*

Proof. Since B is a pseudo ideal of A , we have $0 \in B$. Let $y, x * y, x \diamond y \in B$ for some $x \in X$. If $x \in A$, then $x \in B$ since B is a pseudo ideal of A . If $x \in X - A$, then $y, x * y, x \diamond y \in B \subseteq A$ and so $x \in A$ because A is a pseudo ideal of X . Thus $x \in B$ since B is a pseudo ideal of A . This completes the proof. \square

Proposition 3.11. *Let I be a pseudo ideal of a pseudo Q -algebra X . Then*

$$(\forall x \in X)(x \in I \Rightarrow 0 * (0 \diamond x) \in I \text{ and } 0 \diamond (0 * x) \in I).$$

Proof. Let $x \in I$. Then

$$0 = (0 \diamond x) * (0 \diamond x) = (0 * (0 \diamond x)) \diamond x$$

and

$$0 = (0 * x) \diamond (0 * x) = (0 \diamond (0 * x)) * x$$

which imply that $0 * (0 \diamond x), 0 \diamond (0 * x) \in I$. This completes the proof. \square

Theorem 3.12. *Let I be a pseudo ideal of a pseudo Q -algebra X and let*

$$I^\# := \{x \in X \mid 0 * (0 \diamond x), 0 \diamond (0 * x) \in I\}.$$

Then $I^\#$ is a pseudo ideal of X and $I \subseteq I^\#$.

Proof. Obviously, $0 \in I^\#$. Let $a \in X, y \in I^\#$ such that $a * y, a \diamond y \in I^\#$. Then

$0 * (0 \diamond (a * y)), 0 \diamond (0 * (a * y)), 0 * (0 \diamond (a \diamond y)) \in I$, and $0 \diamond (0 * (a \diamond y)) \in I$. Using Proposition 3.6 (iii) and (iv), we have

$$\begin{aligned} (0 * (0 \diamond a)) * (0 \diamond (0 * y)) &= 0 \diamond ((0 \diamond a) \diamond (0 * y)) \\ &= 0 \diamond (0 * (a * y)) \in I \end{aligned}$$

and

$$\begin{aligned} (0 \diamond (0 * a)) \diamond (0 * (0 \diamond y)) &= 0 * ((0 * a) * (0 \diamond y)) \\ &= 0 * (0 \diamond (a \diamond y)) \in I. \end{aligned}$$

Since $0 * (0 \diamond y), 0 \diamond (0 * y) \in I$, it follows from (PI2) that $0 * (0 \diamond a), 0 \diamond (0 * a) \in I$. Hence $a \in I^\#$. Thus $I^\#$ is a pseudo ideal of X . By Proposition 3.11, we know that $I \subseteq I^\#$. This completes the proof. \square

Let X be a pseudo Q -algebra. For any non-empty subset S of X , we define

$$G(S) := \{x \in S \mid 0 * x = x = 0 \diamond x\}.$$

In particular, if $S = X$ then we say that $G(X)$ is the G -part of X .

Proposition 3.13. *If X is a pseudo Q -algebra, then a left cancellation law holds in $G(X)$.*

Proof. Let $x, y, z \in G(X)$ satisfy $x * y = x * z$ and $x \diamond y = x \diamond z$. By Proposition 3.5, we have $0 * y = 0 * z$ and $0 \diamond y = 0 \diamond z$. Since $y, z \in G(X)$, it follows that $y = z$. \square

Proposition 3.14. *Let X be a pseudo Q -algebra. Then $x \in G(X)$ if and only if $0 * x \in G(X)$ and $0 \diamond x \in G(X)$.*

Proof. If $x \in G(X)$, then $0 * x = x = 0 \diamond x$ and $0 * (0 * x) = 0 * x$ and $0 \diamond (0 \diamond x) = 0 \diamond x$. Hence $0 * x \in G(X)$ and $0 \diamond x \in G(X)$. Conversely, assume that $0 * x \in G(X)$ and $0 \diamond x \in G(X)$. Then $0 * (0 * x) = 0 * x$ and $0 \diamond (0 \diamond x) = 0 \diamond x$. By applying Proposition 3.13, we obtain $0 * x = x = 0 \diamond x$. Therefore $x \in G(X)$. \square

Proposition 3.15. *Let X be a pseudo Q -algebra. If $x, y \in G(X)$, then $x * y = y \diamond x$.*

Proof. If $x, y \in G(X)$, then $0 * x = x = 0 \diamond x$ and $0 * y = y = 0 \diamond y$. Using (P3), we have

$$x * y = (0 \diamond x) * (0 \diamond y) = (0 * (0 \diamond y)) \diamond x = (0 * y) \diamond x = y \diamond x.$$

This completes the proof. \square

Theorem 3.16. *If a pseudo Q -algebra X satisfies $x * (x \diamond y) = x * y$ or $x \diamond (x * y) = x \diamond y$ for all $x, y \in X$, then it is a trivial algebra.*

Proof. Assume that X satisfies $x * (x \diamond y) = x * y$ for all $x, y \in X$. Putting $x = y$ in the equation $x * (x \diamond y) = x * y$, we obtain

$$x = x * 0 = x * (x \diamond x) = x * x = 0.$$

Concerning the other case, the argument is similar. Hence X is a trivial algebra. \square

Theorem 3.17. *Let X be a pseudo Q -algebra such that*

$$(3.1) \quad x * (y \diamond z) = (x * y) \diamond z \text{ (resp., } x \diamond (y * z) = (x \diamond y) * z)$$

for all $x, y, z \in X$. Then X is a group under operation \diamond (resp. $*$).

Proof. Putting $x = y = z$ in (3.1) and using (P1) and (P2), we obtain $x = x * 0 = 0 \diamond x$ (resp., $x = x \diamond 0 = 0 * x$). This means that 0 is the zero element of X with respect to the operation \diamond (resp., $*$). Since $x \diamond x = 0 = x * x$, we know that the inverse of x is itself with respect to the operation \diamond (resp., $*$). Hence X is a group under operation \diamond (resp., $*$). \square

Definition 3.18. A Pseudo Q -algebra X is said to be \diamond -*medial* if it satisfies the following identity

$$(M1) \quad (x * y) \diamond (z * u) = (x * z) \diamond (y * u), \forall x, y, z, u \in X.$$

Proposition 3.19. *Every \diamond -medial pseudo Q -algebra X satisfies the following identities: for any $x, y \in X$*

$$(i) \quad x * y = 0 \diamond (y * x),$$

$$(ii) \quad 0 \diamond (0 * x) = x,$$

$$(iii) \quad x \diamond (x * y) = y.$$

Proof. (i) For any $x, y \in X$, we have $x * y = (x * y) \diamond 0 = (x * y) \diamond (x * x) = (x * x) \diamond (y * x) = 0 \diamond (y * x)$.

(ii) If we put $y = 0$ in (i), then we have (ii).

(iii) Using (ii), (P1) and (P2), we have $x \diamond (x * y) = (x * 0) \diamond (x * y) = (x * x) \diamond (0 * y) = 0 \diamond (0 * y) = y$. \square

4. Pseudo Atom

Definition 4.1. An element a of a pseudo Q -algebra X is called a *pseudo atom* of X if for every $x \in X$, $x \preceq a$ implies $x = a$.

Obviously, 0 is a pseudo atom of X . Let $L(X)$ denote the set of all pseudo atoms of X , we call it the *center* of X .

Theorem 4.2. *Let X be a pseudo Q -algebra. Then the following are equivalent: for all $x, y, z, w, u \in X$*

$$(i) \quad w \text{ is a pseudo atom,}$$

$$(ii) \quad w = x \diamond (x * w) \text{ and } w = x * (x \diamond w)$$

$$(iii) \quad (x * y) \diamond (x * w) = w * y \text{ and } (x \diamond y) * (x \diamond w) = w \diamond y$$

$$(iv) \quad w * (x \diamond y) = y \diamond (x * w) \text{ and } w \diamond (x * y) = y * (x \diamond w),$$

$$(v) \ 0 \diamond (y * w) = w * y \text{ and } 0 * (y \diamond w) = w \diamond y,$$

$$(vi) \ 0 \diamond (0 * w) = w \text{ and } 0 * (0 \diamond w) = w,$$

$$(vii) \ 0 \diamond (0 * (w \diamond z)) = w \diamond z \text{ and } 0 * (0 \diamond (w * z)) = w * z,$$

$$(viii) \ z \diamond (z * (w \diamond u)) = w \diamond u \text{ and } z * (z \diamond (w * u)) = w * u.$$

Proof. (i) \Rightarrow (ii): Let w be a pseudo atom of X . Since $x \diamond (x * w) \preceq w$ and $x * (x \diamond w) \preceq w$ by Proposition 3.6 (i), we have $w = x \diamond (x * w)$ and $w = x * (x \diamond w)$.

(ii) \Rightarrow (iii): For every $x \in X$, we obtain $(x * y) \diamond (x * w) = (x \diamond (x * w)) * y = w * y$ and $(x \diamond y) * (x \diamond w) = (x * (x \diamond w)) \diamond y = w \diamond y$ by (P3) and (ii).

(iii) \Rightarrow (iv): Replacing y by $x \diamond y$ in (iii), we get

$$\begin{aligned} w * (x \diamond y) &= (x * (x \diamond y)) \diamond (x * w) \\ &= (x \diamond (x * w)) * (x \diamond y) \\ &= y \diamond (x * w) \end{aligned}$$

and

$$\begin{aligned} w \diamond (x * y) &= (x \diamond (x * y)) * (x \diamond w) \\ &= (x * (x \diamond w)) \diamond (x * y) \\ &= y * (x \diamond w). \end{aligned}$$

(iv) \Rightarrow (v): Put $y := 0$ in (iv). Then $w * (x \diamond 0) = 0 \diamond (x * w)$ and $w \diamond (x * 0) = 0 * (x \diamond w)$. Hence $w * x = 0 \diamond (x * w)$ and $w \diamond x = 0 * (x \diamond w)$ by (P2).

(v) \Rightarrow (vi): Set $y := 0$ in (v). Then $0 \diamond (0 * w) = w * 0 = w$ and $0 * (0 \diamond w) = w \diamond 0 = w$.

(vi) \Rightarrow (vii): For any $w, z \in X$, we have

$$\begin{aligned} 0 \diamond (0 * (w \diamond z)) &= 0 * (0 * (w \diamond z)) \\ &= 0 * (0 \diamond (w \diamond z)) \\ &= 0 * [(0 * w) * (0 \diamond z)] \\ &= (0 \diamond (0 * w)) \diamond (0 * (0 \diamond z)) \\ &= w \diamond z \end{aligned}$$

and

$$\begin{aligned} 0 * (0 \diamond (w * z)) &= 0 * (0 * (w * z)) \\ &= 0 * [(0 \diamond w) \diamond (0 * z)] \\ &= 0 \diamond [(0 \diamond w) \diamond (0 * z)] \\ &= (0 * (0 \diamond w)) * (0 \diamond (0 * z)) \\ &= w * z. \end{aligned}$$

(vii) \Rightarrow (viii): For any $u, w, z \in X$, we have

$$\begin{aligned}
w \diamond u &= 0 \diamond (0 * (w \diamond u)) \\
&= 0 \diamond ((z \diamond z) * (w \diamond u)) \\
&= 0 \diamond [(z * (w \diamond u)) \diamond z] \\
&= (0 * (z * (w \diamond u))) * (0 \diamond z) \\
&= (0 \diamond (z * (w \diamond u))) * (0 \diamond z) \\
&= (0 * (0 \diamond z)) \diamond (z * (w \diamond u)) \\
&= (0 * (0 \diamond (z \diamond 0))) \diamond (z * (w \diamond u)) \\
&= (0 \diamond (0 * (z \diamond 0))) \diamond (z * (w \diamond u)) \\
&= (z \diamond 0) \diamond (z * (w \diamond u)) \\
&= z \diamond (z * (w \diamond u)).
\end{aligned}$$

By a similar way, we obtain $z * (z \diamond (w * u)) = w * u$.

(viii) \Rightarrow (i): If $z * x = z \diamond x = 0$, then by (viii) we have $x = x * 0 = z * (z \diamond (x * 0)) = z * (z \diamond x) = z * 0 = z$. This shows that z is a pseudo atom of X . This completes the proof. \square

Corollary 4.3. *Let X be a pseudo Q -algebra. If a is a pseudo atom of X , then for all x of X , $a * x$ and $a \diamond x$ are pseudo atoms. Hence $L(X)$ is a pseudo subalgebra of X .*

Corollary 4.4. *Let X be a pseudo Q -algebra. For every x of X , there is a pseudo atom a such that $a \preceq x$, i.e., every pseudo Q -algebra is generated by a pseudo atom.*

Proposition 4.5. *A non-zero element $a \in X$ is a pseudo atom of a pseudo Q -algebra X if $\{0, a\}$ is a pseudo ideal of X .*

Proof. Let $x \preceq a$ for any $x \in X$. Then $x * a = x \diamond a = 0 \in \{0, a\}$. Since $x \in \{0, a\}$ is a pseudo ideal of X , we have $x = 0$ or $x = a$. Since $a \neq 0$, we obtain $x = a$. Hence a is a pseudo atom of X . \square

Proposition 4.6. *If non-zero element of a pseudo Q -algebra X is a pseudo atom, then any pseudo subalgebra of X is a pseudo ideal of X .*

Proof. Let S be a pseudo subalgebra of X and let $x, y * x, y \diamond x \in S$. By Theorem 4.2, we have $y = x * (x \diamond y) = x * (0 * (y \diamond x))$. Since $0, y \diamond x \in S$ and S is a pseudo subalgebra of X , we obtain $0 * (y \diamond x) \in S$. Hence $y = x * (0 * (y \diamond x)) \in S$. Thus any pseudo subalgebra of X is a pseudo ideal of X . \square

For pseudo atom a of a Pseudo Q -algebra X ,

$$V(a) := \{x \in X \mid a \preceq x\}$$

is called a *pseudo branch* of X .

Theorem 4.7. *Let X be a pseudo Q -algebra. Suppose that a and b are pseudo atoms of X . Then the following properties hold:*

- (i) For all $x \in V(a)$ and all $y \in V(b)$, $x * y \in V(a * b)$ and $x \diamond y \in V(a \diamond b)$.
- (ii) For all x and $y \in V(a)$, $x \diamond y, x * y \in K(X)$, where $K(X) = \{x \in X \mid 0 \preceq x\}$.
- (iii) If $a \neq b$, then for all $x \in V(a)$ and $y \in V(b)$, we have $x * y, x \diamond y \in K(X)$.
- (iv) For all $x \in V(b)$, $a * x = a * b$ and $a \diamond x = a \diamond b$.
- (v) If $a \neq b$, then $V(a) \cap V(b) = \emptyset$.

Proof. (i) For all $x \in V(a)$ and all $y \in V(b)$, by Proposition 3.6 and Theorem 4.2 we have

$$\begin{aligned}
(a * b) \diamond (x * y) &= [0 * (0 \diamond (a * b))] \diamond (x * y) \\
&= (0 \diamond (x * y)) * (0 \diamond (a * b)) \\
&= (0 * (x * y)) * (0 \diamond (a * b)) \\
&= ((0 \diamond x) \diamond (0 * y)) * (0 \diamond (a * b)) \\
&= ((0 * (0 \diamond (a * b))) \diamond x) \diamond (0 * y) \\
&= ((a * b) \diamond x) \diamond (0 * y) \\
&= ((a \diamond x) * b) \diamond (0 * y) \\
&= (0 * b) \diamond (0 * y) \\
&= (0 \diamond b) \diamond (0 * y) \\
&= 0 * (b * y) = 0 * 0 = 0
\end{aligned}$$

and

$$\begin{aligned}
(a \diamond b) * (x \diamond y) &= [0 \diamond (0 * (a \diamond b))] * (x \diamond y) \\
&= (0 * (x \diamond y)) \diamond (0 * (a \diamond b)) \\
&= (0 \diamond (x \diamond y)) \diamond (0 \diamond (a \diamond b)) \\
&= ((0 * x) * (0 \diamond y)) \diamond (0 \diamond (a \diamond b)) \\
&= ((0 \diamond (0 \diamond (a \diamond b))) * x) * (0 \diamond y) \\
&= ((0 \diamond (0 * (a \diamond b))) * x) * (0 \diamond y) \\
&= ((a \diamond b) * x) * (0 \diamond y) \\
&= ((a * x) \diamond b) * (0 \diamond y) \\
&= (0 \diamond b) * (0 \diamond y) \\
&= (0 * b) * (0 \diamond y) \\
&= 0 \diamond (b \diamond y) = 0 \diamond 0 = 0.
\end{aligned}$$

Hence $x * y \in V(a * b)$ and $x \diamond y \in V(a \diamond b)$.

(ii) and (iii) are simple consequences of (i).

(iv) For all $x \in V(b)$, by Theorem 4.2 we have $(a * x) \diamond (a * b) = (a \diamond (a * b)) * x = b * x = 0$. Moreover, $a * b$ is a pseudo atom by Corollary 4.3. Therefore $a * x = a * b$. Also we

get $(a \diamond x) * (a \diamond b) = (a * (a \diamond b)) * x = b * x = 0$. Moreover, $a \diamond b$ is a pseudo atom by Corollary 4.3. Therefore $a \diamond x = a \diamond b$.

(v) Let $a \neq b$ and $V(a) \cap V(b) \neq \emptyset$. Then there exists $c \in V(a) \cap V(b)$. By (i), we have $0 = c * c = c \diamond c \in V(a * b), V(a \diamond b)$. Hence $a * b = a \diamond b = 0$, which is a contradiction. Thus (v) is true. \square

References

- [1] S. S. Ahn and H. S. Kim, *On QS-algebras*, J. Chungcheong. Math. Soc., **12**(1999), 33–41.
- [2] M. A. Chaudhry, *On BCH-algebras*, Math. Japon., **4**(1991), 665–676.
- [3] J. M. Font, A. J. Rodríguez and A. Torrens, *Wajsbergs Algebras*, Stochastica, **8**(1984), 5–31.
- [4] G. Georgescu and A. Iorgulescu, *Pseudo BCK-algebras*, in: *Proc. DMTS'01: Combinatorics, Computability and Logics*, Springer, London, 2001, 97–114.
- [5] Q. P. Hu and X. Li, *On BCH-algebras*, Math. Seminar Notes, **11**(1983), 313–320.
- [6] Q. P. Hu and X. Li, *On proper BCH-algebras*, Math. Japon., **30**(1985), 659–661.
- [7] K. Iséki, *On BCI-algebras*, Math. Seminar Notes, **8**(1980), 125–130.
- [8] K. Iséki and S. Tanaka, *An introduction to the theory of BCK-algebras*, *Math. Japon.*, **23**(1978), 1–26.
- [9] Y. B. Jun, H. S. Kim and J. Neggers, *On pseudo d-algebras*, Inform. Sci., **179**(2009), 1751–1759.
- [10] Y. B. Jun, *Characterizations of pseudo BCK-algebras*, Sci. Math. Japon. Online, **7**(2002), 225–230.
- [11] J. Meng, *Implicative commutative semigroups are equivalent to a class of BCK-algebras*, Semigroup Forum, **50**(1995), 89–96.
- [12] J. Meng and Y. B. Jun, *BCK-algebras*, Kyung Moon Sa, Seoul, 1994.
- [13] D. Mundici, *MV-algebras are categorically equivalent to bounded commutative BCK-algebras*, Math. Japon., **31**(1986), 889–894.
- [14] J. Neggers, S. S. Ahn and H. S. Kim, *On Q-algebras*, Inter. J. Math. and Math. Sci., **27**(2001), 749–757.
- [15] E. H. Roh and S. Y. Kim, *On BH*-subalgebras of transitive BH*-algebras*, Far East J. Math. Sci., **1**(1999), 255–263.