

Aloe-emodin inhibits Pam₃CSK₄-induced MAPK and NF-κB signaling through TLR2 in macrophages*

Lee, Mi Jin^{1**} · Park, Mi-Young^{2**} · Kim, Soon-Kyung^{1†}

¹Department of Food Science and Nutrition, College of Natural Science, Soonchunhyang University, Asan 31538, Korea

²Department of Food & Nutrition Education, Graduate School of Education, Soonchunhyang University, Asan 31538, Korea

ABSTRACT

Purpose: Aloe-emodin (AE), an ingredient of aloe, is known to exhibit anti-inflammatory activities. However, little is known about the underlying molecular mechanisms of its inflammatory modulatory activity *in vitro*. In the present study, we investigated the anti-inflammatory potential of AE using Pam₃CSK₄-stimulated macrophages. **Methods:** RAW 264.7 macrophages were treated with AE (0~20 μM) for 1 h, followed by treatment with Pam₃CSK₄ for 1 h. After incubation, mRNA expression levels of cytokines were measured. The effect of AE on TLR2-related molecules was also investigated in Pam₃CSK₄-stimulated RAW 264.7 macrophages. **Results:** AE attenuated Pam₃CSK₄-stimulated expression of proinflammatory cytokines, including tumor necrosis factor-α (TNF-α), interleukin-6 (IL-6), and interleukin-1β (IL-1β) in RAW 264.7 macrophages. Two concentrations of AE (10 μM and 20 μM) effectively reduced mRNA expression of TLR2 by 41.18% and 54.43%, respectively, compared to that in control cells (*p* < 0.05). AE also decreased nuclear factor-κB (NF-κB) activation and mitogen-activated protein kinase (MAPK) phosphorylation. Phosphorylation levels of ERK1/2, p38, and JNK were markedly reduced by 20 μM AE. In particular, AE decreased phosphorylation of ERK in a dose-dependent manner in Pam₃CSK₄-stimulated RAW 264.7 macrophages. **Conclusion:** Our data indicate that AE exerts its anti-inflammatory effect by suppressing TLR2-mediated activation of NF-κB and MAPK signaling pathways in macrophages.

KEY WORDS: Aloe-emodin (AE), macrophage, mitogen-activated protein kinase (MAPK), nuclear factor-κB (NF-κB), Toll-like receptor 2 (TLR2)

INTRODUCTION

Aloe leaves are used in the treatment of several medical conditions, including burns, cancer, and inflammatory bowel disease.¹⁻³ Aloe-emodin (AE), a major anthraquinone present in aloe, exhibits antibacterial, antiviral, anti-inflammatory, and anticancer effects.⁴⁻⁶ The levels of AE and aloin, a C-glycoside derivative of AE, range from 0.1% to 21.5% of dry weight in leaf exudates of 68 Aloe species.^{7,8} Previous studies have used *in vitro* and *in vivo* models to study the anti-inflammatory activity of emodin. Park et al. reported that AE dose-dependently inhibited the levels of nitric oxide (NO) and prostaglandin E₂ (PGE₂) by blocking the mRNA expression of inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) in LPS-stimulated macrophages.⁹ Yin et al. reported that emodin

ameliorated lung injury via the inhibition of cytokine production and inhibition of the p38 mitogen-activated protein kinase (MAPK) pathway in an animal model.¹⁰ Iwanowycz et al. have reported that emodin bidirectionally regulates macrophage phagocytosis and migration by blocking the nuclear factor kappa B (NF-κB)/interferon regulatory factor 5 (IRF5)/signal transducer and activator of transcription 1 (STAT1) and interferon regulator factor 4 (IRF4)/signal transducer and activator of transcription 6 (STAT6) signaling pathways.¹¹ However, few studies have investigated the biological activity of AE, and its exact mechanisms have not been fully elucidated.

Chronic inflammation is associated with several diseases. Since macrophages are implicated in the initiation of inflammatory responses, they play an important role in inflammatory diseases.^{12,13} Therefore, the inhibition of

Received: July 12, 2016 / Revised: August 3, 2016 / Accepted: August 14, 2016

*This study was supported by Soonchunhyang University Research Fund.

**These authors contributed equally to this work.

†To whom correspondence should be addressed.

tel: +82-41-530-1261, e-mail: soon56@sch.ac.kr

© 2016 The Korean Nutrition Society

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

macrophage-mediated inflammatory responses is a useful therapeutic approach against several inflammatory diseases. Toll-like receptors (TLRs) are pattern recognition receptors that recognize several pathogen-associated molecular pattern (PAMP) molecules involved in pathogenic invasions.^{14,15} PAMP-recognizing TLRs stimulate signaling pathways that involve NF- κ B and MAPKs.

The aim of this study was to investigate the anti-inflammatory activity of AE against synthetic triacylated lipoprotein Pam₃CSK₄-stimulated RAW 264.7 macrophages.

MATERIALS AND METHODS

Reagents

AE and Pam₃CSK₄ were obtained from Sigma-Aldrich Co. (St. Louis, MO, USA). Antibodies against extracellular signal-regulated kinase (ERK), p38, c-Jun N-terminal kinase (JNK), phospho (p)-ERK (Thr²⁰²/Tyr²⁰⁴), p-p38 (Thr¹⁸⁰/Tyr¹⁸²), and p-JNK (Thr¹⁸³/Tyr¹⁸⁵) were purchased from Cell Signaling Technology (Beverly, MA, USA). All other chemicals were purchased from Sigma-Aldrich Co. (St. Louis, MO, USA).

Cell culture

The mouse macrophage cell line (RAW 264.7) was purchased from American Type Culture Collection (Manassas, VA, USA). Cells were grown in DMEM supplemented with 10% heat-inactivated fetal bovine serum, and 1% penicillin/streptomycin. Cells were incubated at 37°C, 5% CO₂ for 5–7 days until monolayers of macrophages were observed.

Real-time reverse transcription-polymerase chain reaction analysis (RT-PCR)

Total RNA was extracted using the TRIzol reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. Real-time quantitative polymerase chain reaction (PCR) was performed using a Quantitect™ SYBR Green PCR kit (Quantitect™ SYBR Green PCR, Qiagen, CA, USA). The specific primer sets were as follows: 5'-AAC-ATCCAACCTTCCCAAACG-3'/5'-CTCTTAACCCCGAATCCCAG-3' for the tumor necrosis factor alpha (TNF- α) gene, 5'-TCACCTCTTCAGAACGAATTGACA-3'/5'-AGTGCCTCTTGCTGCTTTCAC-3' for the interleukin 6 (IL-6) gene, 5'-ATTGGGATCATCTTGCTGGT-3'/5'-CCTGCTGTTACAGTTGCC-3' for the interleukin 1 β

(IL-1 β) gene, and 5'-GAGCGCAAGTACTCTGTGTG-3'/5'-CGGACTCATCGTACTCCTG-3' for the β -actin gene used as an endogenous control. The relative mRNA expression levels of cytokines were normalized to that β -actin using the $\Delta\Delta$ CT method.¹⁶

NF- κ B activity

To determine the NF- κ B activity, the nucleus of macrophages were extracted using a Nuclear Extract kit (Active Motif, Carlsbad, CA, USA) and analyzed using a PathScan Total NF- κ B p65 assay kit (Cell Signaling Technology, MA, USA) according to the manufacturer's instructions.

Western blotting

Whole cell extract was prepared by suspending in an extraction lysis buffer and the cellular proteins were extracted with Laemmli sample buffer. Proteins were separated on 10% sodium dodecyl sulfate (SDS) polyacrylamide gels electrophoresis and then transferred to polyvinylidene fluoride (PVDF) membrane. The membranes were incubated with 5% nonfat dry milk, followed by incubation with appropriate primary antibodies in 5% nonfat dry milk in 0.05% Tris-buffered saline with Tween 20 (TBS-T) at 4°C overnight. The membranes were washed three times with TBS-T and then incubated for 1 h at room temperature with appropriate horseradish peroxidase-conjugated secondary antibodies. The membranes were visualized by chemiluminescence (ECL) and the density of the blots was quantified by ChemiDoc MP imaging system (Bio-Rad, Hercules, CA, USA).

Statistical analysis

All of the data are expressed as mean \pm standard deviation (SD). We analyzed our data with the by one-way analysis of variance, followed by Duncan's multiple range tests. $P < 0.05$ was considered to be significant.

RESULTS

TNF- α mRNA expression

To evaluate whether AE can inhibit the gene expression of cytokines, we measured the TNF- α mRNA level in AE-pretreated RAW 264.7 macrophages after their stimulation with Pam₃CSK₄. We observed that Pam₃CSK₄ upregulated the TNF- α mRNA expression, but the overexpression was

inhibited by both 10 μ M and 20 μ M AE (Fig. 1).

TNF- α , IL-6 and IL-1 β mRNA expression

Next, we measured the mRNA expression levels of pro-inflammatory cytokines in Pam₃CSK₄-stimulated RAW 264.7 macrophages by real-time PCR. Our results showed that AE at concentrations of 1-20 μ M significantly inhibited the mRNA expression levels of all cytokines studied (Fig. 2; $p < 0.05$). AE significantly decreased the mRNA expression levels of IL-6 and IL-1 β in a dose-dependent manner in this cell model (Fig. 2B, C; $p < 0.05$). Among the three proinflammatory cytokines, TNF- α was most effectively blocked by AE treatment of macrophages (Fig. 2A; $p < 0.05$).

TLR2 mRNA expression

Next, we evaluated the inhibitory effect of AE on the Pam₃CSK₄-stimulated TLR2 mRNA expression in macrophages. AE significantly inhibited the Pam₃CSK₄-stimulated TLR2 mRNA overexpression at both 10 μ M and 20 μ M




Fig. 1. Effect of AE on Pam₃CSK₄-induced TNF- α mRNA expression in RAW 264.7 macrophages. RAW 264.7 macrophages were pretreated with 0~20 μ M AE for 1 h. The cells were further stimulated with Pam₃CSK₄ (1 μ g/mL). After 1 h, the TNF- α mRNA expression was analyzed in an agarose gel.

concentrations (Fig. 3; $p < 0.05$). Pretreatment with 10 μ M and 20 μ M AE decreased the mRNA expression of TLR2 by 41.18% and 54.43%, respectively, compared to that in the control cells (Fig. 3; $p < 0.05$).

NF- κ B activation and MAPK phosphorylation

To understand the molecular mechanism(s) underlying the TLR2-blocking effect, NF- κ B and MAPK activation were examined in RAW 264.7 macrophages. Our results showed that the NF- κ B activity was dramatically upregulated by the Pam₃CSK₄ treatment compared with the activity in the Pam₃CSK₄-stimulated control group (Fig. 4A). AE at 10 μ M and 20 μ M reduced the NF- κ B activity by 26.83% and 46.34%, respectively, compared with that in




Fig. 3. Effect of AE on the Pam₃CSK₄-induced TLR2 mRNA expression in RAW 264.7 macrophages. RAW 264.7 macrophages were pretreated with 0~20 μ M AE for 1 h and further stimulated with Pam₃CSK₄ (1 μ g/mL). After 1 h, TLR2 mRNA expression was determined. The data are expressed as a fold induction relative to the vehicle-treated cells. The values are the mean \pm SD (n = 4). *Significantly different from the Pam₃CSK₄-stimulated control, $p < 0.05$.




Fig. 2. Effect of AE on the Pam₃CSK₄-induced TNF- α (A), IL-6 (B), and IL-1 β (C) mRNA expression in RAW 264.7 macrophages. RAW 264.7 macrophages were pretreated with 0~20 μ M AE for 1 h and further stimulated with Pam₃CSK₄ (1 μ g/mL). After 1 h, TNF- α , IL-6, and IL-1 β mRNA expression was determined. The data are expressed as a fold induction compared with the vehicle-treated cells. The values are the mean \pm SD (n = 3). *Significantly different from the Pam₃CSK₄-stimulated control ($p < 0.05$).




Fig. 4. Effect of AE on the NF- κ B activation (A) and MAPK expression (B) in RAW 264.7 macrophages. RAW 264.7 macrophages were pretreated with 0–20 μ M AE for 1 h and further stimulated with Pam₃CSK₄ (1 μ g/ml). After 1 h, NF- κ B activity was determined (A), and cell lysates were analyzed by MAPK immunoblotting (B). The NF- κ B activity data are expressed as % activation compared with the vehicle-treated cells. The values are the mean \pm SD (n = 4). *Significantly different from the Pam₃CSK₄-stimulated control, p < 0.05.

the Pam₃CSK₄-stimulated control cells (Fig. 4A; p < 0.05).

Next, we examined the MAPK phosphorylation in Pam₃CSK₄-stimulated RAW 264.7 macrophages. The phosphorylation of ERK1/2, p38, and JNK was reduced by 20 μ M AE (Fig. 4B). In particular, AE decreased the ERK phosphorylation in a dose-dependent manner in the Pam₃CSK₄-stimulated RAW 264.7 macrophages. These results indicated that the anti-inflammatory efficacy of AE on RAW 264.7 is associated with the inactivation of NF- κ B, as well as with blocking of MAPK phosphorylation.

DISCUSSION

TLRs are pattern recognition molecules, which represent major components of the innate immune response. Their specific signaling pathways are associated with several inflammatory diseases.¹⁷ In particular, TLR2 is widely distributed on the surface of several types of immune cells, including macrophages, dendritic cells, and mast cells.^{18–20} The receptor is the main sensor for PAMP recognition of gram-positive bacteria. Pam₃CSK₄ binds to the host TLR2 and leads to an inflammatory reaction.²¹ In this study, we investigated whether AE can inhibit the Pam₃CSK₄-induced, TLR2-regulated signaling in RAW 264.7 macrophages.

To test the anti-inflammatory potential of AE, the mRNA expression level of the inflammatory marker TNF- α was measured in AE-pretreated RAW 264.7 macrophages after stimulation with Pam₃CSK₄. Our results showed that AE downregulated the TNF- α mRNA expression at concentrations of 10 μ M and 20 μ M (Fig. 1). Our results also

showed that AE at concentrations of 5–20 μ M effectively suppressed the mRNA levels for all the cytokines (TNF- α , IL-6, and IL-1 β) in RAW 264.7 macrophages (Fig. 2). These data indicated that AE protects against the Pam₃CSK₄-induced inflammatory response in macrophages. Related *in vivo* studies have shown that emodin significantly ameliorated inflammatory responses through inhibition of cytokine overproduction. Nemmar et al. have suggested that emodin administration protects from diesel exhaust particle-induced lung inflammation via inhibition of TNF- α , IL-6, and IL-1 β in mice.²² According to Han et al., the treatment with emodin decreased the IL-1 β secretion by blocking the activation of the NLRP3 inflammasome in alipolysaccharide-induced endotoxin mouse model.²³

To understand the mechanism of AE in regulating Pam₃CSK₄-induced cytokine gene expression, we examined the TLR2 mRNA expression in Pam₃CSK₄-stimulated macrophages. We found that AE effectively reduced upregulated expression of TLR2 (Fig. 3). These results indicate that AE down-regulates TLR2-mediated cytokine induction. There are a few studies evaluating the inhibitory effect of AE on TLR2-mediated signaling pathways. Li et al. reported that emodin from the Chinese herb Radix et Rhizoma Rhe significantly decreased the expression of cytokine genes by blocking TLR2 signaling pathways in rat kidney epithelial cells.²⁴

The effect of AE in TLR-2 pathways may have associated with the inhibition of the NF- κ B signaling pathway, which is the major transcription pathway for inflammatory responses.²⁵ The results showed that NF- κ B activity was dramatically enhanced by Pam₃CSK₄ treatment compared

with the activity in the Pam₃CSK₄-negative control group; however, this enhancement of the activity was effectively alleviated in the 10 μM and 20 μM AE-treated groups (Fig. 4A). A previous study indicated that AE effectively decreased the activation of p38 and NF-κB in a concanavalin A-induced animal hepatitis model.²⁶ Although there are few data on AE, several *in vivo* studies have reported that emodin effectively ameliorates inflammatory diseases via inactivation of NF-κB.^{27,28} This evidence indicates that AE is a potential anti-inflammatory agent that inhibits both NF-κB activation and gene expression of proinflammatory cytokines.

Because the activation of NF-κB as well as MAPK pathways are implicated in TLR2 signaling, we evaluated the protein levels of p-ERK1/2, p-p38, and p-JNK in Pam₃CSK₄-stimulated macrophages.

Pam₃CSK₄ stimulation upregulated the phosphorylations of the MAPKs in RAW 264.7 macrophages. We have also observed that 20 μM AE treatment prevented an increased expression of MAPKs (Fig. 4B), which is consistent with previous observations showing that emodin suppresses the activation of p38 and ERK1/2.^{29,30} These results demonstrated that AE shows an anti-inflammatory action that is mediated by a TLR2-dependent MAPK signaling pathway in macrophages.

SUMMARY

Our data established that AE exhibited anti-inflammatory effects against Pam₃CSK₄-induced inflammation in macrophages. Its activity was partially due to the suppression of the NF-κB and MAPKs signaling pathways.

REFERENCES

1. Farzadnia P, Jofreh N, Khatamsaz S, Movahed A, Akbarzadeh S, Mohammadi M, Bargahi A. Anti-inflammatory and wound healing activities of aloe vera, honey and milk ointment on second-degree burns in rats. *Int J Low Extrem Wounds* 2016; 15(3): 241-247.
2. Al-Oqail MM, El-Shaibany A, Al-Jassas E, Al-Sheddi ES, Al-Massarani SM, Farshori NN. In vitro anti-proliferative activities of Aloe perryi flowers extract on human liver, colon, breast, lung, prostate and epithelial cancer cell lines. *Pak J Pharm Sci* 2016; 29(2 Suppl): 723-729.
3. Park MY, Kwon HJ, Sung MK. Dietary aloin, aloesin, or aloe-gel exerts anti-inflammatory activity in a rat colitis model. *Life Sci* 2011; 88(11-12): 486-492.
4. Andersen DO, Weber ND, Wood SG, Hughes BG, Murray BK, North JA. In vitro virucidal activity of selected anthraquinones and anthraquinone derivatives. *Antiviral Res* 1991; 16(2): 185-196.
5. Arosio B, Gagliano N, Fusaro LM, Parmeggiani L, Tagliabue J, Galetti P, De Castri D, Moscheni C, Annoni G. Aloe-Emodin quinone pretreatment reduces acute liver injury induced by carbon tetrachloride. *Pharmacol Toxicol* 2000; 87(5): 229-233.
6. Esmat AY, Tomasetto C, Rio MC. Cytotoxicity of a natural anthraquinone (Aloin) against human breast cancer cell lines with and without ErbB-2: topoisomerase IIalpha coamplification. *Cancer Biol Ther* 2006; 5(1): 97-103.
7. Groom QJ, Reynolds T. Barbaloin in aloe species. *Planta Med* 1987; 53(4): 345-348.
8. van Wyk BE, van Rheede van Oudtshoorn MC, Smith GF. Geographical variation in the major compounds of Aloe ferox leaf exudate. *Planta Med* 1995; 61(3): 250-253.
9. Yin JT, Wan B, Liu DD, Wan SX, Fu HY, Wan Y, Zhang H, Chen Y. Emodin alleviates lung injury in rats with sepsis. *J Surg Res* 2016; 202(2): 308-314.
10. Iwanowycz S, Wang J, Altomare D, Hui Y, Fan D. Emodin bidirectionally modulates macrophage polarization and epigenetically regulates macrophage memory. *J Biol Chem* 2016; 291(22): 11491-11503.
11. Nathan C, Ding A. Nonresolving inflammation. *Cell* 2010; 140(6): 871-882.
12. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 2008; 8(12): 958-969.
13. Kaisho T, Akira S. Critical roles of Toll-like receptors in host defense. *Crit Rev Immunol* 2000; 20(5): 393-405.
14. Barton GM, Medzhitov R. Toll-like receptor signaling pathways. *Science* 2003; 300(5625): 1524-1525.
15. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001; 25(4): 402-408.
16. Salvador B, Arranz A, Francisco S, Córdoba L, Punzón C, Llamas MÁ, Fresno M. Modulation of endothelial function by Toll like receptors. *Pharmacol Res* 2016; 108(9): 46-56.
17. McCurdy JD, Olynych TJ, Maher LH, Marshall JS. Cutting edge: distinct Toll-like receptor 2 activators selectively induce different classes of mediator production from human mast cells. *J Immunol* 2003; 170(4): 1625-1629.
18. Redecke V, Häcker H, Datta SK, Fermin A, Pitha PM, Broide DH, Raz E. Cutting edge: activation of Toll-like receptor 2 induces a Th2 immune response and promotes experimental asthma. *J Immunol* 2004; 172(5): 2739-2743.
19. Takeda K, Akira S. Toll-like receptors in innate immunity. *Int Immunol* 2005; 17(1): 1-14.
20. Takeda K, Akira S. TLR signaling pathways. *Semin Immunol* 2004; 16(1): 3-9.
21. Nemmar A, Al-Salam S, Yuvaraju P, Beegam S, Ali BH. Emodin mitigates diesel exhaust particles-induced increase in airway resistance, inflammation and oxidative stress in mice. *Respir Physiol Neurobiol* 2015; 215(5): 51-57.
22. Han JW, Shim DW, Shin WY, Heo KH, Kwak SB, Sim EJ, Jeong JH, Kang TB, Lee KH. Anti-inflammatory effect of emodin via attenuation of NLRP3 inflammasome activation. *Int J Mol Sci* 2015; 16(4): 8102-8109.
23. Gambhir V, Yildiz C, Mulder R, Siddiqui S, Guzzo C, Szewczuk

- M, Gee K, Basta S. The TLR2 agonists lipoteichoic acid and Pam3CSK4 induce greater pro-inflammatory responses than inactivated *Mycobacterium butyricum*. *Cell Immunol* 2012; 280(1): 101-107.
24. Li Y, Xiong W, Yang J, Zhong J, Zhang L, Zheng J, Liu H, Zhang Q, Ouyang X, Lei L, Yu X. Attenuation of inflammation by emodin in lipopolysaccharide-induced acute kidney injury via inhibition of toll-like receptor 2 signal pathway. *Iran J Kidney Dis* 2015; 9(3): 202-208.
 25. Dahiya Y, Pandey RK, Sodhi A. Nod2 downregulates TLR2/1 mediated IL1 β gene expression in mouse peritoneal macrophages. *PLoS One* 2011; 6(11): e27828.
 26. Xue J, Chen F, Wang J, Wu S, Zheng M, Zhu H, Liu Y, He J, Chen Z. Emodin protects against concanavalin A-induced hepatitis in mice through inhibiting activation of the p38 MAPK-NF- κ B signaling pathway. *Cell Physiol Biochem* 2015; 35(4): 1557-1570.
 27. Yao WY, Zhou YF, Qian AH, Zhang YP, Qiao MM, Zhai ZK, Yuan YZ, Yang SL. Emodin has a protective effect in cases of severe acute pancreatitis via inhibition of nuclear factor κ B activation resulting in antioxidation. *Mol Med Rep* 2015; 11(2): 1416-1420.
 28. Xiao M, Zhu T, Zhang W, Wang T, Shen YC, Wan QF, Wen FQ. Emodin ameliorates LPS-induced acute lung injury, involving the inactivation of NF- κ B in mice. *Int J Mol Sci* 2014; 15(11): 19355-19368.
 29. Gao Y, Zhang J, Li G, Xu H, Yi Y, Wu Q, Song M, Bee YM, Huang L, Tan M, Liang S, Li G. Protection of vascular endothelial cells from high glucose-induced cytotoxicity by emodin. *Biochem Pharmacol* 2015; 94(1): 39-45.
 30. Yin X, Gong X, Jiang R, Kuang G, Wang B, Zhang L, Xu G, Wan J. Emodin ameliorated lipopolysaccharide-induced fulminant hepatic failure by blockade of TLR4/MD2 complex expression in D-galactosamine-sensitized mice. *Int Immunopharmacol* 2014; 23(1): 66-72.