

SHORT REPORT

Open Access



Characterization of the complete mitochondrial genome of Mauritian sardinella, *Sardinella jussieu* (Lacepède, 1803), collected in the Banten Bay, Indonesia

Sinar Pagi Sektiana^{1,3}, Sapto Andriyono^{1,4,5} and Hyun-Woo Kim^{1,2*} 

Abstract

Fishes in genus *Sardinella* are small pelagic species, which plays an important role in marine ecosystem as the first consumer. Those species are also commercially important, whose total catch reaches 278,600 tons in 2011 in Indonesia, but their identification has been difficult for their morphological similarity. In this study, we reported *Sardinella jussieu* for the first time in Indonesian coastal area (Banten Bay, Indonesia, 6° 0' 50.00" S–106° 10' 21.00" E). We were able to confirm the species by both its morphological characteristics including the black spot at dorsal fin origin, the dusky pigmentation at caudal fin, 31 total scute numbers, and DNA sequence identity in the GenBank database by the molecular analysis. Its total mitochondrial genome was determined by the combination of next-generation sequencing and typical PCR strategy. The total mitochondrial genome of *Sardinella jussieu* (16,695 bp) encoded 13 proteins, 2 ribosomal RNAs, 22 transfer RNAs, and the putative control region. All protein-coding genes started with ATG and typical stop codon and ended with TAA or TAG except for ND4 in which AGA is used. Phylogenetic analyses of both COI region and full mitochondrial genome showed that *S. jussieu* is most closely related to *Sardinella albella* and *Sardinella gibbosa*.

Keywords: Barcode, Mitochondrial genome, *Sardinella*, Indonesia, Next-generation sequencing

Background

Sardinella is a genus of fish in the family Clupeidae found in the Atlantic, Indian, and the Pacific Ocean. The paddle-shaped supramaxilla bones are major characteristics, which help distinguish *Sardinella* from other genera. Morphological characters distinguish *Sardinella* from all other *clupeoid* genera with the presence of two fleshy outgrowths on the hind margin of the gill opening (Whitehead 1985). According to FishBase (<http://www.fishbase.org/>), there are currently 22 recognized species in the genus

Sardinella. *Sardinella* is important not only in marine food webs as a base consumer supporting tuna, seabirds, and marine mammals (Willette et al. 2011) but also in industry as the protein source with a low cost using as a bait for large fish or a feed in aquaculture.

Seven species in the genus *Sardinella* are currently known in Indonesian waters including *Sardinella fimbriata*, *Sardinella gibbosa*, *Sardinella lemuru*, *Sardinella albella*, *Sardinella atricauda*, *Sardinella branchysoma*, and *Sardinella melanura*, whose total catch in Indonesia reaches 278,600 tons in 2011 (MMAF 2012). Morphological identification in *Sardinella* is mainly characterized by their gill raker, pelvic scute, scales, and otolith (Homayuni et al. 2013; Bräger and Moritz

* Correspondence: kimhw@pkn.ac.kr

¹Interdisciplinary Program of Biomedical, Mechanical and Electrical Engineering, Pukyong National University, Busan 48513, Republic of Korea

²Department of Marine Biology, Pukyong National University, Busan 48513, Republic of Korea

Full list of author information is available at the end of the article



2016; Begg and Waldman 1999). However, species identification in the genus *Sardinella* is often hard for its broad geographical ranges, overlapping distributions (Willette et al. 2011) and morphological similarities (Sivakumaran et al. 1987) especially in larval stages (Ditty et al. 1994), which makes it difficult to manage the *Sardinella* resources in Indonesia.

In addition to the traditional morphological identification, the genetic information is now alternatively used for the species identification for its fast and exact results. The most widely used genetic markers are partial mitochondrial DNA sequences such as cytochrome C oxidase I (COI) or cytochrome B (CytB) (Palumbi et al. 1991; Ward et al. 2005; Vrijenhoek 1994). However, full mitochondrial genome sequences provide more information about its biogeographical or evolutionary information than those fragmental sequences. Therefore, more than 5000 mitochondrial genomes have been deposited in GenBank database (www.ncbi.nlm.nih.gov) from 33,500 species identified based on morphological characteristics (www.fishbase.org).

In this study, we report the Mauritian sardinella, *Sardinella jussieu*, for the first time in Indonesian coastal waters, which was collected from the Banten Bay. *S. jussieu* was previously reported only in the Western Indian Ocean, Taiwan, Hong Kong, and Vietnam (www.fishbase.org). Morphological characteristics of *Sardinella jussieu* are distinguished within other *Sardinella* species with the presence of black spot at dorsal fin origin and dusky pigmentation at caudal fin, total scute measurement which is 31, and vertical striae on a scale not meeting at center and

no perforation on hind part (Whitehead 1985). After confirmation of the species by the molecular COI markers, its total mitochondrial genome sequence was determined by the combination of the traditional PCR methods and next-generation sequencing (NGS) techniques.

Methods

Sample collection and morphological measurement

Five individuals of *S. jussieu* were collected in the Banten Bay, Indonesia ($6^{\circ} 0' 50.00''$ – S $106^{\circ} 10' 21.00''$ E), in January 2016 as the part of the regular fish survey (Fig. 1). Collected fish were directly stored in 96% ethanol and kept at $-20^{\circ}C$ until the further analysis (Knebelsberger and Stöger 2012). Morphological identification was made by their body shape, type of scale, fin feature, morphometric (i.e., standard length, body width, and head length), and meristic characteristic (total number of scutes) (Whitehead 1985; Strauss and Bond 1990).

Genomic DNA extraction and next-generation sequencing

Genomic DNA was extracted using an AccuPrep® Genomic DNA Extraction Kit (Bioneer) according to the manufacturer's instruction. A small portion of tail fin was dissected, which was further homogenized by the TissueLyser II (Qiagen). Purified genomic DNA was quantified with nanoDrop (ThermoFisher Scientific D1000), aliquoted, and stored at the $-70^{\circ}C$ for further analysis.

Two universal primer sets targeting cytochrome c oxidase I (COI) region, Fish F1 and Fish R1 (Ward et al. 2005), and targeting cytochrome b (cyt-B) region,




Fig. 1 Sampling location in Banten Bay, Indonesia (red square)

Table 1 Primers used for the mitochondrial genome of *Sardinella jussieu*

Name	Sequence (5' to 3')	Product size (bp)
Fish F1	TCAACCAACCACAAAGACATTGGCAC	652
Fish R1	TAGACTTCTGGGTGGCCAAAGAATCA	
GLUDG-L	TGACTTGAARAAYCGTTG	451
GB2-H	CCCTCAGAATGATATTTGTCCTCA	
CYT-B-F	GCCTACGAAAAACCCACCCGCTCC	8.2 k
CO1-R	GTAAGTCTACGGATGCCCTGCG	
CYT-B-R	AACGGAGGAGAAAGCGGTTGCGATG	8.7 k
CO1-F	CTTCCTGCTTCTCTGGCCTCCTC	
SARD F	TTAAAGTCTCCCTGAGGCC	683
SARD R	TTAGGAGGGAGTCGTCAAATGC	

GLUDG-L and CB2-H (Palumbi et al. 1991), were used to obtain the partial sequences of each gene, respectively (Table 1). The quality of all the primers used in this experiment was analyzed by the OligoAnalyzer 3.1 (<http://sg.idtdna.com/calc/analyzer>) and commercially synthesized by Bioneer Co. (Korea). Each PCR mixture (20 μ L) contained 12.8 μ L ultrapure water, 1 μ L primer (0.5 μ M, forward and reverse), 0.2 μ L Ex Taq DNA polymerase (TaKaRa, Japan), 2 μ L 10 \times Buffer, 2 μ L dNTPs (1 μ M, TaKaRa, Japan), and 100 ng genomic DNA as template. PCR was carried out under the following condition: initial denaturation step at 95 $^{\circ}$ C for 3 min, followed by 35 cycles of denaturation at 95 $^{\circ}$ C for 30 s, annealing at 50 $^{\circ}$ C for 30 s, and extension at 72 $^{\circ}$ C for 45 s (COI target sequence) or 30 s (Cyt-B target sequence). The process was completed with a final extension at 72 $^{\circ}$ C for 10 min. Two PCR products targeting partial sequences of COI and Cyt B were then purified with AccuPrep Gel purification kit (Bioneer, Korea) and ligated into a cloning vector (Promega, USA), sequenced in both directions.

In order to obtain two large PCR products (~8 kb), two pairs of sequence-specific primer sets (CYT-F and CO1-R and CO1-F and CYT B-R) were designed based on the obtained partial sequences of each region

(Table 1). Each PCR reaction (30 μ L) contained 19.7 μ L ultrapure water, 1 μ L of each primer (0.5 μ M), 0.3 μ L Ex Taq Hot Start Version DNA polymerase (TAKARA, Japan), 3 μ L 10 \times Buffer, 3 μ L dNTPs (1 mM, Takara, Japan), and 100 ng genomic DNA as template. PCR was carried out with two-step PCR protocol for long PCR under the following condition: initial denaturation step at 94 $^{\circ}$ C for 3 min, followed by 30 cycles of denaturation at 98 $^{\circ}$ C for 10s, and annealing and extension at 68 $^{\circ}$ C for 10 min. The process was completed with a final extension at 72 $^{\circ}$ C for 10 min. Two large PCR products were pooled together in equal concentration and fragmented to ~350 bp in length by Covaris M220 (Covaris Inc.). TruSeq[®] sample preparation kit V2 (Illumina, USA) was used for the construction of a library from fragmented sequence and quality and quantity of the constructed library was measured using 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA). Sequencing was performed by Illumina Miseq platform (2 \times 300 bp pair ends) (Illumina, USA).

Assembly of mitochondrial genome by the bioinformatic analysis

Raw reads from MiSeq sequencer, with under Qv 20 and more than ambiguous nucleotides, were removed from raw read using CLC Genomic Workbench v 7.5 (CLC BIO Aarhus, Denmark). Mothur software was used to pairing forward and reverse sequence with more than 7 bp overlapped and without any mismatch. Paired sequence then assembled using Geneious R8 with minimum 20 bp of overlapping sequence and 100% overlap identity. Ambiguous sequences of the D-loop region were reconfirmed by the typical end-point PCR and with sequence-specific primers (Sard_F and Sard_R) and DNA sequencing of its PCR products by Sanger sequencing method (Table 1).

Results and discussion

Morphological and molecular identification of *Sardinella jussieu*

As the result of morphometric measurements, we determined that the collected five fish were *S. jussieu*.

Table 2 General morphometric and meristic (total scute) of *S. jussieu*

Sample	Measurement								Total scute
	Standard length/SL (mm)	Body depth/BD (mm)	Head length/HL (mm)	Eye diameter/ED (mm)	BD/SL (%)	HL/SL (%)	ED/SL (%)	HL/ED (%)	
1	50	13.5	12	3.5	27.0	24.0	7.0	29.2	30
2	47	12.5	11	3	26.6	23.4	6.4	27.3	31
3	45	12.5	11	3	27.8	24.4	6.7	27.3	31
4	41	11.5	11	3	28.0	26.8	7.3	27.3	31
5	52	14.5	13	3.5	27.9	25.0	6.7	26.9	32
Average	47	12.9	11.6	3.2	27.5	24.7	6.8	27.6	31




Fig. 2 Mauritian sardinella (*S. jussieu*) collected from Banten Bay, Indonesia (a). The fish scale of *S. jussieu* presents no perforations and vertical striated with not meeting at center (b) according to Whitehead (1985) (c). Black scale bar = 1 cm

Among the morphologically similar fish *Sardinella* species including *S. albelli*, *S. atricauda*, *S. fimbriata*, *S. marquesensis*, *S. sindensis*, and *S. gibbosa*, the scale and pigmentation patterns are useful characteristics to identify species (Bräger and Moritz 2016; Strauss and Bond 1990). The average ratio of body depth (BD) to standard length (SL) of the collected samples was 27.5%, and total scute numbers were 31 (Table 2). Vertical striae on scales did not meet at center with no perforations on hind part of the scale, and the pigmented dorsal and caudal fins were also identified (Fig. 2). Those morphological characteristics suggested that the collected samples were *S. jussieu*. The most closely related *Sardinella* species, *S. albelli* and *S. gibbosa*, are distinguished from *S. jussieu* in the presence of scale perforations

(Table 3). Molecular identification of five *Sardinella* samples confirmed the morphological identification. The COI region of five individuals (652 bp) exhibited 100% sequence identity to *Sardinella* sp. (GenBank accession number: KJ566769) collected from the coastal water in Thailand and 99% to *S. jussieu* (GenBank accession no.: HQ231358) collected from the Philippines (Quilang et al. 2011). Based on the morphological characteristics and DNA sequence identity, we concluded that five *Sardinella* samples collected in the Banten Bay, Indonesia, were Mauritian sardinella, *Sardinella jussieu*.

Complete mitochondrial genome of the *Sardinella jussieu*
 In order to have additional information of *S. jussieu*, the complete mitochondrial genome sequence was

Table 3 Comparison of morphological characteristic of seven *Sardinella* species

Name	Scale		Fin	
	Striae connected/overlapped	Perforations	Dark spot at dorsal fin origin	Dark spot at caudal fin
<i>S. fimbriata</i>		√	√	√
<i>S. gibbosa</i>		√	√	√
<i>S. albelli</i>		√	√	√
<i>S. atricauda</i>		√	√	√
<i>S. brachysoma</i>	√	√	√	√
<i>S. melanura</i>	√	√		√
<i>S. jussieu</i>			√	√

Table 4 Organization of the full-length mitochondrial genome of *Sardinella jussieu*

Feature	Position numbers	Size (bp)	Strand	Intergenic nucleotides	Codon		Anticodon/position
					Start	Stop	
tRNA-Phe	1–68	68	H	–	–	–	GAA/31–33
12S rRNA	69–1019	951	H	0	–	–	–
tRNA-Val	1020–1091	72	H	0	–	–	TAC/1053–1055
16S rRNA	1092–2777	1686	H	0	–	–	–
tRNA-Leu	2779–2853	75	H	1	–	–	TAA/2814–2816
ND1	2854–3828	975	H	0	ATG	TAG	–
tRNA-Ile	3837–3908	72	H	8	–	–	GAT/3869–3871
tRNA-Gln	3908–3978	71	L	–1	–	–	TTG/3944–3946
tRNA-Met	3978–4046	69	H	–1	–	–	CAT/4008–4010
ND2	4020–5093	1074	H	–27	ATG	TAG	–
tRNA-Trp	5092–5163	72	H	–2	–	–	TCA/5125–5127
tRNA-Ala	5165–5233	69	L	1	–	–	TGC/5173–5175
tRNA-Asn	5236–5308	73	L	2	–	–	GTT/5273–5275
tRNA-Cys	5345–5410	66	L	36	–	–	GCA/5366–5368
tRNA-Tyr	5414–5484	71	L	3	–	–	GTA/5450–5452
COX1	5678–7036	1359	H	193	ATG	TAA	–
tRNA-Ser	7037–7107	71	L	0	–	–	TGA/7073–7075
tRNA-Asp	7112–7180	69	H	4	–	–	GTC/7142–7144
COII	7193–7897	705	H	12	ATG	TAA	–
tRNA-Lys	7884–7957	74	H	–14	–	–	TTT/7918–7920
ATP8	7959–8126	168	H	1	ATG	TAA	–
ATP6	8117–8800	684	H	–10	ATG	TAA	–
COIII	8800–9585	786	H	–1	ATG	TAA	–
tRNA-Gly	9585–9656	72	H	0	–	–	TCC/9618–9620
ND3	9600–10007	408	H	–57	ATG	TAG	–
tRNA-Arg	10006–10075	70	H	–2	–	–	TCG/1037–10039
ND4L	10076–10372	297	H	0	ATG	TAA	–
ND4	10366–11751	1386	H	–7	ATG	AGA	–
tRNA-His	11747–11815	69	H	–5	–	–	GTG/11747–11815
tRNA-Ser	11816–11883	68	H	0	–	–	GCT/11842–11844
tRNA-LEu	11884–11955	72	H	0	–	–	TAG/11916–11918
ND5	11956–13791	1836	H	0	ATG	TAA	–
ND6	13788–14309	522	L	–4	ATG	TAG	–
tRNA-Glu	14310–14378	69	L	0	–	–	TTC/14346–14348
CYTB	14385–15581	1197	H	6	ATG	TAA	–
tRNA-Thr	15526–15597	72	H	–56	–	–	TGT/15558–15560
tRNA-Pro	15597–15666	70	L	–1	–	–	TGG/15620–15622
Control region	15667–16695	1029	H	0	–	–	–

longiceps (958 bp) (GenBank accession number: NC033407), *S. albella* (986 bp) (GenBank accession number: NC016726), and *S. maderensis* (986 bp) (GenBank accession number: NC009587).

Total mitochondrial DNA sequence of *S. jussieu* showed 84–93% identity with those of currently known three other *Sardinella* species among which *S. albella* is the most closely related to *S. jussieu* (Fig. 5a). In




Fig. 4 Putative secondary structure tRNA genes in mitochondrial genomic. Proposed structure of 22 tRNA genes encoded in the mitochondrial of *Sardinella jussieu*

order to know the better evolutionary relationship of *S. jussieu*, its COI sequence was compared with those of the other 12 *Sardinella* species (Fig. 5b). As shown in the analysis by the full mitochondrial genomes, *S. jussieu* showed the most closely related to *S. albella* with 96% sequence identity. In fact, DNA sequence identity of two species *S. albella* and *Sardinella gibbosa* was too high to be distinct with each other in the COI region (Figs. 5b). Although morphological keys to discriminate two species were proposed, the post-pelvic ventral scutes and gill rakers number on a lower limb (Stern et al. 2016), *S. albella* and *S. gibbosa* frequently misidentified as shown in the COI barcodes. From the reason, it is required to compare full-length mitochondrial sequences of two species for the better classification. As the lowest sequence identity to other *Sardinella* species, control region of *S. jussieu*

mitochondrial genome would be the good candidate to discriminate them.

In this study, we identified that *S. jussieu* inhabits in Java island, Indonesia, as well as the two previously known *Sardinella* species, *S. albella* and *S. gibbosa*. Although *S. jussieu* is originally distributed in the western Indian Ocean from the western coast of southern India from Bombay South to Sri Lanka also Madagascar and Mauritius, the recent information is also caught in Taiwan (Hu et al. 2015), Hong Kong (Leung 1997), and the Philippines (Quilang et al. 2011). The result strongly supported that *S. jussieu* is more widely distributed than we have thought and the large-scale survey should be made to know the spatiotemporal distribution of four *Sardinella* species in Indonesia. We here reported the full-length mitochondrial genome sequence of *S. jussieu* collected from Java island, which would provide the




Fig. 5 a Phylogenetic tree of mitochondrial genome of four species belonging to *Sardinella*. The phylogenetic tree was constructed using molecular evolutionary genetic analysis ver.6.0 (MEGA 6, MEGA Inc. Englewood, NJ), program with the minimum evolution algorithm, the evolutionary distances were computed using Kimura 2-Parameter method and **b** Phylogenetic tree of CO1 sequences of 18 species belonging to genus *Sardinella*. The phylogenetic tree was constructed using molecular evolutionary genetic analysis ver.6.0 (MEGA 6, MEGA Inc. Englewood, NJ), program with the minimum evolution algorithm, the evolutionary distances were computed using Kimura 2-Parameter method

important information for the scientific management of *Sardinella* species in Indonesia. We expect that more *Sardinella* species may exist in Java island and more information about the mitochondrial genome of the other unreported *Sardinella* species such as *S. gibbosa* would be a useful information for the molecular biological tools to discriminate different *Sardinella* species in Indonesia.

Conclusion

This study determined the complete mitochondrial DNA (mtDNA) sequence of *S. jussieu* in Java Island,

Indonesia, for the first time. The mtDNA sequence is 16.695 bp in length and comprises the typical set of 2 rRNAs, 22 tRNA genes, 13 protein-coding genes, and putative control region. Mitochondrial genome structure of *S. jussieu* was identical to those in other *Sardinella* genus. Phylogenetic analysis using full mitochondrial genome exhibits that *S. jussieu* were most closely related to *S. albella*. However, comparison in the COI region showed that relationship between *S. albella* and *S. gibbosa* was ambiguous and determination of the complete mitochondrial DNA sequence of

S. gibbosa is required for the better understanding of evolutionary relationship between *S. jussieu* and those species. Those information would provide the basic information for the scientific management of *Sardinella* species in Indonesia.

Abbreviations

COI region: Cytochrome c oxidase subunit 1 region; Cyt-B: Cytochrome B subunit; mtDNA: Mitochondrial DNA; ND4: NADH dehydrogenase subunit 4; ND5: NADH dehydrogenase subunit 5; ND6: NADH dehydrogenase subunit 6; NGS: Next-generation sequencing

Acknowledgements

All authors thank the funding of grant from the Pukyong National University, and authors also thank the Jakarta fisheries University students who helped for collecting the fish samples.

Funding

This work was supported by a grant from the Pukyong National University in 2016.

Availability of data and materials

The dataset(s) supporting the conclusions of this article is(are) included within the article.

Authors' contributions

SPS carried out the whole process, participated in the whole experiment, and drafted the manuscript. SA added and drafted the final phylogenetic analysis. HWK participated in the design of the study and edited the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Interdisciplinary Program of Biomedical, Mechanical and Electrical Engineering, Pukyong National University, Busan 48513, Republic of Korea. ²Department of Marine Biology, Pukyong National University, Busan 48513, Republic of Korea. ³Aquaculture Technology Study Program, Jakarta Fisheries University, Jl. AUP Pasar Minggu Jakarta Selatan, Jakarta 12520, Indonesia. ⁴Fisheries and Marine Faculty, C Campus Jl. Mulyorejo, Surabaya 60115, Indonesia. ⁵Universitas Airlangga, Surabaya, East Java, Indonesia.

Received: 12 July 2017 Accepted: 28 September 2017

Published online: 23 October 2017

References

- Begg GA, Waldman JR. An holistic approach to fish stock identification. *Fish Res.* 1999;43(1):35–44.
- Bräger Z, Moritz T. A scale atlas for common Mediterranean teleost fishes. *Vertebr Zool.* 2016;66
- Ditty JG, Houde ED, Shaw RF. Egg and larval development of Spanish sardine, *Sardinella aurita* (Family Clupeidae), with a synopsis of characters to identify clupeid larvae from the northern Gulf of Mexico. *Bull Mar Sci.* 1994;54(2):367–80.
- Homayuni H, Marjani M, Mousavi-Sabet H. Descriptive key to the otoliths of three *Sardinella* species (Pisces, Clupeidae) from the northern Oman Sea. *AAFL Bioflux.* 2013;6(3):211–21.

- Hu W, et al. Study on fish life history traits and variation in the Taiwan Strait and its adjacent waters. *Acta Oceanol Sin.* 2015;34(2):45.
- Kneibelsberger T, Stöger I. DNA extraction, preservation, and amplification. In: *DNA barcodes: methods and protocols*; 2012. p. 311–38.
- Lavoué S, et al. Phylogenetic relationships among anchovies, sardines, herrings and their relatives (Clupeiformes), inferred from whole mitogenome sequences. *Mol Phylogenet Evol.* 2007;43(3):1096–105.
- Leung A. The epibenthic ichthyofauna of Tolo Harbour and Hong Kong's northeastern waters: a long term record of change. In: *Proceedings of the Eighth International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China*, Hong Kong University Press, Hong Kong; 1997.
- MMAF. Capture fisheries statistics of Indonesia, 2011. Jakarta: Directorate General of Capture Fisheries; 2012. p. 190.
- Palumbi S, et al. The simple fool's guide to PCR, version 2.0, privately published document compiled by S. Palumbi. Dept. Honolulu: Zoology, Univ. Hawaii; 1991. p. 96822.
- Quilang JP, et al. DNA barcoding of the Philippine endemic freshwater sardine *Sardinella tawilis* (Clupeiformes: Clupeidae) and its marine relatives. *Philipp Agric Sci.* 2011;94(3)
- Sivakumaran K, Manickasundaram M, Ramaiyan V. Problems of identification among species of *Sardinella*. In: *CMFRI Bulletin National Symposium on Research and Development in Marine Fisheries Sessions I & II 1987*. Kochi: CMFRI; 1987.
- Stern N, Rinkevich B, Goren M. Integrative approach revises the frequently misidentified species of *Sardinella* (Clupeidae) of the Indo-West Pacific Ocean. *J Fish Biol.* 2016;89(5):2282–305.
- Strauss RE, Bond CE. Taxonomic methods: morphology. *Methods for fish biology*. Bethesda: American Fisheries Society; 1990. p. 109–40.
- Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol.* 1994;3(5):294–9.
- Ward RD, et al. DNA barcoding Australia's fish species. *Philos Trans R Soc Lond B Biol Sci.* 2005;360(1462):1847–57.
- Whitehead P. *FAO species catalogue vol. 7. Clupeoid fishes of the world (suborder Clupeoidei): an annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, shads, anchovies and wolf-herrings*. Rome: Food and Agriculture Organization of the United Nations; 1985.
- Willette D, Santos M, Aragon M. First report of the Taiwan sardinella *Sardinella huaiensis* (Clupeiformes: Clupeidae) in the Philippines. *J Fish Biol.* 2011;79(7): 2087–94.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

