

# EVOLUTION OF ORBIT AND ROTATION OF A PSEUDO-SYNCHRONOUS BINARY SYSTEM ON THE MAIN SEQUENCE

LIN-SEN LI

School of Physics, Northeast Normal University, Changchun 130024, China; [lils653@nenu.edu.cn](mailto:lils653@nenu.edu.cn)

Received August 5, 2017; accepted January 8, 2018

**Abstract:** We study the pseudo-synchronous orbital motion of a binary system on the main sequence. The equations of the pseudo-synchronous orbit are derived up to  $O(e^4)$  where  $e$  is the eccentricity of the orbit. We integrate the equations to present their solutions. The theoretical results are applied to the evolution of the orbit and spin of the binary star Y Cygni, which has a current eccentricity of  $e_0 = 0.142$ . We tabulate our numerical results for the evolution of the orbit and spin per century. The numerical results for the semi-major axes and rotational angular velocities in the evolutionary time scales of three stages (synchronization, circularization, and collapse time scale) are also tabulated. Synchronization is achieved in about  $5 \times 10^3$  years followed by circularization lasting about  $1 \times 10^5$  years before decaying in  $2 \times 10^5$  years.

**Key words:** stars: binaries — stars: kinematics and dynamics — stars: individual: Y Cygni

## 1. INTRODUCTION

In a close detached binary system, tidal interaction couples the spin and orbital angular momenta of the component stars. The orbit and spin of a binary system evolve from a non-synchronous orbit to a synchronous one gradually under the influence of tidal friction. It is important and meaningful to study the synchronization process of the orbit and spin. The evolution of the orbit and spin is always from non-synchronization to synchronization, where the synchronized orbit is the final destination of the evolution. This is an inevitable tendency of the evolution from the younger binary stars to old binary stars.

Generally speaking, the synchronization of binary stars requires that the rotational angular velocity  $\Omega$  equals the orbital angular velocity  $\omega$ . However, strictly speaking, truly synchronous binary stars are found only very rarely among binary systems. What is observed instead is that each component star settles in a state of pseudo-synchronization  $\Omega \neq \omega$ . All binary systems appear to pass through a phase of pseudo-synchronous orbital motion. Therefore, pseudo-synchronous orbits are quite common in binary systems. Hence, it is very important to study the evolution of orbital and spin periods of binary stars.

Zahn (1977, 1978, 1989) and Zahn & Bouchet (1989) established the set of equations describing the orbital evolution of non-synchronization of binary systems. However, they did not investigate the orbit and spin of the pseudo-synchronization. Hut (1980, 1981) defines pseudo-synchronization and showed that the spin period converges to a value slightly smaller than the orbital period by an order of  $e^2$ .

Li (2012, 2013, 2015) studied the orbit and spin synchronization of binary stars using analytical and nu-

merical solutions, but this study does not cover the pseudo-synchronization process. We investigate the orbit and spin of pseudo-synchronization of binary stars and obtain the solution of the set of the equations of pseudo-synchronization.

## 2. DEFINITION OF PSEUDO-SYNCHRONIZATION

Zahn (1977) wrote down the equations for the orbit and spin of a non-synchronized binary system and in Zahn (1978) also revised the equations for the time rate change of eccentricity  $de/dt$  and spin rate  $d(I\Omega)/dt$  of the primary star as follows.

$$\frac{1}{a} \frac{da}{dt} = -12 \frac{k_2}{t_f} q(1+q) \left(\frac{R}{a}\right)^8 \left[ \left(1 - \frac{\Omega}{\omega}\right) + e^2 \left(23 - \frac{27\Omega}{2\omega}\right) + O(e^4) \right] \quad (1)$$

$$\frac{1}{e} \frac{de}{dt} = -3 \frac{k_2}{t_f} q(1+q) \left(\frac{R}{a}\right)^8 \left[ \left(18 - \frac{11\Omega}{\omega}\right) + O(e^2) \right] \quad (2)$$

$$\frac{d(I\Omega)}{dt} = 6 \frac{k_2}{t_f} q^2 M R^2 \left(\frac{R}{a}\right)^6 [(\omega - \Omega) + e^2 \left(\frac{27\omega}{2} - \frac{15\Omega}{2}\right) + O(e^4)]. \quad (3)$$

Here,  $M$ ,  $R$  and  $I$  denote the mass, radius, and moment of inertia of the primary star and  $M_2$  denotes the mass of the secondary star. The parameter  $q = M_2/M$  is the mass ratio of the primary and secondary stars. The parameters  $a, e, \omega$  denote semi-major axis, eccentricity and mean motion. We denote the apsidal motion by  $k_2$  and the friction time by  $t_f$  introduced by Zahn &

**Table 1**  
Orbital elements and derived parameters for Y Cygni

| Y Cyg<br>Orbit elements | P(d)       | a( $R_\odot$ )          | M( $M_\odot$ ) | q        | R( $R_\odot$ ) | e     | log L( $L_\odot$ ) | $k_2$    | V(km s $^{-1}$ ) |
|-------------------------|------------|-------------------------|----------------|----------|----------------|-------|--------------------|----------|------------------|
|                         | 2.996      | 28.49                   | 17.57          | 0.97     | 5.93           | 0.142 | 4.653              | 0.005745 | 146              |
| Y Cyg<br>Derived values | $t_F$ (yr) | $k_2/t_F$ (yr $^{-1}$ ) | $MR^2/I$       | $\omega$ | $\Omega$       | $N$   |                    |          |                  |
|                         | 0.1038     | 0.0553                  | 8.2508         | 2.0965   | 3.5735         | 49.59 |                    |          |                  |

$P$ ,  $a$ ,  $M$ ,  $q$ ,  $R$ ,  $L$  and  $e$  are from Simon et al. (1994) and Ibanoglu & Soyudugan (2006). Data for  $k_2$  are from Peraiah (1965). The value of  $t_F$  is calculated using Equation (4).

Bouchet (1989)

$$t_F = \left( \frac{MR^2}{L} \right)^{1/3}. \quad (4)$$

Here,  $L$  is the stellar luminosity.

Hut (1981) gives the definition of the pseudo-synchronization, describing the phenomenon of near synchronization of revolution and rotation around periastron, where the tidal interaction is the strongest. Pseudo-synchronism is therefore important in binary systems with substantial eccentricity. As Hut (1981) discusses, Mercury apparently achieved pseudo-synchronism with the Sun where its spin rate is near its orbital revolution near perihelion.

According to the definition by Hut (1981),  $\Omega = \Omega_{ps}$  if  $d\Omega/dt = 0$  in Equation (3). It can be obtained for definition of the condition of the pseudo-synchronization, from which  $(\omega - \Omega) + e^2 \left( \frac{27\omega}{2} - \frac{15\Omega}{2} \right) = 0$  and  $6(k_2/t_f)q^2 MR^2 (R/a)^6 \neq 0$ . Hence, the condition of the pseudo-synchronization can be written as

$$\frac{\Omega}{\omega} = \frac{2 + 27e^2}{2 + 15e^2}. \quad (5)$$

This shows that the departure from strict synchronism ( $\Omega = \omega$ ) is of order  $e^2$ .

### 3. PSEUDO-SYNCHRONOUS EQUATIONS AND THEIR SOLUTIONS

Substituting the condition for the pseudo-synchronization (5) into Equations (1) and (2), we obtain the following two equations for pseudosynchronization

$$\frac{1}{a} \frac{da}{dt} = -6K_1 e^2 \left( \frac{14 - 39e^2}{2 + 15e^2} \right), \quad (6)$$

and

$$\frac{1}{e} \frac{de}{dt} = -3K_1 \left( \frac{14 - 27e^2}{2 + 15e^2} \right). \quad (7)$$

Here, the parameter  $K_1$  is given by

$$K_1 = (k_2/t_F)q(1+q)(R/a)^8. \quad (8)$$

With the presence of the term  $O(e^4)$ , Equations (6) and (7) are valid up to the order of  $e^3$ . As we will present in Section 4, we will apply our theoretical model to the binary system Y Cyg, for which the eccentricity  $e = 0.142$ ,  $e^4 = 0.0004$  is small enough. Hence Equations (6) and (7) may be neglected for the order  $O(e^4)$ .

From Equations (6) and (7), we have

$$\frac{da}{de} = \frac{da}{dt} \frac{dt}{de} = 2ae \left( \frac{14 - 39e^2}{14 - 27e^2} \right), \quad (9)$$

leading to

$$\begin{aligned} \frac{da}{a} &= \left( \frac{14 - 39e^2}{14 - 27e^2} \right) d(e^2), \\ &= \frac{1}{14} [14 - 12e^2 + O(e^4)] d(e^2). \end{aligned} \quad (10)$$

Integrating this equation, we have

$$\ln \left( \frac{a}{a_0} \right) = \int_{e_0}^e \left( 1 - \frac{6}{7}e^2 \right) d(e^2) \quad (11)$$

which leads to

$$\ln(a/a_0) = (e^2 - e_0^2) - \frac{6}{7}(e^4 - e_0^4) + O(e^6). \quad (12)$$

Neglecting terms of order  $e^4$ , we obtain

$$a = a_0 \exp(e^2 - e_0^2). \quad (13)$$

Next, we solve Equation (7) in order to obtain variation of the eccentricity with time. Writing Equation (7) in the following form

$$\frac{1}{e} \frac{de}{dt} = -3K_2 \left( \frac{14 - 27e^2}{2 + 15e^2} \right) a^{-8}, \quad (14)$$

where the parameter  $K_2$  is defined by

$$K_2 = (k_2/t_F)q(1+q)R^8. \quad (15)$$

Substituting for Equation (10) into Equation (11), it can be written as

$$\frac{1}{e} \frac{de}{dt} = -\frac{3k_2}{a_0^8} \left( \frac{14 - 27e^2}{2 + 15e^2} \right) e^{-8(e^2 - e_0^2)} \quad (16)$$

which can be rearranged to

$$\left( \frac{2 + 15e^2}{14 - 27e^2} \right) e^{8e^2} \frac{de}{e} = -3K_2 a_0^8 e^{8e_0^2} dt. \quad (17)$$

Neglecting terms of order  $e^4$  and higher, one can obtain

$$\left[ \left( \frac{31}{28} + \frac{27}{196} \right) + \frac{1}{14e^2} \right] de^2 = -3K_2 a_0^{-8} \exp(8e_0^2) dt. \quad (18)$$

**Table 2**

Numerical results for the change of the orbital elements and spin of Y Cyg during 100 years

| Orbital Elements | $a/a_0$ | $a(R_\odot)$ | $\delta a(R_\odot)$ | $e/e_0$ | $e$    | $\delta e$ | $\delta\Omega$ |
|------------------|---------|--------------|---------------------|---------|--------|------------|----------------|
| Y Cyg            | 0.9999  | 28.487       | -0.0004             | 0.9996  | 0.1419 | -0.0001    | 0.00005        |

Integration from  $t_0$  and  $e_0$  to  $t$  and  $e$  yields

$$\begin{aligned} & \left( \frac{31}{28} + \frac{27}{196} \right) (e^2 - e_0^2) + \frac{1}{14e^2} N(e) (e^2 - e_0^2) de^2 \\ & = -3K_2 a_0^{-8} \exp(8e_0^2) dt, \end{aligned} \quad (19)$$

where the expression  $N(e)$  is

$$N = (\ln e^2 - \ln e_0^2) / (e^2 - e_0^2). \quad (20)$$

Substitution of the expression of  $K_2$  leads to

$$e^2 - e_0^2 = -\frac{3k_2 q(1+q)(R/a_0)^8 \exp(8e_0^2)(t-t_0)}{t_f \left( \frac{N}{7} + \frac{31}{28} + \frac{27}{196} \right) e_0^2}, \quad (21)$$

from which

$$e = e_0 \left[ 1 - \frac{3k_2 q(1+q)(R/a_0)^8 \exp(8e_0^2)(t-t_0)}{t_f \left( \frac{N}{7} + \frac{31}{28} + \frac{27}{196} \right) e_0^2} \right]^{1/2}, \quad (22)$$

Substitution into Equation (13) yields

$$a = a_0 \exp \left[ -\frac{3k_2 q(1+q)(R/a_0)^8 \exp(8e_0^2)(t-t_0)}{t_f \left( \frac{N}{7} + \frac{31}{28} + \frac{27}{196} \right)} \right]. \quad (23)$$

From Equation (5) with  $e = e_0$  we may write

$$\delta\Omega = \left( \frac{2 + 27e_0^2}{2 + 15e_0^2} \right) \delta\omega. \quad (24)$$

Combining it with the Kepler's third law  $\delta\omega = -\frac{3}{2} \left( \frac{\omega}{a} \right) da$ , we obtain

$$\delta\Omega = -\frac{3}{2} \left( \frac{2 + 27e_0^2}{2 + 15e_0^2} \right) \left( \frac{\omega}{a} \right) \delta a. \quad (25)$$

Zahn (1977) defined the circulatization time  $t_{\text{cir}}$  by means of his non-revised Equation (2) for  $de/dt$  as

$$t_{\text{cir}}^{-1} = -e^{-1} \frac{de}{dt} = \frac{63 k_2}{4 t_f} q(1+q)(R/a)^8 \text{ yr}^{-1} \quad (26)$$

assuming that  $\Omega = \omega$ . However, in this work, we use the revised equation for  $de/dt$  with  $\Omega = \omega$ , in which case the circularization time should be written as

$$t_{\text{cir}}^{-1} = 21 \frac{k_2}{t_f} q(1+q)(R/a)^8 \text{ yr}^{-1}. \quad (27)$$

According to Equation (3), the synchronization time  $t_{\text{syn}}$  becomes

$$\frac{1}{t_{\text{syn}}} = -\frac{1}{\Omega - \omega} \frac{d\Omega}{dt} = 6 \frac{k_2}{t_f} q^2 \left( \frac{MR^2}{I} \right) (R/a)^6 \text{ yr}^{-1} \quad (28)$$

In turn, the orbital decay time  $t_{\text{decay}}$  or the collapse time of the system can be written as

$$\frac{1}{t_{\text{decay}}} = -\frac{1}{a} \frac{da}{dt} = 12 \frac{k_2}{t_f} q(1+q)(R/a)^8 \quad (29)$$

#### 4. PSEUDO-SYNCHRONIZATION OF ORBIT AND SPIN IN Y CYGNI

We choose the binary system Y Cygni as our exemplary model for pseudo-synchronization. The spectral type of Y Cyg is traditionally given as B0 V in the literature, but has been found to be earlier, O9.8 V (Simon et al. 1994). Y Cyg is an eclipsing binary and is also very well known for its apsidal motion. According to Giuricin (1984), the rotational velocity  $V = 146 \text{ km s}^{-1}$  and the estimated synchronized velocity and  $V_k = 88 \text{ km s}^{-1}$  for the binary star Y Cyg, which shows that this binary system is not a synchronous binary system. The orbital elements and the deduced parameters are listed in Table 1.

Noting that  $e \simeq e_0$ , we use l'Hospital's rule to approximate Equation (20) by

$$N(e) = \frac{\ln e^2 - \ln e_0^2}{e^2 - e_0^2} \simeq \frac{1}{e_0^2}. \quad (30)$$

For Y Cyg, we have  $e_0 = 0.142$ , which, in turn, leads to  $N \simeq e_0^{-2} = 49.6$ . The use of l'Hospital's rule is justified because the variation  $\delta e$  is very small.

Substituting these values for data of  $q, e, R, N$  while taking  $t - t_0 = 100 \text{ yr}$  into Equations (16)–(18) and (19)–(20) and (22), we obtain the change of the orbit and spin per century for Y Cyg, which is summarized in Table 2.

We estimate the changes of the orbital semi-major axis and spin of Y Cyg when the binary system achieves synchronization, circularization, and collapse. Substituting these values of data in Table 1 into Equations 28, 27 and 29, we obtain the values of  $t_{\text{cir}}$ ,  $t_{\text{syn}}$  and  $t_{\text{decay}}$ .

$$\begin{aligned} t_{\text{syn}} &= 4.774 \times 10^5 \text{ yr}, \\ t_{\text{cir}} &= 1.279 \times 10^5 \text{ yr} \\ t_{\text{decay}} &= 2.239 \times 10^5 \text{ yr} \end{aligned} \quad (31)$$

Substituting Equation (31) into Equations (23), (24), and (25), we obtain the values for the three cases of evolutionary stages of pseudo-synchronization of Y Cyg listed in Table 3.

#### 5. CONCLUSION

It can be seen from Table 2 that the orbital parameters and spin exhibit very slow changes for 100 years due to the action of the tidal friction for Y Cyg. We may also find from Table 3 that Y Cyg achieves or enters the synchronization stage firstly, and then, circularization stage and finally enter the orbital collapse stage.

From Table 3, we may assert that in the system of Y Cyg the orbital semi-major axis,  $a$ , decreases with age monotonically and the spin rate,  $\Omega$ , increases with age

**Table 3**  
Evolution of semi-major axis and spin of Y Cyg

| Stage | $t$<br>(yr)         | $a/a_0$ | $a$<br>( $R_\odot$ ) | $\delta a$<br>( $R_\odot$ ) | $\delta\Omega$<br>(rad/d) |
|-------|---------------------|---------|----------------------|-----------------------------|---------------------------|
| A     | $4.774 \times 10^3$ | 0.9991  | 28.4643              | -0.0256                     | 0.0031                    |
| B     | $1.279 \times 10^5$ | 0.9811  | 27.9515              | -0.05384                    | 0.0065                    |
| C     | $2.239 \times 10^5$ | 0.9669  | 26.852               | -0.9430                     | 0.1998                    |

Numerical results for the three evolutionary stages of the orbital semi-major axis and spin of Y Cyg. The stages A, B and C stand for synchronization, circularization and collapse, respectively.

continuously in each stage. In the synchronous stage the decrease is large for  $\delta a_{\text{syn}}$ , and the increase is large for  $\delta\Omega_{\text{syn}}$ . In the circularization stage, we find that  $\delta a_{\text{cir}} < \delta a_{\text{syn}}$  and  $\delta\Omega_{\text{syn}} > \delta\Omega_{\text{cir}}$ . However, in the orbital collapse stage,  $\delta a < \delta a_{\text{cir}} < \delta a_{\text{syn}}$  and  $\delta\Omega > \delta\Omega_{\text{cir}} > \delta\Omega_{\text{syn}}$ .

It should be noted that Equations (1) and (2) are valid up to second order of eccentricity  $e$  and that higher order terms  $O(e^4)$  are neglected. Hence these equations should be used for binary systems with a small eccentricity, and may be inadequate to those with a large eccentricity. For the eccentricity  $e = 0.142$  of the eclipsing binary system Y Cyg, we have  $e^4 = 0.0004$ , which is sufficiently small validating our use of Equations (1) and (2).

#### REFERENCES

Giuricin, G., Mardirossian, F., & Mezzetti, M. 1984, Synchronization in Eclipsing Binary Stars, *A&A*, 131, 152

- Hut, P. 1980, Stability of Tidal Equilibrium, *A&A*, 92, 167  
Hut, P. 1981, Tidal Evolution in Close Binary Systems, *A&A*, 99, 126  
Ibanoglu, C., Soydugan, F., Soydugan, E., & Dervisoglu, A. 2006. Angular Momentum Evolution of Algol Binaries, *MNRAS*, 373, 435  
Li, L.-S. 2012, Orbit and Spin Evolution of Synchronous Binary Stars on the Main Sequence, *Res. Astron. Astrophys.*, 12, 1673  
Li, L.-S. 2013, Numerical Solution and Evolutionary Treads of the Orbit and Rotation of a Synchronous Binary Star System at the Orbital Circularization Time, *Astron. Res & Tech.* 10 (3), 249  
Li, L.-S. 2015, Orbit and Spin Evolution of Synchronous Binary Stars on the Main Sequence (A Theoretical Improvement to the Analytical Method), *Res. Astron. Astrophys.*, 15, 1695  
Peraiah, A. 1965, Rotation of the Components in Close Binary Systems, *ZA*, 62, 48  
Simon, K. P., Sturm, E., & Fielder, A. 1994, Spectroscopic Analysis of Hot Binaries. 2: The Components of Y Cygni, *A&A*, 292, 507  
Zahn, J. P. 1977, Tidal Friction in Close Binary Stars, *A&A*, 57, 383  
Zahn, J. P. 1978, Erratum: Tidal Friction in Close Binary Stars, *A&A*, 67, 162  
Zahn, J. P. 1989, Tidal Evolution of Close Binary Stars. I – Revisiting the Theory of the Equilibrium Tide, *A&A*, 220, 112  
Zahn, J. P., & Bouchet, L. 1989, Tidal Evolution of Close Binary Stars. II – Orbital Circularization of Late-Type Binaries, *A&A*, 223, 112