

# Will Platelet Rich Plasma Open New Era for Treatment of Rotator Cuff Tear?

Sungwook Choi<sup>✉</sup>

Department of Orthopaedic Surgery, Jeju National University School of Medicine, Jeju, Korea

About 30% to 70% of chronic shoulder pain is developed by rotator cuff tear, impairing daily living activities, and quality of life.<sup>1-3)</sup> It is reported that progression of tears is not only characterized by cellular but also extracellular matrix changes including reduced cell number, decreased vascularization, poor matrix organization, changed gene expression and alterations in proteoglycans and collagens, which play an important role in rotator cuff tear pathology.<sup>4-6)</sup> Some of the up-to-date literatures point out that platelet rich plasma (PRP) provides potential healing environment for tendon to bone attachment.<sup>7,8)</sup> However, there has been no comparative study of PRP effect on tenocytes collected from the rotator cuff between normal and degenerative torn tendon.

It is well known by many other literatures state that the cell proliferation, or density of tenocytes are much more higher in teared rotator cuff than normal rotator cuff.<sup>4,9)</sup> Moreover, Anitua et al.<sup>10,11)</sup> reported that 20% PRP releasate (vol/vol) increased cell proliferation and the production of vascular endothelial growth factor and hepatocyte growth factor in human healthy hamstring tendon. It is proven in the research by Jo et al.<sup>12)</sup> that PRP activated calcium with or without thrombin significantly stimulated the proliferation of tenocytes from human rotator cuff tendons with degenerative tears in a dose-dependent manner.

The mechanism of proteoglycan transformation is also found to be critical in pathology of rotator cuff tear. Lo et al.<sup>13)</sup> revealed changes in proteoglycans; aggrecan increased while decorin decreased in the ruptured rotator cuff tendon when compared with normal cadaveric controls. Elevated level of proteoglycan, tenascin-C and fibronectin, in ruptured rotator cuff was reported.<sup>14,15)</sup>

The enhancement of gene expression of makers for collagen type I, and decrease of collagen type III level is carried out by

PRP, and resultant positive collagen I/III ratio was achieved in equine tendon explants.<sup>16)</sup> de Mos et al.<sup>17)</sup> claimed that when PRP was injected to normal tendon of 13 to 15 aged children, number of cell, and total level of collagen were significantly increased, but decreased the gene expressions of type I and III collagen without affecting the III/I ratio was identified.

The effect of PRP was quiet different from that of de Mos et al.,<sup>17)</sup> since the gene expression of type I and III collagen in both tenocytes from normal and degenerative tendon were increased, and Yoon et al.<sup>18)</sup> showed that there were no statistically significant differences between normal and degenerative tendon. The gene expression of type III collagen was only significantly increased in degenerative tendon. However, the fold-change of type I collagen in degenerative tendon was higher than in normal tendon. Although there were no statistical significances, this increased rate can be considered to reflect a greater impact of PRP on the degenerative tendon.

The retear rate after rotator cuff repair varies depending on tear size of tendon, number of tendons involved, and the age of patients. It is described that rates of retear are up to 80% to 90% exist in the radiology literature and are as high as 57% in the orthopaedic literature.<sup>19-21)</sup> However, according to the previously published studies, when PRP technique is added, decrease of the retear rate of rotator cuff is expected.

PRP, which promotes cellular proliferation and regeneration, is being utilized in many field of medical science and related researches are in full swing. PRP expedites tenocyte proliferation in both normal and degeneratively torn rotator cuff and facilitates tendon matrix synthesis. Not only PRP has a positive effect on cell proliferation, but also elevated matrix gene expression and synthesis in tenocytes from degenerative torn tendon than tenocytes from normal tendon. Applying properly administrated PRP

<sup>✉</sup>Correspondence to: Sungwook Choi

Department of Orthopaedic Surgery, Jeju National University Hospital, Jeju National University School of Medicine, 15 Aran 13-gil, Jeju 63241, Korea

Tel: +82-64-717-1690, Fax: +82-64-717-1131, E-mail: swchoi1115@gmail.com

Editorial does not need an IRB approval.

Financial support: None. Conflict of interests: None.

in treating rotator cuff tear will promise excellent functional outcome.

## References

1. Bongers PM. The cost of shoulder pain at work. *BMJ*. 2001;322(7278):64-5.
2. Urwin M, Symmons D, Allison T, et al. Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at different anatomical sites, and the relation to social deprivation. *Ann Rheum Dis*. 1998;57(11):649-55.
3. Jo CH, Shin WH, Park JW, Shin JS, Kim JE. Degree of tendon degeneration and stage of rotator cuff disease. *Knee Surg Sports Traumatol Arthrosc*. 2017;25(7):2100-8.
4. Matthews TJ, Hand GC, Rees JL, Athanasou NA, Carr AJ. Pathology of the torn rotator cuff tendon. Reduction in potential for repair as tear size increases. *J Bone Joint Surg Br*. 2006;88(4):489-95.
5. Chaudhury S, Carr AJ. Lessons we can learn from gene expression patterns in rotator cuff tears and tendinopathies. *J Shoulder Elbow Surg*. 2012;21(2):191-9.
6. Riley GP. Gene expression and matrix turnover in overused and damaged tendons. *Scand J Med Sci Sports*. 2005;15(4):241-51.
7. Jo CH, Shin JS, Shin WH, Lee SY, Yoon KS, Shin S. Platelet-rich plasma for arthroscopic repair of medium to large rotator cuff tears: a randomized controlled trial. *Am J Sports Med*. 2015;43(9):2102-10.
8. Kim SJ, Lee SM, Kim JE, Kim SH, Jung Y. Effect of platelet-rich plasma with self-assembled peptide on the rotator cuff tear model in rat. *J Tissue Eng Regen Med*. 2017;11(1):77-85.
9. Lundgreen K, Lian OB, Engebretsen L, Scott A. Tenocyte apoptosis in the torn rotator cuff: a primary or secondary pathological event? *Br J Sports Med*. 2011;45(13):1035-9.
10. Anitua E, Sánchez M, Orive G, Andía I. The potential impact of the preparation rich in growth factors (PRGF) in different medical fields. *Biomaterials*. 2007;28(31):4551-60.
11. Anitua E, Sánchez M, Zalduendo MM, et al. Fibroblastic response to treatment with different preparations rich in growth factors. *Cell Prolif*. 2009;42(2):162-70.
12. Jo CH, Kim JE, Yoon KS, Shin S. Platelet-rich plasma stimulates cell proliferation and enhances matrix gene expression and synthesis in tenocytes from human rotator cuff tendons with degenerative tears. *Am J Sports Med*. 2012;40(5):1035-45.
13. Lo IK, Boorman R, Marchuk L, Hollinshead R, Hart DA, Frank CB. Matrix molecule mRNA levels in the bursa and rotator cuff of patients with full-thickness rotator cuff tears. *Arthroscopy*. 2005;21(6):645-51.
14. Riley GP, Harrall RL, Cawston TE, Hazleman BL, Mackie EJ. Tenascin-C and human tendon degeneration. *Am J Pathol*. 1996;149(3):933-43.
15. Tillander B, Franzén L, Norlin R. Fibronectin, MMP-1 and histologic changes in rotator cuff disease. *J Orthop Res*. 2002;20(6):1358-64.
16. McCarrel T, Fortier L. Temporal growth factor release from platelet-rich plasma, trehalose lyophilized platelets, and bone marrow aspirate and their effect on tendon and ligament gene expression. *J Orthop Res*. 2009;27(8):1033-42.
17. de Mos M, van der Windt AE, Jahr H, et al. Can platelet-rich plasma enhance tendon repair? A cell culture study. *Am J Sports Med*. 2008;36(6):1171-8.
18. Yoon JY, Lee SY, Shin S, Yoon KS, Jo CH. Comparative analysis of platelet-rich plasma effect on tenocytes from normal human rotator cuff tendon and human rotator cuff tendon with degenerative tears. *Clin Shoulder Elbow*. 2018;21(1):3-14.
19. Charoussat C, Bellaïche L, Kalra K, Petrover D. Arthroscopic repair of full-thickness rotator cuff tears: is there tendon healing in patients aged 65 years or older? *Arthroscopy*. 2010;26(3):302-9.
20. Liem D, Bartl C, Lichtenberg S, Magosch P, Habermeyer P. Clinical outcome and tendon integrity of arthroscopic versus mini-open supraspinatus tendon repair: a magnetic resonance imaging-controlled matched-pair analysis. *Arthroscopy*. 2007;23(5):514-21.
21. Liem D, Lichtenberg S, Magosch P, Habermeyer P. Magnetic resonance imaging of arthroscopic supraspinatus tendon repair. *J Bone Joint Surg Am*. 2007;89(8):1770-6.