

Toxicol. Res. Vol. 34, No. 1, p. 83 (2018) https://doi.org/10.5487/TR.2018.34.1.083

Open Access

Erratum to "Anti-Inflammatory Effect of 3-Bromo-4,5-Dihydroxy-benzaldehyde, a Component of *Polysiphonia morrowii*, *In Vivo* and *In Vitro*" [Toxicol. Res. 33 (2017) 325-332]

Na-Jin Kang¹, Sang-Chul Han¹, Hyun-Jae Kang¹, Geum Ko¹, Weon-Jong Yoon², Hee-Kyoung Kang¹ and Eun-Sook Yoo¹

On page 327 and 328, the authors found labeling errors in Fig. 2B and Fig. 3B. Fig. 2B is originally result for "IgE levels" but was erroneously expressed as "ear thickness". Therefore author wanted to change from Fig. 2B ear thickness results to IgE levels and from Fig. 3B to corrected "ear thickness" results.




Fig. 2. 3-Bromo-4,5-dihydroxybenzaldehyde suppresses the expression of serum IgE.




Fig. 3. 3-Bromo-4,5-dihydroxybenzaldehyde suppresses experimental atopic dermatitis.

Correspondence to: Eun-Sook Yoo, Department of Medicine, School of Medicine, Jeju National University, Jeju 63243, Korea E-mail: eunsyoo@jejunu.ac.kr

DOI of original article: http://dx.doi.org/10.5487/TR.2017.33.4.325

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

¹Department of Medicine, School of Medicine, Jeju National University, Jeju, Korea

²Jeju Biodiversity Research Institute (JBRI), Jeju Technopark (JTP), Jeju, Korea