

DERIVATIONS OF A NON-ASSOCIATIVE GROWING ALGEBRA

SEUL HEE CHOI

Abstract. There are various papers on finding all the derivations of a non-associative algebra and an anti-symmetrized algebra. We find all the derivations of a growing algebra in the paper. The dimension of derivations of the growing algebra is one and every derivation of the growing algebra is outer. We show that there is a class of purely outer algebras in this work.

1. Introduction

Let \mathbb{N} be the set of all non-negative integers and \mathbb{Z} be the set of all integers. Let \mathbb{N}^+ be the set of all positive integers. Let \mathbb{F} be a field of characteristic zero and \mathbb{F}^\bullet the set of all non-zero elements in \mathbb{F} . Throughout the paper, we will assume that e is not the element of the field \mathbb{F} . For $n, t \in \mathbb{N}$, throughout the paper, m denotes a non-negative integer such that $m \leq n + t$. For fixed integers, i_1, \dots, i_m and for given irreducible polynomials $f_1, \dots, f_m \in \mathbb{F}[x_1, \dots, x_{n+t}]$, define $[f_1^{i_1}, \dots, f_m^{i_m}]$ as the set $Poly_m = P_m = \{f_1^{i_1} \cdots f_m^{i_m}, f_1^{i_1} \cdots f_{m-1}^{i_{m-1}}, \dots, f_2^{i_2} \cdots f_m^{i_m}, \dots, f_1^{i_1}, \dots, f_m^{i_m}\}$. For any subset P of P_m , define the \mathbb{F} -algebra $\mathbb{F}[e^{\pm[P]}, n, t] := \mathbb{F}[e^{\pm[P]}, x_1^{\pm 1}, \dots, x_n^{\pm 1}, x_{n+1}, \dots, x_{n+t}]$, which is spanned by

$$\mathbf{B} = \{e^{a_1 f_1} \cdots e^{a_r f_r} x_1^{j_1} \cdots x_{n+t}^{j_{n+t}} | f_1, \dots, f_r \in P, a_1, \dots, a_r \in \mathbb{Z}, j_1, \dots, j_n \in \mathbb{Z}, j_{n+1}, \dots, j_{n+t} \in \mathbb{N}\}$$

We then denote $\partial_{h_1}^{r_1} \cdots \partial_{h_r}^{r_r}$ by the composition of the partial derivatives $\partial_{h_1}, \dots, \partial_{h_r}$ on $\mathbb{F}[e^{\pm[P]}, n, t]$ with appropriate exponents where $1 \leq h_1, \dots, h_r \leq n + t$ and ∂_h^0 , $1 \leq h \leq n + t$, denotes the identity map on $\mathbb{F}[e^{\pm[P]}, n, t]$. For any $\alpha_u \in P \subset P_m$, let \mathfrak{A}_{α_u} be an additive subgroup

Received October 17, 2017. Revised May 1, 2018. Accepted May 12, 2018.

2010 Mathematics Subject Classification. 17B40, 17B56.

Key words and phrases. outer, non-associative algebra, derivation.

of \mathbb{F} such that \mathfrak{A}_{α_u} contains \mathbb{Z} . Consider now the (free) \mathbb{F} -vector space $N(e^{\mathfrak{A}_P}, n, t)_k$ (resp. $N(e^{\mathfrak{A}_P}, n, t)_{k+}$) whose basis is the set

$$(1) \quad \mathbf{B}_1 = \{e^{a_1 f_1} \cdots e^{a_r f_r} x_1^{j_1} \cdots x_{n+t}^{j_{n+t}} \partial_{h_1}^{r_1} \cdots \partial_{h_r}^{r_r} \mid a_1 \in \mathfrak{A}_{\alpha_1}, \dots, a_r \in \mathfrak{A}_{\alpha_r}, \\ f_1, \dots, f_r \in P, h_1, \dots, h_r \leq n+t, r_1 + \dots + r_r \leq k \in \mathbb{N} \quad (\text{resp. } \mathbb{N}^+)\}$$

If we define the multiplication $*$ on $N(e^{\mathfrak{A}_P}, n, t)_k$ as follows:

$$(2) \quad f \partial_{h_1}^{p_1} \cdots \partial_{h_r}^{p_r} * g \partial_{u_1}^{v_1} \cdots \partial_{u_q}^{v_q} = f(\partial_{h_1}^{p_1} \cdots \partial_{h_r}^{p_r}(g)) \partial_{u_1}^{v_1} \cdots \partial_{u_q}^{v_q}$$

for any $f \partial_{h_1}^{p_1} \cdots \partial_{h_r}^{p_r}, g \partial_{u_1}^{v_1} \cdots \partial_{u_q}^{v_q} \in N(e^{\mathfrak{A}_P}, n, t)_k$, then we define the combinatorial non-associative algebra $WN(e^{\mathfrak{A}_P}, n, t)_k$ whose underlying vector space is $N(e^{\mathfrak{A}_P}, n, t)_k$ and whose multiplication is $*$ in (2) (see [1], [5], [13] and [14]). The non-associative subalgebra $WN(e^{\mathfrak{A}_P}, n, t)_{<k>}$ of the algebra $WN(e^{\mathfrak{A}_P}, n, t)_k$ is generated by

$$(3) \quad \{f \partial_{h_1}^{r_1} \cdots \partial_{h_r}^{r_r} \mid f \in \mathbf{B}, 1 \leq h_1, \dots, h_r \leq n+t, r_1 + \dots + r_r = k \in \mathbb{N}^+\}.$$

The non-associative subalgebra $WN(e^{\mathfrak{A}_P}, n, t)_{[k]}$ of the algebra $WN(e^{\mathfrak{A}_P}, n, t)_k$ is generated by

$$(4) \quad \{f \partial_h^k \mid f \in \mathbf{B}, 1 \leq h \leq n+t\}.$$

For an algebra A and $l \in A$, an element $l_1 \in A$ is a right (resp. left) identity of l , if $l * l_1 = l$ (resp. $l_1 * l = l$) holds. The set of all right identities of $WN(e^{\mathfrak{A}_P}, n, t)_{[1]}$ is $\{\sum_{1 \leq u \leq n+t} x_u \partial_u + \sum_{1 \leq u \leq n+t} c_u \partial_u \mid c_u \in \mathbb{F}\}$. There is no left identity of $WN(e^{\mathfrak{A}_P}, n, t)_{k+}$. The algebra $WN(e^{\mathfrak{A}_P}, n, t)_k$ has the left identity 1. If A is an associative \mathbb{F} -algebra, then the anti-symmetrized algebra of A is a Lie algebra relative to the commutator $[x, y] := xy - yx$, (See [9]). For a general non-associative \mathbb{F} -algebra N we define in the same way its antisymmetrized algebra N^- . In case N^- is a Lie algebra we shall say that N is Lie admissible. For $S \subset N^-$, an element l is ad-diagonal with respect to S if for any $l_1 \in S$, $[l, l_1] = cl_1$ for $c \in \mathbb{F}$. The algebra $WN(e^{\mathfrak{A}_P}, n, t)_{[1]}$ is Lie admissible (see [8] and [15]). Since the cardinality $|P|$ of P is 2^m , for all $\alpha \in P_m$, if \mathfrak{A}_α is \mathbb{Z} , then the algebra $WN(e^{\mathfrak{A}_{P_m}}, n, t)_k$ is \mathbb{Z}^{2^m} -graded as follows:

$$(5) \quad WN(e^{\mathfrak{A}_{P_m}}, n, t)_k = \bigoplus_{(a_1, \dots, a_{m^2})} N_{(a_1, \dots, a_{m^2})}$$

where $N_{(a_1, \dots, a_{m^2})}$ is the vector subspace of $WN(e^{\mathfrak{A}_{P_m}}, n, t)_k$ spanned by

$$\{e^{a_1 f_1} \cdots e^{a_r f_r} x_1^{j_1} \cdots x_{n+t}^{j_{n+t}} \mid j_1, \dots, j_n \in \mathbb{Z}, j_{n+1}, \dots, j_{n+t} \in \mathbb{N}\}.$$

This implies that $WN(e^{\mathfrak{A}_P}, n, t)_k$ and $WN(e^{\mathfrak{A}_P}, n, t)_{k+}$ are appropriate graded algebras as (5) (see [11]). Thus throughout the paper, the

$(0, \dots, 0)$ -homogeneous component N_0 of $WN(e^{\mathfrak{A}P}, n, t)_k$ is the subalgebra $WN(0, n, t)_k$ of $WN(e^{\mathfrak{A}P}, n, t)_k$. For any standard basis element $e^{a_1 f_1} \dots e^{a_r f_r} x_1^{j_1} \dots x_{n+t}^{j_{n+t}} \partial_{t_1}^{r_1} \dots \partial_{t_r}^{r_r}$ of $WN(e^{\mathfrak{A}P}, n, t)_k$, define the homogeneous degree as follows:

$$hd(e^{a_1 f_1} \dots e^{a_r f_r} x_1^{j_1} \dots x_{n+t}^{j_{n+t}} \partial_{t_1}^{r_1} \dots \partial_{t_r}^{r_r}) = \sum_{u=1}^{n+t} |j_u|$$

where $|j_u|$ is the absolute value of j_u for $1 \leq u \leq n + t$. For any element $l \in WN(e^{\mathfrak{A}P}, n, t)_k$, define $hd(l)$ as the highest homogeneous degree of each monomial of l . Note that the set of all right annihilators of $WN(e^{\mathfrak{A}P}, n, t)_k$ is the subalgebra T_{n+t} of $WN(e^{\mathfrak{A}P}, n, t)_k$ which is spanned by $\{\partial_{t_1}^{r_1} \dots \partial_{t_r}^{r_r} | 1 \leq t_1, \dots, t_r \leq n + t, r_1 + \dots + r_r \leq k \in \mathbb{N}\}$. For a given algebra A , $Out(A)$ (resp. $Inn(A)$) is the set of all the outer (resp. inner) derivations of A and $Der(A)$ is the set of all the derivations of A . An algebra A is purely outer, if every derivation of A is outer i.e., $Der(A) = Out(A)$. There are various papers on studying the derivations of a non-associative algebra and an anti-symmetrized algebra (see [1], [2], [3], [5], [6], [7], [10], [12], [14]). We find all the derivations of a growing algebra in section 2.

2. Derivations of the non-associative algebra

$$WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$$

For this section, the set of all right annihilators T_3 of

$$WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$$

is spanned by $\{\partial_1, \partial_2, \partial_3\}$. The algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ is \mathbb{Z} -graded (see [5]). The algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ and the Lie algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}^-$ are simple (see [3] and [11]).

Note 1. For any basis elements $\partial_u, x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_u, e^{p x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_u, r_i \geq 1, 1 \leq u \leq 3$, of $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_1$, and for any $c \in \mathbb{F}, p \in \mathbb{Z}$, if we define an \mathbb{F} -linear map D_c from the algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_1$ to itself as follows:

$$\begin{aligned} D_c(\partial_u) &= 0, \\ (6) \quad D_c(x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_u) &= 0, \\ D_c(e^{p x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_u) &= p c e^{p x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_u, \end{aligned}$$

then the map D_c can be linearly extended to a non-associative algebra derivation of $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ (see [4], [6], [8] and [10]). \square

Lemma 2.1. *For any derivation D of $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ and for any basis elements $\partial_u, x_1^i x_2^j x_3^k \partial_u, 1 \leq u \leq 3$, of $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$, we have that*

$$\begin{aligned} D(\partial_u) &= 0, \\ D(x_1^i x_2^j x_3^k \partial_u) &= ic_{0,0,0,1} x_1^{i-1} x_2^j x_3^k \partial_u + jd_{0,0,0,2} x_1^i x_2^{j-1} x_3^k \partial_u \\ &\quad + kr_{0,0,0,3} x_1^i x_2^j x_3^{k-1} \partial_u \end{aligned}$$

hold with appropriate coefficients where $1 \leq u \leq 3$.

Proof. Let D be the derivation in the lemma. Since the algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ is \mathbb{Z} -graded, $D(\partial_1)$ is the sum of terms in different homogeneous components of $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ in (5). So $D(\partial_1)$ can be written as follows:

$$\begin{aligned} D(\partial_1) &= \sum_{r_1, r_2, r_3, i, j, k \geq 0} \alpha_{r_1, r_2, r_3, i, j, k, 1} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^i x_2^j x_3^k \partial_1 \\ &\quad + \sum_{r_1, r_2, r_3, i, j, k \geq 0} \alpha_{r_1, r_2, r_3, i, j, k, 2} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^i x_2^j x_3^k \partial_2 \\ &\quad + \sum_{r_1, r_2, r_3, i, j, k \geq 0} \alpha_{r_1, r_2, r_3, i, j, k, 3} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^i x_2^j x_3^k \partial_3 \end{aligned}$$

with appropriate coefficients. Since ∂_1 centralizes itself, we have that $D(\partial_1)$ is in the right annihilator of ∂_1 , i.e.,

$$\begin{aligned}
 & \partial_1 * D(\partial_1) \\
 &= \sum_{r_1, r_2, r_3, i, j, k \geq 0} pr_1 \alpha_{r_1, r_2, r_3, i, j, k, 1} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i+r_1-1} x_2^{j+r_2} x_3^{k+r_3} \partial_1 \\
 &+ \sum_{i \geq 1, r_1, r_2, r_3, j, k \geq 0} i \alpha_{r_1, r_2, r_3, i, j, k, 1} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i-1} x_2^j x_3^k \partial_1 \\
 &+ \sum_{r_1, r_2, r_3, i, j, k \geq 0} pr_1 \alpha_{r_1, r_2, r_3, i, j, k, 2} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i+r_1} x_2^{j+r_2} x_3^{k+r_3} \partial_2 \\
 (7) \quad &+ \sum_{i \geq 1, r_1, r_2, r_3, j, k \geq 0} i \alpha_{r_1, r_2, r_3, i, j, k, 2} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i-1} x_2^j x_3^k \partial_2 \\
 &+ \sum_{r_1, r_2, r_3, i, j, k \geq 0} pr_1 \alpha_{r_1, r_2, r_3, i, j, k, 3} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i+r_1} x_2^{j+r_2} x_3^{k+r_3} \partial_3 \\
 &+ \sum_{i \geq 1, r_1, r_2, r_3, j, k \geq 0} i \alpha_{r_1, r_2, r_3, i, j, k, 3} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i-1} x_2^j x_3^k \partial_3 \\
 &= 0
 \end{aligned}$$

with appropriate coefficients. By (7), we have that $\alpha_{r_1, r_2, r_3, i, j, k, 1}$, $\alpha_{r_1, r_2, r_3, i, j, k, 2}$, and $\alpha_{r_1, r_2, r_3, i, j, k, 3}$, are zeros, $r_1, r_2, r_3, i, j, k \geq 0$. Thus $D(\partial_1)$ is zero. Similarly we can prove that $D(\partial_2)$ and $D(\partial_3)$ are also zeros. Since ∂_1 centralizes $x_1 \partial_1$, we can also prove that

$$D(x_1 \partial_1) = c_1 \partial_1 + c_2 \partial_2 + c_3 \partial_3.$$

Since $x_1 \partial_1$ is an idempotent, we can prove that $c_2 = 0, c_3 = 0$. This implies that $D(x_1 \partial_1) = c_1 \partial_1$. Since $D(\partial_1 * x_1^2 \partial_1) = 2D(x_1 \partial_1)$, we are also able to prove that

$$\begin{aligned}
 D(x_1^2 \partial_1) &= 2c_1 x_1 \partial_1 + \sum_{j, k} t_{0, j, k, 1} x_2^j x_3^k \partial_1 \\
 &+ \sum_{j, k} t_{0, j, k, 2} x_2^j x_3^k \partial_2 + \sum_{j, k} t_{0, j, k, 3} x_2^j x_3^k \partial_3
 \end{aligned}$$

where $t_{0, j, k, 1}, t_{0, j, k, 2}, t_{0, j, k, 3} \in \mathbb{F}$ for all j and k . Since $D(x_1 \partial_1 * x_1^2 \partial_1) = 2D(x_1^2 \partial_1)$, we have that $t_{0, j, k, 1} = t_{0, j, k, 2} = t_{0, j, k, 3} = 0$ for all j and k . This implies that

$$\begin{aligned}
 D(x_1 \partial_1) &= c_1 \partial_1, \\
 D(x_1^2 \partial_1) &= 2c_1 x \partial_1
 \end{aligned}$$

hold. By $D(\partial_1 * x_1^3 \partial_1) = 3D(x_1^2 \partial_1)$, we have that

$$\begin{aligned} D(x_1^3 \partial_1) &= 3c_1 x_1^2 \partial_1 + \sum_{j,k} s_{0,j,k,1} x_2^j x_3^k \partial_1 \\ &+ \sum_{j,k} s_{0,j,k,2} x_2^j x_3^k \partial_2 + \sum_{j,k} s_{0,j,k,3} x_2^j x_3^k \partial_3, \end{aligned}$$

where $s_{0,j,k,1}, s_{0,j,k,2}, s_{0,j,k,3} \in \mathbb{F}$ for all j and k . By $D(x_1 \partial_1 * x_1^3 \partial_1) = 3D(x_1^3 \partial_1)$, we have that $s_{0,j,k,1} = s_{0,j,k,2} = s_{0,j,k,3} = 0$ for all j and k and $D(x_1^3 \partial_1) = 3c_1 x_1^2 \partial_1$. Since $D(x_1^2 \partial_1 * x_1^{i-1} \partial_1) = (i-1)D(x_1^i \partial_1)$, by induction on i of $x_1^i \partial_1$, we are able to prove that

$$D(x_1^i \partial_1) = ic_1 x_1^{i-1} \partial_1.$$

Similarly we are also able to prove that

$$\begin{aligned} D(x_2^j \partial_2) &= jd_2 x_2^{j-1} \partial_2, \\ D(x_3^k \partial_3) &= kh_3 x_3^{k-1} \partial_3. \end{aligned}$$

Since $\partial_u, 1 \leq u \leq 3$, is in the left annihilator of $x_1 \partial_2$, we can prove that $D(x_1 \partial_2) = \alpha_1 \partial_1 + \alpha_2 \partial_2 + \alpha_3 \partial_3$. By $D(x_1 \partial_1 * x_1 \partial_2) = D(x_1 \partial_2)$, we can also prove that $\alpha_1 = \alpha_3 = 0, \alpha_2 = c_1$. This implies that $D(x_1 \partial_2) = c_1 \partial_2$. Since $D(x_1^2 \partial_1 * x_1^{i-1} \partial_2) = (i-1)D(x_1^i \partial_2)$, by induction on i of $x_1^i \partial_2$, we can prove that

$$D(x_1^i \partial_2) = ic_1 x_1^{i-1} \partial_2.$$

Similarly we are able to prove that

$$\begin{aligned} D(x_1^i \partial_3) &= ic_1 x_1^{i-1} \partial_3, \\ D(x_2^j \partial_u) &= jd_2 x_2^{j-1} \partial_u, \\ D(x_3^k \partial_u) &= kh_3 x_3^{k-1} \partial_u \end{aligned}$$

where $1 \leq u \leq 3$. By $D(x_1^i \partial_2 * x_2^{j+1} \partial_u) = (j+1)D(x_1^i x_2^j \partial_u)$, we have that

$$D(x_1^i x_2^j \partial_u) = ic_1 x_1^{i-1} x_2^j \partial_u + jd_2 x_1^i x_2^{j-1} \partial_u$$

where $1 \leq u \leq 3$. Since $D(x_1^i x_2^j \partial_3 * x_3^{k+1} \partial_u) = (k+1)D(x_1^i x_2^j x_3^k \partial_u)$, we are also able to prove that

$$D(x_1^i x_2^j x_3^k \partial_u) = ic_1 x_1^{i-1} x_2^j x_3^k \partial_u + jd_2 x_1^i x_2^{j-1} x_3^k \partial_u + kh_3 x_1^i x_2^j x_3^{k-1} \partial_u.$$

where $1 \leq u \leq 3$. So we have proven the lemma. \square

Lemma 2.2. *For any derivation D of the algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ and for basis elements $x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_u$, $e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u$, $1 \leq u \leq 3$, of $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$, we have that*

$$\begin{aligned} D(x_1^i x_2^j x_3^k \partial_u) &= 0, \\ D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u) &= c_{r_1, r_2, r_3} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u, \end{aligned}$$

hold where $c \in \mathbb{F}$.

Proof. Let D be the derivation in the lemma. Since the algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ is \mathbb{Z} -graded, $D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1)$ is the sum of terms in different homogeneous components of $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ in (5). Assume that

$$\begin{aligned} D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) &= \sum_{r_1, r_2, r_3, i, j, k \geq 0} a_{r_1, r_2, r_3, i, j, k, 1} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^i x_2^j x_3^k \partial_1 \\ &+ \sum_{r_1, r_2, r_3, i, j, k \geq 0} a_{r_1, r_2, r_3, i, j, k, 2} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^i x_2^j x_3^k \partial_2 \\ &+ \sum_{r_1, r_2, r_3, i, j, k \geq 0} a_{r_1, r_2, r_3, i, j, k, 3} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^i x_2^j x_3^k \partial_3 \end{aligned}$$

with appropriate coefficients. We have that

$$\begin{aligned} (8) \quad D(\partial_1 * e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) &= r_1 D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1-1} x_2^{r_2} x_3^{r_3} \partial_1) \\ &= \sum_{r_1, r_2, r_3, i, j, k \geq 0} pr_1 a_{r_1, r_2, r_3, i, j, k, 1} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1+i-1} x_2^{r_2+j} x_3^{r_3+k} \partial_1 \\ &+ \sum_{i \geq 1, r_1, r_2, r_3, j, k \geq 0} ia_{r_1, r_2, r_3, i, j, k, 1} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i-1} x_2^j x_3^k \partial_1 \\ &+ \sum_{r_1, r_2, r_3, i, j, k \geq 0} pr_1 a_{r_1, r_2, r_3, i, j, k, 2} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1+i-1} x_2^{r_2+j} x_3^{r_3+k} \partial_2 \\ &+ \sum_{i \geq 1, r_1, r_2, r_3, j, k \geq 0} ia_{r_1, r_2, r_3, i, j, k, 2} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i-1} x_2^j x_3^k \partial_2 \\ &+ \sum_{r_1, r_2, r_3, i, j, k \geq 0} pr_1 a_{r_1, r_2, r_3, i, j, k, 3} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1+i-1} x_2^{r_2+j} x_3^{r_3+k} \partial_3 \\ &+ \sum_{i \geq 1, r_1, r_2, r_3, j, k \geq 0} ia_{r_1, r_2, r_3, i, j, k, 3} e^{px_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i-1} x_2^j x_3^k \partial_3 \end{aligned}$$

and

$$\begin{aligned}
 (9) \quad & D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1 * x_1^{r_1} x_2^{r_2} x_3^{r_3} \partial_1) = r_1 D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1-1} x_2^{r_2} x_3^{r_3} \partial_1) \\
 & = D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) * x_1^{r_1} x_2^{r_2} x_3^{r_3} \partial_1 + e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1 * D(x_1^{r_1} x_2^{r_2} x_3^{r_3} \partial_1) \\
 & = r_1 \sum_{r_1, r_2, r_3, i, j, k \geq 0} a_{i, j, k, 1} e^{p x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i+r_1-1} x_2^{j+r_2} x_3^{k+r_3} \partial_1 \\
 & \quad + r_2 \sum_{r_1, r_2, r_3, i, j, k \geq 0} a_{r_1, r_2, r_3, i, j, k, 2} e^{p x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i+r_1} x_2^{j+r_2-1} x_3^{k+r_3} \partial_1 \\
 & \quad + r_3 \sum_{r_1, r_2, r_3, i, j, k \geq 0} a_{r_1, r_2, r_3, i, j, k, 3} e^{p x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{i+r_1} x_2^{j+r_2} x_3^{k+r_3-1} \partial_1 \\
 & \quad + (r_1 - 1) r_1 c_1 e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1-2} x_2^{r_2} x_3^{r_3} \partial_1 \\
 & \quad + r_1 r_2 d_2 e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1-1} x_2^{r_2-1} x_3^{r_3} \partial_1 \\
 & \quad + r_1 r_3 h_3 e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1-1} x_2^{r_2} x_3^{r_3-1} \partial_1.
 \end{aligned}$$

By comparing (8) and (9), we have that

$$\begin{aligned}
 & p = 1, \\
 & a_{r_1, r_2, r_3, i, j, k, 2} = a_{r_1, r_2, r_3, i, j, k, 3} = 0, \quad r_1, r_2, r_3, i, j, k \geq 0, \\
 & a_{r_1, r_2, r_3, i, j, k, 1} = 0, \quad i \geq 1, \quad \text{and} \\
 & c_1 = d_2 = h_3 = 0.
 \end{aligned}$$

This implies that

$$(10) \quad D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) = \sum_{r_1, r_2, r_3, j, k \geq 0} a_{r_1, r_2, r_3, 0, j, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_2^j x_3^k \partial_1,$$

and we also have

$$D(x_1^i x_2^j x_3^k \partial_u) = 0$$

where $1 \leq u \leq 3$. Similarly we can prove that

$$(11) \quad D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u) = \sum_{r_1, r_2, r_3, j, k \geq 0} a_{r_1, r_2, r_3, 0, j, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_2^j x_3^k \partial_u.$$

where $2 \leq u \leq 3$. Since

$$\begin{aligned}
 & D(\partial_2 * e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) = r_2 D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1} x_2^{r_2-1} x_3^{r_3} \partial_1) \\
 & = r_2 \sum_{r_1, r_2, r_3, j, k \geq 0} a_{r_1, r_2, r_3, 0, j, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1} x_2^{r_2+j-1} x_3^{r_3+k} \partial_1 \\
 & \quad + \sum_{j \geq 1, r_1, r_2, r_3, k \geq 0} j a_{r_1, r_2, r_3, 0, j, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_2^{j-1} x_3^k \partial_1
 \end{aligned}$$

and

$$\begin{aligned} D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_2 * x_1^{r_1} x_2^{r_2} x_3^{r_3} \partial_1) &= r_2 D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1} x_2^{r_2-1} x_3^{r_3} \partial_1) \\ &= r_2 \sum_{r_1, r_2, r_3, j, k \geq 0} a_{0, j, k, 1} e^{x_1^{r_1} x_1^{r_1} x_2^{r_2+j-1} x_3^{r_3+k}} \partial_1, \end{aligned}$$

we have that $a_{r_1, r_2, r_3, 0, j, k, 1} = 0, j \geq 1$. This implies that

$$(12) \quad D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) = \sum_{r_1, r_2, r_3, k \geq 0} a_{r_1, r_2, r_3, 0, 0, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_3^k \partial_1.$$

Similarly we can prove that

$$(13) \quad D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u) = \sum_{r_1, r_2, r_3, k \geq 0} a_{r_1, r_2, r_3, 0, 0, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_3^k \partial_u.$$

where $2 \leq u \leq 3$. Since

$$\begin{aligned} D(\partial_3 * e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) &= r_3 D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1} x_2^{r_2} x_3^{r_3-1} \partial_1) \\ &= \sum_{r_1, r_2, r_3, k \geq 0} a_{r_1, r_2, r_3, 0, 0, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1} x_2^{r_2} x_3^{r_3+k-1} \partial_1 \\ &+ \sum_{r_1, r_2, r_3 \geq 0, k \geq 1} k a_{r_1, r_2, r_3, 0, 0, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_3^{k-1} \partial_1 \end{aligned}$$

and

$$\begin{aligned} D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_3 * x_1^{r_1} x_2^{r_2} x_3^{r_3} \partial_1) &= r_3 D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1} x_2^{r_2} x_3^{r_3-1} \partial_1) \\ &= \sum_{r_1, r_2, r_3, k \geq 0} r_3 a_{r_1, r_2, r_3, 0, 0, k, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} x_1^{r_1} x_2^{r_2} x_3^{r_3+k-1} \partial_1. \end{aligned}$$

Thus we have that $a_{r_1, r_2, r_3, 0, 0, k, 1} = 0, k \geq 1$. This implies that

$$(14) \quad D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1) = a_{r_1, r_2, r_3, 0, 0, 0, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1.$$

By (14) and $D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_1 * x_1 \partial_u) = D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u)$, we can prove that

$$D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u) = a_{r_1, r_2, r_3, 0, 0, 0, 1} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u,$$

where $2 \leq u \leq 3$. Putting $c_{r_1, r_2, r_3} = a_{r_1, r_2, r_3, 0, 0, 0, 1}$, we have that $D(e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u) = c_{r_1, r_2, r_3} e^{x_1^{r_1} x_2^{r_2} x_3^{r_3}} \partial_u$. Therefore we have proven the lemma. \square

Theorem 2.3. For any derivation D of the algebra $WN(e^{\pm x_1^{r_1} x_2^{r_2} x_3^{r_3}}, 0, 3)_{[1]}$ and for basis elements

$e^{px_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u$, $1 \leq u \leq 3$, of $WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]}$, we have that

$$D(e^{px_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u) = pc_{r_1,r_2,r_3}e^{px_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u$$

hold where $p \in \mathbb{Z}$ and $c \in \mathbb{F}$.

Proof. Let D be the derivation in the lemma.

By $D(e^{x_1^{r_1}x_2^{r_2}x_3^{r_3}}\partial_1 * x_1^{i+1}x_2^jx_3^k\partial_u) = (i+1)D(e^{x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u)$, we are able to prove that $D(e^{x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u) = c_{r_1,r_2,r_3}e^{x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u$ for $1 \leq u \leq 3$, with appropriate coefficients. By

$$D(e^{x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^{i-r_1+1}x_2^{j-r_2}x_3^{k-r_3}\partial_1 * e^{x_1^{r_1}x_2^{r_2}x_3^{r_3}}\partial_u) = r_1D(e^{2x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u),$$

we prove that

$$D(e^{2x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u) = 2c_{r_1,r_2,r_3}e^{2x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u.$$

By induction on $p \in \mathbb{Z}$ of $e^{px_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u$ and

$$D(e^{(p-1)x_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^{i-r_1+1}x_2^{j-r_2}x_3^{k-r_3}\partial_1 * e^{x_1^{r_1}x_2^{r_2}x_3^{r_3}}\partial_u) = r_1D(e^{px_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u),$$

we are able to prove that

$$D(e^{px_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u) = pc_{r_1,r_2,r_3}e^{px_1^{r_1}x_2^{r_2}x_3^{r_3}}x_1^ix_2^jx_3^k\partial_u.$$

Therefore the proof is completed. □

Theorem 2.4. For any $D \in Der_{non}(WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]})$, D is the linear sum of the derivations D_c as shown in Note 1 where $c \in \mathbb{F}$. Every derivation of the algebra $WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]}$ is outer.

Proof. The proofs of the theorem are straightforward by Lemma 2.2, Theorem 2.3, and the fact that the derivation of Note 1 cannot be inner. This completes the proof of the theorem. □

Corollary 2.5. The dimension of $Der_{non}(WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]})$ of the algebra $WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]}$ is one. For any derivation D of $Der_{non}(WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]})$, $D(\mathfrak{A}) = 0$ holds where \mathfrak{A} is the zero-homogeneous component of $WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]}$ in (5) (see [9]).

Proof. The proofs of the corollary are straightforward by Lemma 2.2 and Note 1. □

Proposition 2.6. If A is not a purely outer algebra, then algebra A and $WN(e^{\pm x_1^{r_1}x_2^{r_2}x_3^{r_3}}, 0, 3)_{[1]}$ are not isomorphic.

Proof. The proof of the proposition is straightforward by Theorem 2.4. □

References

- [1] Seul Hee Choi and Ki-Bong Nam, *The Derivation of a Restricted Weyl Type Non-Associative Algebra*, Hadronic Journal, **28(3)**, (2005), 287-295.
- [2] Seul Hee Choi, *A growing algebra containing the polynomial ring*, Honam Mathematical Journal, **32(3)**, (2010), 467-480.
- [3] Seul Hee Choi, *An algebra with right identities and its antisymmetrized algebra*, Honam Mathematical Journal, **29(2)**, (2007), 213-222.
- [4] Seul Hee Choi, *New algebras using additive abelian groups I*, Honam Mathematical Journal, **31(3)**, (2009), 407-419.
- [5] Seul Hee Choi and Ki-Bong Nam, *Weyl type non-associative algebra using additive groups I*, Algebra Colloquium, **14(3)** (2007), 479-488.
- [6] Seul Hee Choi and Ki-Bong Nam, *Derivations of a restricted Weyl Type Algebra I*, Rocky Mountain Math. Journals, **37(6)**, (2007), 67-84.
- [7] Seul Hee Choi, Hong Goo Park, Moon-Ok Wang, and Ki-Bong Nam, *Combinatorial algebra and its antisymmetrized algebra I*, Algebra Colloquium, **22(1)**, (2015), 823-834.
- [8] Seul Hee Choi, Jongwoo Lee, and Ki-Bong Nam, *Derivations of a restricted Weyl type algebra containing the polynomial ring*, Communication in Algebra, **36(9)**, (2008), 3435 - 3446.
- [9] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag, New York, (1987), 7-21.
- [10] T. Ikeda, N. Kawamoto and Ki-Bong Nam, *A class of simple subalgebras of Generalized W algebras*, Proceedings of the International Conference in 1998 at Pusan (Eds. A. C. Kim), Walter de Gruyter GmbH Co. KG, (2000), 189-202.
- [11] V. G. Kac, *Description of the filtered Lie algebras with which graded Lie algebras of Cartan type are associated*, Izv. Akad. Nauk SSSR, Ser. Mat. Tom, **38**, (1974), 832-834.
- [12] Jongwoo Lee and Ki-bong Nam, *Non-Associative Algebras containing the Matrix Ring*, Linear Algebra and its Applications **429(1)**, (2008), Pages 72-78.
- [13] Ki-Bong Nam, *Generalized W and H Type Lie Algebras*, Algebra Colloquium **6(3)**, (1999), 329-340.
- [14] Ki-Bong Nam, *On Some Non-Associative Algebras Using Additive Groups*, Southeast Asian Bulletin of Mathematics, **27**, Springer Verlag, (2003), 493-500.
- [15] R. D. Schafer, *Introduction to nonassociative algebras*, Dover, (1995), 128-138.

Seul Hee Choi
Department of Mathematics, Jeonju University,
Jeonju, Jeonbuk 55069, Korea.
E-mail: chois@jj.ac.kr