

ANIMAL

Storage stability of dry-aged beef: the effects of the packaging method and storage temperature

Juhui Choe¹, Kwan Tae Kim^{2,3}, Hyun Jung Lee¹, Jungmin Oh¹, Hyun Cheol Kim¹, Bumjin Park¹, Yang Il Choi³, Cheorun Jo^{1,4,*}

¹Department of Agricultural Biotechnology, Center for Food and Bioconvergence, and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea

²Korea Institute for Animal Products Quality Evaluation, Sejong 30100, Korea

³Department of Animal Science, Chungbuk National University, Cheongju 28644, Korea

⁴Institute of Green Bio Science and Technology, Seoul National University, Pyeongchang 25354, Korea

*Corresponding author: cheorun@snu.ac.kr


Abstract

Different packaging methods and storage temperatures were tested to determine the storage stability of beef dry-aged for 21 days based on microbial, physicochemical, and sensory qualities. After completion of the dry aging, the dried surface of beef sirloin was trimmed off, and the beef was packaged using two different methods (oxygen-permeable wrap or vacuum packaging) and stored at different temperatures ($3 \pm 2^\circ\text{C}$ or $-23 \pm 2^\circ\text{C}$) for 0, 7, 14, or 21 days. Lipid oxidation and the sensory quality of the dry-aged beef were not affected by the packaging method and storage temperature during storage. No microbial growth was observed over the storage period in the vacuum-packaged dry-aged beef, regardless of the storage temperature. However, dry-aged beef in the oxygen-permeable wrap packaging showed microbial spoilage with $8.82 \log \text{CFU/g}$ at day 7 of the refrigerated storage. The vacuum-packaged dry-aged beef showed the lowest values ($p < 0.05$) in a^* and chroma at days 14 and 21 at 3°C , and days 7 and 14 at -23°C , respectively. Therefore, it is recommended that dry-aged beef with wrap packaging stored in refrigerated conditions should be consumed as quickly as possible due to microbial growth. For long-term storage, dry-aged beef should be frozen because freezing can extend the color stability up to day 21 of storage without adverse effects on the hygienic or meat quality aspects of dry-aged beef.

Keywords : dry aging, packaging method, storage temperature, microbial quality, sensory quality

Introduction

The aging process is widely used for the enhancement of beef palatability, including tenderness and flavor (Sitz et al., 2006; Lee et al., 2018). There are two types of aging: Wet and dry aging. In wet aging, meat is vacuum-packed and aged under refrigeration (Smith et al., 2008). Dry aging refers to the exposure of meat to air at controlled



OPEN ACCESS

Citation: Choe J, Kim KT, Lee HJ, Oh JM, Kim HC, Park BJ, Choi YI, Jo C. 2018. Storage stability of dry-aged beef: the effects of the packaging method and storage temperature. Korean Journal of Agricultural Science. <https://doi.org/10.7744/kjoas.20180021>

DOI: <https://doi.org/10.7744/kjoas.20180021>

Editor: Ki Teak Lee, Chungnam National University, Korea

Received: September 27, 2017

Revised: February 22, 2018

Accepted: March 20, 2018

Copyright: © 2018 Korean Journal of Agricultural Science



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

temperature, relative humidity, or/and air flow. Dry-aged beef is reported to have a more “Beefy” and “Brown-roasted” flavor compared to wet-aged or non-aged beef (Warren and Kastner, 1992). However, dry aging in beef leads to higher weight loss due to greater water and trim loss compared to wet-aged or non-aged beef (Parrish et al., 1991). Despite the high cost of the process, the application of dry aging in beef has rapidly increased in popularity recently due to the resulting enhanced palatability and unique flavor.

Among consumers who prefer dry-aged meat, there is a particular demand for safety while maintaining the characteristic flavor attributes, because of the high possibility of contamination in dry-aged meat during the aging or/and trimming process. Vacuum packaging, commonly used for the distribution or sale of meat, is utilized to control microbial growth in meat. Wrap packaging with oxygen-permeable polyvinyl chloride (PVC) film is also widely used for fresh beef during retail display (Jayasingh et al., 2001). The optimal storage period for wrap- or vacuum-packaged meat to maintain meat quality, including physicochemical and sensory properties, depends on the storage temperature (Jakobson and Bertelsen, 2000). It is especially important for dry-aged meat to determine the optimal storage conditions (temperature and storage period) for different packaging methods to maintain its inherent sensory characteristics.

To our knowledge, there has been no study determining the microbial, physicochemical, and sensory properties of differently packaged beef after dry aging with subsequent storage at different temperatures. In other words, it is necessary to determine the shelf-life of dry-aged beef in common storage conditions such that there is no adverse effect on meat quality. Doing so will allow the practical needs of producers and distributors in the rapidly growing dry-aged meat market to be met. Therefore, the objective of this study was to investigate the changes in microbial, physicochemical, and sensory properties of dry-aged beef resulting from different packaging methods and storage temperatures during extended storage times.

Materials and Methods

Raw materials and dry aging

A total of 28 sirloins (approximately 54-month-old Hanwoo cows; quality grade 2) were collected and transported in an iced condition (4°C) to Korea Institute for Animal Products Quality Evaluation (Sejong, Korea). Initial pH was measured for all samples prior to the dry aging process. Ten sirloins were designated as non-aged controls, and the other 18 (n = 6 for each of three treatments, described below) were dry aged at $2 \pm 1^\circ\text{C}$ (75% relative humidity) for 28 days. The non-aged control samples were frozen immediately. After 28 days of dry aging, the dried surfaces (crusts) of the dry-aged beef were trimmed off, and the beef was packed and stored using the following packaging methods and temperatures for 21 days: 1) overwrapped with oxygen-permeable PVC film and stored at $3 \pm 2^\circ\text{C}$; 2) vacuum-packaged and stored at $3 \pm 2^\circ\text{C}$; 3) vacuum-packaged and stored at $-23 \pm 2^\circ\text{C}$. Vacuum packaging was carried out

using polyethylene bags (O_2 permeability 2.3 mL/m²/d at 38°C; HFV-600L, Hankook Fudee Co., Ltd., Siheung, Korea). Meat samples were analyzed at days 0, 7, 14, and 21 of storage after the completion of dry aging and packaging.

Total aerobic bacteria (TAB)

Meat samples (5 g) were blended with sterile saline (45 mL, 0.85%) for 2 min using a BagMixer® 400 P blender (Interscience, St. Nom la Bretèche, France), and diluted (Yoon et al., 2017). Each sample (100 µL) was spread on plate count agar (Difco Laboratories, Detroit, MI, USA) to inspect TAB. The agar plates for TAB and were incubated at 37°C for 48 h. After incubation, microbial counts were calculated and expressed as log CFU/g sample.

2-Thiobarbituric acid-reactive substances (TBARS) value

Each sample (3.0 g) was homogenized (T25 Basic, Ika works, Staufen, Germany) in 9 mL of distilled water and 50 µL of butylated hydroxyl toluene as previously described (Khan et al., 2016). The homogenate (1 mL) was mixed with 2-thiobarbituric acid/trichloroacetic acid solution (2 mL). The mixture was heated at 90°C for 30 min in a water bath, cooled, and centrifuged at $2,090 \times g$ (Continent 512R, Hanil Co., Ltd., Incheon, Korea). The absorbance of the supernatant was measured at 532 nm using a spectrophotometer (X-ma 3100; Human Co. Ltd., Seoul, Korea) and 2-thiobarbituric acid-reactive substances (TBARS) value [mg malondialdehyde/kg meat sample] was calculated using a standard curve.

Instrumental color

The samples were bloomed for 30 min and CIE color values (L^* , a^* , and b^*) were determined using a spectrophotometer (CR400; Konica Minolta Censing Inc., Osaka, Japan). Before color measurement, the spectrophotometer was calibrated with a standard tile ($L^*=97.74$, $a^*=-0.06$, $b^*=1.76$). Each sample was measured three times, and the average value was used as one replicate. Chroma [$(a^{*2} + b^{*2})$] and hue angle [$\tan^{-1}(b^*/a^*)$] were derived from CIE L^* , a^* , and b^* values.

Sensory evaluation

Sensory evaluation was carried out with a consumer panel (total 10 panelists) to observe the change in and differences between vacuum-packed beef samples after dry aging with storage at different temperatures. The wrap-packed beef after dry aging could not be analyzed for sensory evaluation owing to microbial spoilage at day 7 of refrigerated storage. Each sample was cut to similar size ($50 \times 20 \times 6$ mm³, length \times width \times height), grilled until the core temperature reached 72°C, and served to the panelists. A 7-point hedonic scale (1, extremely dislike; 7, extremely like) was used to score juiciness, tenderness, flavor, and overall acceptability at day 0, 7, 14, and 21 of storage.

Statistical analysis

The experimental design was a randomized incomplete block design using trial as block. The non-aged meat (control, $n = 10$) and 3 treatments ($n = 6/\text{treatment}$) were assigned and a model was analyzed with fixed effects (packaging and temperature) and random effects (carcass and side of carcass). An interaction effect between storage temperature and storage time was considered for color values (L^* , a^* , b^* , chroma value, and hue angle) and lipid oxidation. A general linear model was generated using SAS 9.3 (SAS Institute Inc., Cary, NC, USA), and mean values \pm standard error of the mean (SEM) reported. Significant differences between the mean values were determined based on the Student-Newman-Keuls multiple comparison test at a level of $p < 0.05$.

Results and Discussion

Total aerobic bacteria

There was no significant difference in TAB counts between non-aged (control, 5.83 log CFU/g) and dry-aged beef (5.72 log CFU/g). The observation might be due to prevention of microbial penetration into the meat by crust formation during dry aging. This is consistent with the previously reported finding that dry aging for 28 days in beef strip loin did not affect the number of total aerobic bacteria ($p > 0.05$) (DeGeer et al., 2006). As presented in Table 1, the TAB counts of dry-aged beef with wrap packaging reached approximately 8.8 log CFU/g at day 7 of refrigerated storage ($3 \pm 2^\circ\text{C}$), indicating meat spoilage (Sofos et al., 2000). In the vacuum-packaged dry-aged beef, TAB counts were below 7 log CFU/g for the entire storage period regardless of storage temperature. This result could be because the application of vacuum packaging eliminated oxygen, led to retardation of aerobic bacteria, and reduced the growth rate of anaerobic and facultative bacteria (Bellés et al., 2017). This result indicates that the application of vacuum packaging to dry-aged beef can control microbial growth up to 21 days of storage at both 3°C and -23°C .

Table 1. Effect of different packaging methods and storage temperatures on total aerobic bacteria and lipid oxidation of dry-aged beef during storage for 21 days.

Traits	Packaging method/ storage temperature after dry aging	Storage days after dry aging				SEM*
		0	7	14	21	
Total aerobic bacteria (log CFU/g)	Wrap/ 3°C	5.72B	8.82Aa	- ^z	- ^z	0.242
	Vacuum/ 3°C	5.72	6.63b	6.63a	6.05	0.244
	Vacuum/ -23°C	5.72B	6.99Ab	5.69Bb	5.42B	0.269
	SEM ^y	0.228	0.258	0.216	0.257	
TBARS (mg MDA/kg meat sample)	Wrap/ 3°C	0.98	0.84	- ^z	- ^z	0.100
	Vacuum/ 3°C	0.98	0.79	1.15	1.07	0.161
	Vacuum/ -23°C	0.98	0.88	1.20	0.96	0.117
	SEM ^y	0.074	0.148	0.313	0.214	

All values are mean \pm standard error of the mean ^x $n = 24$, ^y $n = 18$.

^zThis experiment was not carried out because microbial numbers exceeded 7 log CFU/g at day 7 of storage.

A, B: Different letters within rows indicate statistically significant differences ($p < 0.05$).

a, b: Different letters within columns indicate statistically significant differences ($p < 0.05$).

2-Thiobarbituric acid-reactive substances (TBARS) value

Dry aging of beef led to an increase in lipid oxidation compared with non-aged control ($p < 0.05$; data not shown). This result may be due to the fact that the beef was exposed to air during dry aging. Higher lipid oxidation has previously been observed in dry-aged beef with the same quality grade used in this study compared with non-aged beef (Lee et al., 2015). The packaging method, storage temperature, and storage periods did not influence lipid oxidation in dry-aged beef, with TBARS values ranging from 0.84 to 1.20 ($p > 0.05$; Table 1). The TBARS values showed that there was no negative effect on meat quality as values remained below 1.2, which is considered as a threshold value for detection of “Off” flavor in meat (Younathan and Watts, 1959). A previous study reported that vacuum packaging of lamb meat led to effective reduction in lipid oxidation at 2 - 4°C after storage for 28 days (Fernandes et al., 2014). There was no interaction between storage temperature and storage periods in this study.

Instrumental color

Different L^* , a^* , and b^* values were observed depending on dry aging, packaging method, and storage temperature (Table 2). Dry aging led to significantly lower a^* , b^* , and chroma values in beef sirloin (Fig. 1). The L^* value (data not shown) and hue angle (Fig. 1) of the beef were not affected by dry aging ($p > 0.05$).

Table 2. Effect of different packaging methods and storage temperatures on instrumental color of dry-aged beef during storage for 21 days.

Traits	Packaging method/storage temperature after dry aging	Storage days after dry aging				SEM ^y
		0	7	14	21	
CIE L^*	Wrap/3°C	36.78	34.85	-	-	0.864
	Vacuum/3°C	36.78A	34.50AB	33.62B	35.95A	0.853
	Vacuum/- 23°C	36.78	36.05	35.36	37.22	1.120
	SEM ^z	0.562	1.280	0.703	1.372	
CIE a^*	Wrap/3°C	14.95A	12.56Bc	-	-	0.489
	Vacuum/3°C	14.95A	14.49Aa	12.74B	11.98B	0.471
	Vacuum/- 23°C	14.95A	9.97Bb	9.56B	15.65A	0.722
	SEM ^z	0.391	0.563	0.517	1.276	
CIE b^*	Wrap/3°C	11.09A	9.16B	-	-	0.407
	Vacuum/3°C	11.09A	9.98B	8.95B	9.63B	0.391
	Vacuum/- 23°C	11.09A	9.05B	8.85B	12.25A	0.571
	SEM ^z	0.235	0.507	0.399	0.936	
Chroma value	Wrap/3°C	18.63A	15.56Bb	-	-	0.714
	Vacuum/3°C	18.63A	17.60Aa	15.57Ba	15.39B	0.552
	Vacuum/- 23°C	18.63A	13.55Bc	13.07Bb	19.90A	0.927
	SEM ^z	0.420	0.610	0.555	1.533	
Hue angle	Wrap/3°C	36.65	36.12b	-	-	0.832
	Vacuum/3°C	36.65AB	34.55Bb	35.11Bb	38.85A	0.833
	Vacuum/- 23°C	36.65B	42.29Aa	42.90Aa	38.20AB	1.473
	SEM ^z	0.562	1.805	1.562	1.322	

All values are mean \pm standard error of the mean ^yn = 24, ^zn = 18.

A - B: Different letters within rows indicate statistically significant differences ($p < 0.05$).

a - c: Different letters within columns indicate statistically significant differences ($p < 0.05$).

compared with non-aged control; Fig. 1). The a^* value of the beef was affected by packaging method at day 7 of storage; vacuum-packaged dry-aged beef had higher a^* values than wrap-packaged dry-aged beef ($p < 0.05$). According to Ledward (1985), vacuum packed meat during storage tended to have relatively high a^* values as helping to maintain metmyoglobin reducing activity. Significant differences in a^* and b^* values were observed for each packaging method and storage temperature. During refrigerated storage, vacuum-packaged dry-aged beef maintained a^* and chroma values up to day 7, but significant decrease in a^* values were observed thereafter until day 21. During frozen storage, a^* values of vacuum-packaged dry-aged beef decreased until day 14. Dry-aged beef with vacuum packaging showed significantly higher a^* and chroma values at day 21 of frozen storage compared with day 7 and day 14. In other words, for dry-aged beef under vacuum packaging, higher color stability was observed with refrigerated storage compared to frozen storage over a short-term storage period (7 days). Moreover, an interaction was observed between storage temperature and storage periods for a^* , b^* , and chroma values.

Sensory properties

Storage temperature, storage period, and interaction between storage temperature and storage period did not affect the sensory properties of dry-aged beef under vacuum packaging ($p > 0.05$; Table 3). For frozen stored beef after dry aging, no significant changes in sensory properties were found. Based on the result, dry aged beef with tender and flavorful could be kept until day 21 of storage at -23°C .




Fig. 1. Effect of dry aging on instrumental color of beef. Standard error of the mean ($a^* = 0.409$, $b^* = 0.343$, chroma = 0.493, hue angle = 0.612). a, b: Different letters within traits indicate statistically significant differences ($p < 0.05$).

Table 3. Effect of different packaging methods and storage temperatures on sensory properties of dry-aged beef during storage for 21 days.

Traits	Packaging method/storage temperature after dry aging	Storage days after dry aging				SEM ^y
		0	7	14	21	
Flavor	Vacuum 3°C	4.830	4.950	4.920	5.020	0.172
	Vacuum - 23°C	4.830	4.720	4.720	4.700	0.119
	SEM ^z	0.109	0.134	0.208	0.142	
Juiciness	Vacuum 3°C	4.850	5.100	4.950	4.910	0.212
	Vacuum - 23°C	4.850	4.840	4.390	4.610	0.265
	SEM ^z	0.179	0.325	0.192	0.269	
Tenderness	Vacuum 3°C	5.040	5.140	4.980	4.610	0.240
	Vacuum - 23°C	5.040	4.750	4.560	4.910	0.264
	SEM ^z	0.087	0.308	0.245	0.386	
Overall acceptance	Vacuum 3°C	4.950	4.950	4.800	4.440	0.256
	Vacuum - 23°C	4.950	4.860	4.500	4.560	0.219
	SEM ^z	0.132	0.277	0.223	0.344	

All values are mean \pm standard error of the mean ^yn = 24, ^zn = 12.

Conclusion

The results of this study indicate that dry-aged beef can be stored without, or with minimal, microbial growth and quality deterioration for different periods depending on the packaging method and storage temperature. With respect to microbial growth, it is recommended that dry-aged beef with wrap packaging stored in refrigerated conditions should be consumed as quickly as possible. Vacuum packaging can extend the color stability of displayed beef after dry aging over short periods (less than 14 days) of refrigerated storage without problems related to microbial spoilage, lipid oxidation, or sensory quality. For long-term storage, the results of this study suggest that dry-aged beef should be frozen, as freezing can extend the color stability up to day 21 of storage without adverse effects on hygienic or meat quality aspects of dry-aged beef.

Acknowledgements

This study was supported by “High Value-added Food Technology Development Program (Project No. 316048-03)”, Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries.

References

- Bellés M, Alonso V, Roncalés P, Beltrán JA. 2017. A revise of fresh lamb and preservation. *Small Ruminant Research* 146:41-47.
- DeGeer SL, Hunt MC, Bratcher CL, Crozier-Dodson BA, Johnson DE, Stika JF. 2006. Effects of dry aging of bone-in and boneless strip loins using two aging processes for two aging times. *Meat Science* 83:768-

774.

- Fernandes R, Freire MT, Paula E, Kanashiro A, Catunda F, Rosa A, Balieiro J, Trindade M. 2014. Stability of lamb loin stored under refrigeration and packed in different modified atmosphere packaging systems. *Meat Science* 96:554-561.
- Jakobson M, Bertelsen G. 2000. Colour stability and lipid oxidation of fresh beef. Development of a response surface model for predicting the effects of temperature, storage time, and modified atmosphere composition. *Meat Science* 54:49-57.
- Jayasingh P, Cornforth DP, Carpenter CE, Whittier D. 2001. Evaluation of carbon monoxide treatment in modified atmosphere packaging or vacuum packaging to increase color stability of fresh beef. *Meat Science* 59:317-324.
- Khan MI, Lee HJ, Kim HJ, Yong HI, Lee H, Jo C. 2016. Marination and physicochemical characteristics of vacuum-aged duck breast meat. *Asian Australasian Journal of Animal Science* 29:1639-1945.
- Ledward DA. 1985. Post-slaughter influences on the formation of metmyoglobin in beef muscles. *Meat Science* 15:149-171.
- Lee CW, Lee SH, Min Y, Lee S, Jo C, Jung S. 2015. Quality improvement of strip loin Hanwoo with low quality grade by dry aging. *The Korean Journal of Food and Nutrition* 28:415-421.
- Lee HJ, Choe J, Yoon JW, Kim SJ, Oh H, Yoon Y, Jo C. 2018. Determination of salable shelf-life for wrap-packaged dry-aged beef during cold storage. *Korean Journal for Food Science of Animal Resources* 38:251-258.
- Parrish FC, Boles JA, Rust RE, Oslon DG. 1991. Dry and wet aging effects on palatability attributes of beef loins and rib steaks from three quality grades. *Journal of Food Science* 56:601-603.
- Sitz BM, Calkins CR, Feuz DM, Umberger WJ, Eskridge KM. 2006. Consumer sensory acceptance and value of wet-aged and dry-aged beef steaks. *Journal of Animal Science* 84:1221-1226.
- Smith RD, Nicholson KL, Nicholson JDW, Harris KB, Miller RK, Griffin DB, Savell JW. 2008. Dry versus wet aging of beef: Retail cutting yields and consumer palatability evaluations of streaks from US choice and US select short loins. *Meat Science* 79:631-639.
- Sofos JN, Cabedo L, Zerby H, Belk KE, Smith GC. 2000. Potential interactions between antioxidants and microbial meat quality. pp. 427-453. In *Antioxidant in muscle foods* edited by Decker E, Faustman C, Lopez-Bote CJ. John Wiley & Sons Publishing, NY, USA.
- Warren KE, Kastner CL. 1992. A comparison of dry-aged and vacuum-aged beef strip loins. *Journal of Muscle Foods* 3:151-157.
- Yoon JW, Lee DG, Lee HJ, Choe JH, Jung S, Jo C. 2017. Microbial, physicochemical, and sensory characteristics of quality grade 2 beef enhanced by injection of pineapple concentrate and honey. *Korean Journal for Food Science of Animal Resources* 37:494-501.
- Younathan MT, Watts B. 1959. Relationship of meat pigments to lipid oxidation. *Food Research* 24:728-734.