

Short Communication

Walnut phenolic extracts reduce telomere length and telomerase activity in a colon cancer stem cell model

Phil-Kyung Shin¹, Yoonchae Zoh¹, Jina Choi¹, Myung-Sunny Kim², Yuri Kim³ and Sang-Woon Choi^{1,4,5}

¹CHA University School of Medicine, 120, Haeryong-ro, Pocheon-si, Gyeonggi 13488, Korea

²Division of Nutrition and Metabolism, Korea Food Research Institute, Jeonju, Jeonbuk 55365, Korea

³Department of Nutritional Science and Food Management, Ewha Womans University, Seoul, 03760, Korea

⁴Chaum Life Center CHA University, 442, Dosan-daero, Gangnam-gu, Seoul, 06062, Korea

BACKGROUND/OBJECTIVES: Telomeres are located at the chromosomal ends and progressively shortened during each cell cycle. Telomerase, which is regulated by *hTERT* and *c-MYC*, maintains telomeric DNA sequences. Especially, telomerase is active in cancer and stem cells to maintain telomere length for replicative immortality. Recently we reported that walnut phenolic extract (WPE) can reduce cell viability in a colon cancer stem cell (CSC) model. We, therefore, investigated the effect of WPE on telomere maintenance in the same model.

MATERIALS AND METHODS: CD133⁺CD44⁺ cells from HCT116, a human colon cancer cell line, were sorted by Fluorescence-activated cell sorting (FACS) and treated with WPE at the concentrations of 0, 10, 20, and 40 µg/mL for 6 days. Telomere lengths were assessed by quantitative real-time PCR (qRT-PCR) using telomere specific primers and DNA extracted from the cells, which was further adjusted with single-copy gene and reference DNA (*ddC_i*). Telomerase activity was also measured by qRT-PCR after incubating the PCR mixture with cell protein extracts, which was adjusted with reference DNA (*dC_i*). Transcriptions of *hTERT* and *c-MYC* were determined using conventional RT-PCR.

RESULTS: Telomere length of WPE-treated cells was significantly decreased in a dose-dependent manner (5.16 ± 0.13 at 0 µg/mL, 4.79 ± 0.12 at 10 µg/mL, 3.24 ± 0.08 at 20 µg/mL and 3.99 ± 0.09 at 40 µg/mL; $P = 0.0276$). Telomerase activities concurrently decreased with telomere length (1.47 ± 0.04 , 1.09 ± 0.01 , 0.76 ± 0.08 , and 0.88 ± 0.06 ; $P = 0.0067$). There was a positive correlation between telomere length and telomerase activity ($r = 0.9090$; $P < 0.0001$). Transcriptions of both *hTERT* and *c-MYC* were also significantly decreased in the same manner.

CONCLUSION: In the present cell culture model, WPE reduced telomere maintenance, which may provide a mechanistic link to the effect of walnuts on the viability of colon CSCs.

Nutrition Research and Practice 2019;13(1):58-63; <https://doi.org/10.4162/nrp.2019.13.1.58>; pISSN 1976-1457 eISSN 2005-6168

Keywords: Walnut, colonic neoplasms, stem cells, telomere, telomerase

INTRODUCTION

Nuts are rich in nutrients and bioactive food components that exhibit anti-cancer properties. Recent prospective studies have suggested the protective effects of nut consumption on colorectal cancer [1]. Walnuts (*Juglans regia* L.) are well known to have many health effects such as anti-inflammatory and anti-oxidant properties as well as cardioprotective effect [2-4]. Walnuts, in particular, contain multiple components such as omega-3 fatty acids, phytochemicals, fiber, phenolic compounds, folate, and vitamin E, each of which has been individually associated with reducing the risk of certain cancers. In a mouse study, a walnut-containing diet was shown to inhibit colorectal cancer growth by suppressing angiogenesis; previous observations

also demonstrated walnut extracts to have inhibitory effects on the growth of colon cancer cells [5]. Moreover, we have already demonstrated that walnut phenolic extracts (WPEs) can suppress the growth of colon cancer stem cells [6].

Telomeres, located at the chromosomal ends and getting shorter during each cell cycle, are crucial for chromosomal integrity and stability. Shortened telomere length has been associated with risks of cancers and age-associated diseases. In fact, maintenance of telomere length is essential for cell replication and tumorigenesis and telomere shortening is an initial event in colorectal carcinogenesis [7,8].

Telomerase, an RNA-dependent DNA polymerase that synthesizes telomeric DNA sequences, is induced for cell proliferation and inhibited for cell differentiation, indicating that the

This work was supported by research grants from the California Walnut Commission (S.W.C.) and Korean Food Research Institute (project no. E0150302-04) (S.W.C.). All the authors participated in the conception, design, and conduction of the study as well as interpreting the data and drafting the manuscript. P.K.S and Y.Z. also performed cell culture studies and measured endpoints. J.C., M.S.K., Y.K., and S.W.C supervised the study.

⁵ Corresponding Author: Sang-Woon Choi, Tel. 82-2-3015-5025, Fax. 82-2-3015-5315, Email. sang.choi@cha.ac.kr

Received: February 14, 2018, Revised: March 28, 2018, Accepted: September 19, 2018

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

(ACX), and protein extract from the cells. Primer sequences are shown in Table 1 [17]. The PCR reaction mixture was incubated at room temperature for 30 min to allow the telomerase in the protein extracts to elongate the TS primer by adding TTAGGG repeat sequences. Thereafter, qRT-PCR was conducted with the following PCR condition: 95°C incubation for 10 min followed by 40 cycles of 95°C for 20 sec and 50°C for 30 sec. All samples were run in triplicate. The HEK 293 cell line was used as a reference and telomerase activity (Mean±SE) was represented as dC_t .

hTERT and c-MYC transcriptions

Quantitation of mRNA expression was performed by reverse transcription polymerase chain reaction (RT-PCR). Total RNA was extracted using Nucleozol (MACHEREY-NAGEL GmbH & Co. KG, Postfach, Düren, Germany) from CD133⁺CD44⁺ HCT116 cells treated with various concentrations of WPE. The RNA was transcribed into cDNA using AccuPower[®] RT PreMix (Bioneer, Daejeon, Korea) and amplified using primers specific to each gene and PCR master mix (Bioneer, Daejeon, Korea). The PCR products were separated by 1.5% agarose gel electrophoresis. Each mRNA level was normalized with Ribosomal protein large subunit P 0 (*RPLP0*). All primers are shown in Table 1 [18-22].

Statistical analysis

Differences between the variables of the telomere length and telomerase activity at each concentration of WPE were tested

using ANOVA. Duncan's post hoc analysis using variance was also performed to compare the differences between the groups. Correlations between telomere length and telomerase activity were assessed using Spearman's correlation. $P < 0.05$ was regarded as significant. All statistical analyses were performed using SPSS version 24.0 (IBM Corporation, Armonk, NY, USA).

RESULTS

Telomere length in WPE-treated cells was significantly decreased in a dose-dependent manner up to the 40 $\mu\text{g/mL}$ concentration (5.16 ± 0.13 at 0 $\mu\text{g/mL}$, 4.79 ± 0.12 at 10 $\mu\text{g/mL}$, 3.24 ± 0.08 at 20 $\mu\text{g/mL}$, and 3.99 ± 0.09 at 40 $\mu\text{g/mL}$; $P = 0.0276$) (Fig. 1A). At 40 $\mu\text{g/mL}$, telomere length was still significantly lower than the control, but higher compared to the 20 $\mu\text{g/mL}$ concentration. Interestingly, telomerase activities concurrently decreased with telomere length up to the 40 $\mu\text{g/mL}$ concentration (1.47 ± 0.04 , 1.09 ± 0.01 , 0.76 ± 0.08 , and 0.88 ± 0.06 ; $P = 0.0067$) (Fig. 1B), even though no additive effect was observed at 40 $\mu\text{g/mL}$ compared to 20 $\mu\text{g/mL}$. Significant positive correlation was also observed between the telomere length and telomerase activity (Fig. 1C; $r = 0.9090$, $P < 0.0001$). The transcriptions of *hTERT* showed a significant stepwise decreasing pattern by stepwise increasing WPE concentrations (Fig. 2A; $P < 0.0001$). The transcription of *c-MYC*, which is known to involve in the transcription of *hTERT*, also showed a significant stepwise decreasing pattern by WPE (Fig. 2B; $P <$




Fig. 1. WPE reduced both telomere length and telomerase activity. (a) Relative telomere length (ddC_t) of HCT116 CD133⁺CD44⁺ cells with respect to the WPE concentration. One-way ANOVA was used to determine the association between telomere length and WPE. Values are represented as Mean±SE, $P < 0.05$. Significant differences are observed for values having bars with different letters, (b) Relative telomerase activity (dC_t) of HCT116 CD133⁺CD44⁺ cells with respect to the WPE concentration. One-way ANOVA was used to determine the association between telomerase activity and WPE. Values are represented as Mean±SE, $P < 0.05$. Significant differences are observed for values having bars with different letters, (c) Correlation between telomere length and telomerase activity. Correlation was assessed using Spearman's correlation. The correlation coefficient is observed to be $r = 0.9090$ and $P < 0.0001$.




Fig. 2. WPE reduced the transcriptions of *hTERT* and *c-MYC*. To determine the transcription of *hTERT* and *c-MYC* genes RT-PCR was conducted with *RPLPO* gene as an endogenous control, WPE downregulated the transcriptions of *hTERT* (a) and *c-MYC* (b) in a dose dependent manner. Values are represented as Mean \pm SE, $P < 0.05$ by one-way ANOVA. Significant differences are observed for values having bars with different letters.

0.0001). Collectively, WPE treatment simultaneously reduced the transcriptions of *c-MYC* and *hTERT* as well as telomerase activity and telomere length.

DISCUSSION

Telomere maintenance is a prerequisite for indefinite proliferation in cancer. Therefore, inhibiting the maintenance of telomeres may prove to be a useful strategy for cancer therapy. The telomere and telomerase connection can obviously offer several targets for inducing cancer cell death [23]. In the present study, we demonstrated that WPE reduced telomerase activity and telomere length as well as transcriptions of *hTERT* and *c-MYC*, both of which are known to control telomerase activity. It could be a mechanism by which walnuts inhibit the growth of colon cancer cells [6]. To the best of our knowledge,

this is the first study that attempts to determine the effect of walnuts on telomere maintenance.

Telomere length, which is crucial for individual cell senescence, is also highly associated with cancer. Individuals with short telomeres are known to have an increased risk of cancer because short telomeres lead to genomic instability; however, individuals with long telomeres paradoxically can have an increased risk of major cancers [24]. The association between the blood cell telomere length and cancer risk is also supported by a meta-analysis which demonstrated that ever smokers have shorter telomeres compared with never smokers. Furthermore, current smokers have shorter telomeres compared with never or former smokers [25]. In a systemic review that analyzed the association between tissue telomere length and breast cancer prognosis, longer telomeres were associated with better prognosis and shorter telomeres were associated with higher local recurrence rates and higher tumor grade [26]. Collectively, telomere length can be a good biomarker for the development and prognosis of cancer, suggesting that telomere and telomere metabolism can be a potential target for cancer treatment. We, therefore, investigated the effects of walnut on telomere length and telomerase activity as well as the expression of upstream genes using a colon cancer stem cell model in which walnut has shown to reduce the cell proliferation and viability [6].

Telomerase-targeted treatments for cancer have garnered a lot of attention because of the peculiar characteristics of telomerases, which are detected in nearly all cancer cells but not expressed in most normal cells [27]. Numerous studies have shown that the inhibition of telomerase induces cancer cell death. Sulforaphane, a bioactive food component found abundantly in broccoli and broccoli sprouts, can inhibit cell viability and induce apoptosis, simultaneously downregulating the protein expression and enzymatic activity of telomerase in colon cancer cell lines owing to its epigenetic effect as a histone deacetylase [28]. In a previous study, a certain form of L-asparaginases, which were used to treat acute lymphoblastic leukemia with some limitations, was found to suppress telomerase activity in human T-cell lymphoma Jurkat cells; this is considered to be an additional anticancer mechanism of these compounds, and inhibition of the telomerase activity may be considered to be a promising approach for cancer treatment [29]. It appears that the decreased telomerase activity demonstrated in this study can be a mechanism by which WPE reduces telomere length and thereby reduces cell viability. Interestingly, the positive correlation between telomere length and telomerase activity further supports this mechanism. A growing body of evidence has also demonstrated that telomere stability can be affected more than the main genome by conventional cancer chemotherapeutic agents, suggesting that cancer chemotherapies with genotoxic properties preferably target telomere [30].

Collectively, our observation that WPE reduced telomerase activity and telomere length represents the potential effect to assist cancer chemotherapy.

We also found that the repression of *hTERT* and *c-MYC* by WPE was in parallel with decreasing telomerase length and telomerase activity, suggesting that the effect of WPE on telomere maintenance could be conveyed through those genes.

Because *hTERT* is a catalytic subunit of the enzyme telomerase, it is not surprising that the *hTERT* expression was downregulated by WPE simultaneously with the reduction of telomerase activity. On the other hand, the observation that WPE reduced the expression of *c-MYC*, which can affect the expression of *hTERT*, is quite interesting, because *c-MYC* plays many other important roles in cancer including the transformation of normal human cells into cancer cells [11]. The *c-MYC* transcription factor is involved in cellular growth, proliferation and metabolism by controlling the expression of numerous target genes [31]. It suggests that walnut may have another anticancer property through the inhibition of *c-MYC* transcription.

In conclusion, WPE demonstrates the ability to disrupt telomere maintenance in a colon cancer stem cells model. Further studies are warranted to determine the cause and effective relationship between the inhibition of telomere maintenance by WPE and the colon cancer stem cell viability.

CONFLICT OF INTEREST

The authors declare no potential conflicts of interests.

ORCID

Phil-Kyung Shin: <https://orcid.org/0000-0002-3935-3132>

Yoonchae Zoh: <https://orcid.org/0000-0002-0707-3523>

Jina Choi: <https://orcid.org/0000-0002-7297-4256>

Myung-Sunny Kim: <https://orcid.org/0000-0002-5020-3397>

Yuri Kim: <https://orcid.org/0000-0001-7606-8501>

Sang-Woon Choi: <https://orcid.org/0000-0003-0713-379X>

REFERENCES

- Sabaté J, Ang Y. Nuts and health outcomes: new epidemiologic evidence. *Am J Clin Nutr* 2009;89:1643S-1648S.
- Yu Z, Malik VS, Keum N, Hu FB, Giovannucci EL, Stampfer MJ, Willett WC, Fuchs CS, Bao Y. Associations between nut consumption and inflammatory biomarkers. *Am J Clin Nutr* 2016;104:722-8.
- Calcabrini C, De Bellis R, Mancini U, Cucchiari L, Stocchi V, Potenza L. Protective effect of *Juglans regia* L. walnut extract against oxidative DNA damage. *Plant Foods Hum Nutr* 2017;72:192-7.
- Feldman EB. The scientific evidence for a beneficial health relationship between walnuts and coronary heart disease. *J Nutr* 2002;132:1062S-1101S.
- Tsoukas MA, Ko BJ, Witte TR, Dincer F, Hardman WE, Mantzoros CS. Dietary walnut suppression of colorectal cancer in mice: mediation by miRNA patterns and fatty acid incorporation. *J Nutr Biochem* 2015;26:776-83.
- Lee J, Kim YS, Lee J, Heo SC, Lee KL, Choi SW, Kim Y. Walnut phenolic extract and its bioactive compounds suppress colon cancer cell growth by regulating colon cancer stemness. *Nutrients* 2016;8:E439.
- Rampazzo E, Bertorelle R, Serra L, Terrin L, Candiottio C, Pucciarelli S, Del Bianco P, Nitti D, De Rossi A. Relationship between telomere shortening, genetic instability, and site of tumour origin in colorectal cancers. *Br J Cancer* 2010;102:1300-5.
- Zhang C, Chen X, Li L, Zhou Y, Wang C, Hou S. The association between telomere length and cancer prognosis: evidence from a meta-analysis. *PLoS One* 2015;10:e0133174.
- Zhao Y, Cheng D, Wang S, Zhu J. Dual roles of c-Myc in the regulation of *hTERT* gene. *Nucleic Acids Res* 2014;42:10385-98.
- Rana C, Piplani H, Vaish V, Nehru B, Sanyal SN. Downregulation of telomerase activity by diclofenac and curcumin is associated with cell cycle arrest and induction of apoptosis in colon cancer. *Tumour Biol* 2015;36:5999-6010.
- Leão R, Apolónio JD, Lee D, Figueiredo A, Tabori U, Castelo-Branco P. Mechanisms of human telomerase reverse transcriptase (*hTERT*) regulation: clinical impacts in cancer. *J Biomed Sci* 2018;25:22.
- Miller DM, Thomas SD, Islam A, Muench D, Sedoris K. c-Myc and cancer metabolism. *Clin Cancer Res* 2012;18:5546-53.
- Choi J, Shin PK, Kim Y, Hong CP, Choi SW. Metabolic influence of walnut phenolic extract on mitochondria in a colon cancer stem cell model. *Eur J Nutr*. Forthcoming 2018.
- Anderson KJ, Teuber SS, Gobeille A, Cremin P, Waterhouse AL, Steinberg FM. Walnut polyphenolics inhibit *in vitro* human plasma and LDL oxidation. *J Nutr* 2001;131:2837-42.
- Min SJ, Lim JY, Kim HR, Kim SJ, Kim Y. Sasa quelpaertensis leaf extract inhibits colon cancer by regulating cancer cell stemness *in vitro* and *in vivo*. *Int J Mol Sci* 2015;16:9976-97.
- Cawthon RM. Telomere measurement by quantitative PCR. *Nucleic Acids Res* 2002;30:e47.
- Hou M, Xu D, Björkholm M, Gruber A. Real-time quantitative telomeric repeat amplification protocol assay for the detection of telomerase activity. *Clin Chem* 2001;47:519-24.
- Sadr M, Noori Mugahi SM, Hassanzadeh G, Nadj SA, Kiani A, Abedini A, Javadi A, Mohammadi F, Masjedi MR, Bahadori M. Telomere shortening in blood leukocytes of patients with chronic obstructive pulmonary disease. *Tanaffos* 2015;14:10-6.
- Herbert BS, Hochreiter AE, Wright WE, Shay JW. Nonradioactive detection of telomerase activity using the telomeric repeat amplification protocol. *Nat Protoc* 2006;1:1583-90.
- Liu T, Liang X, Li B, Björkholm M, Jia J, Xu D. Telomerase reverse transcriptase inhibition stimulates cyclooxygenase 2 expression in cancer cells and synergizes with celecoxib to exert anti-cancer effects. *Br J Cancer* 2013;108:2272-80.
- Duangmano S, Dakeng S, Jiratchariyakul W, Suksamrarn A, Smith DR, Patmasiriwat P. Antiproliferative effects of cucurbitacin B in breast cancer cells: down-regulation of the c-Myc/*hTERT*/telomerase pathway and obstruction of the cell cycle. *Int J Mol Sci* 2010;11:5323-38.
- Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. *Methods Mol Biol* 2000;132:365-86.
- Gilley D, Tanaka H, Herbert BS. Telomere dysfunction in aging and cancer. *Int J Biochem Cell Biol* 2005;37:1000-13.
- Aviv A, Anderson JJ, Shay JW. Mutations, cancer and the telomere length paradox. *Trends Cancer* 2017;3:253-8.
- Astuti Y, Wardhana A, Watkins J, Wulaningsih W, Network PR, PILAR Research Network. Cigarette smoking and telomere length: a systematic review of 84 studies and meta-analysis. *Environ Res* 2017;158:480-9.
- Ennour-Idrissi K, Maunsell E, Diorio C. Telomere length and breast cancer prognosis: a systematic review. *Cancer Epidemiol Biomarkers Prev* 2017;26:3-10.
- Fang X, Hu T, Yin H, Yang J, Tang W, Hu S, Xu X. Differences in telomerase activity and the effects of AZT in aneuploid and euploid cells in colon cancer. *Int J Oncol* 2017;51:525-32.
- Martin SL, Kala R, Tollefsbol TO. Mechanisms for the inhibition of

- colon cancer cells by sulforaphane through epigenetic modulation of microRNA-21 and human telomerase reverse transcriptase (hTERT) down-regulation. *Curr Cancer Drug Targets* 2018;18:97-106.
29. Zhdanov DD, Pokrovsky VS, Pokrovskaya MV, Alexandrova SS, Eldarov MA, Grishin DV, Basharov MM, Gladilina YA, Podobed OV, Sokolov NN. Inhibition of telomerase activity and induction of apoptosis by rhodospirillum rubrum L-asparaginase in cancer Jurkat cell line and normal human CD4+ T lymphocytes. *Cancer Med* 2017;6:2697-712.
30. Lu Y, Leong W, Guérin O, Gilson E, Ye J. Telomeric impact of conventional chemotherapy. *Front Med* 2013;7:411-7.
31. Marfil V, Blazquez M, Serrano F, Castell JV, Bort R. Growth-promoting and tumorigenic activity of c-Myc is suppressed by Hhex. *Oncogene* 2015;34:3011-22.