

# Infection Status with *Clonorchis sinensis* Metacercariae in Fish from Yangcheon (Stream) in Sancheong-gun, Gyeongsangnam-do, Korea

Woon-Mok Sohn<sup>1,\*</sup>, Byoung-Kuk Na<sup>1</sup>, Shin-Hyeong Cho<sup>2</sup>, Jung-Won Ju<sup>2</sup>

<sup>1</sup>Department of Parasitology and Tropical Medicine, and Institute of Health Sciences, Gyeongsang National University College of Medicine, Jinju 52727, Korea; <sup>2</sup>Division of Vectors and Parasitic Diseases, Centers for Disease Control and Prevention, Osong 28159, Korea

**Abstract:** The infection status with *Clonorchis sinensis* metacercariae (CsMc) was examined in freshwater fishes from Yangcheon (a branch of Gyeongho-gang), which is located in Sancheong-gun, Gyeongsangnam-do, the Republic of Korea. Total 2,201 fishes in 26 species were examined by the artificial digestion method through 7 years. CsMc were detected in 1,171 (53.2%) fishes in 21 spp. (80.8%) and their density was 85 per fish infected. Total 532 (99.6%) out of 534 *Pungtungia herzi* (index fish) examined were infected with 147 CsMc per fish infected. Metacercarial densities in this fish were highest in 2015 (179 CsMc), followed by 2012 (168), 2013 (152), 2016 (145), 2014 (114), and 2017 (89) respectively. In the gobioninid fish group, i.e., *P. herzi*, *Sarcocheilichthys* spp., *Squalidus* spp., *Pseudogobio esocinus*, *Hemibarbus longirostris*, and *Hemibarbus labeo*, 841 (92.7%) fishes were infected with 117 CsMc per fish infected. Total 250 (54.7%) acheilognathinid fish (bitterlings), *Acheilognathus* spp. and *Acanthorhodeus* spp. were infected with 5.8 CsMc. In the rasborinid fish (chubs) group, i.e., *Zacco platypus*, *Zacco temminckii*, and *Zacco koreanus*, 77 (13.7%) out of 563 fish examined were infected with 2.4 CsMc in average. The susceptibility indices of CsMc were 49.09 in the overall positive fish group, 104.15 in the gobioninid group, 3.17 in the acheilognathinid group and 0.35 in the rasborinid fish group respectively. Only 1 CsMc was detected in 3 fish species, *Coreoperca herzi*, *Channa argus*, and *Lepomis macrochirus*, respectively. Conclusively, it was confirmed that CsMc are moderately prevalent in fishes from Yangcheon in Sancheong-gun, Gyeongsangnam-do, Korea.

**Key words:** *Clonorchis sinensis*, metacercaria, susceptibility index, Gobioninae, Acheilognathinae, Rasborinae, Cyprinidae, fish host, Yangcheon

## INTRODUCTION

Clonorchiasis, *Clonorchis sinensis* (Digenea: Opisthorchiidae) infection, is most important among endemic parasitic diseases in the Republic of Korea (Korea). Although the prevalence of this endemic disease was 1.86% in the nationwide survey on the helminthic infection in Korea, about 932,540 Korean people are estimated to be infected with this endemic trematode. Nowadays, it is the highest value among the prevalences of parasitic diseases in Korea [1]. The prevalence of clonorchiasis has maintained at relatively high levels in the residents of riverside areas in Korea [2-6]. Recently, a team of Korean CDC (Division of Vectors and Parasitic Diseases, Centers for Disease

Control and Prevention) [4-6] reported the prevalences of clonorchiasis in the adjacent residents of 5 major rivers, i.e., Nakdong-gang (gang means river), Seomjin-gang, Geum-gang, Yeongsan-gang and Han-gang, in Korea. The riverside area of Gyeongho-gang (a branch stream of Nakdong-gang) in Sancheong-gun, Gyeongsangnam-do has been known as a high endemic area of clonorchiasis [6-10].

Many Korean workers epidemiologically surveyed the freshwater fishes, the human infection sources, to estimate the endemicities of clonorchiasis [11-17]. Especially, Kim et al. [11] widely surveyed freshwater fishes from 34 localities to know the infection status with *Clonorchis sinensis* metacercariae (CsMc) in Korea. Cho et al. [12] investigated the infection status of CsMc in freshwater fish from 3 wide regions, which were tentatively divided by the latitudinal levels of Korean peninsula. Cho et al. [13] also surveyed on the prevalence of zoonotic trematode metacercariae in freshwater fish from Gangwon-do (do = Province), Korea. Sohn et al. [14] investigated the infection status of digenetic trematode metacercariae including *C.*

•Received 14 March 2019, accepted 25 March 2019.

\*Corresponding author (wmsohn@gnu.ac.kr)

© 2019, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

*sinensis* in freshwater fish from the water systems of Hantangan and Imjingang located in relatively northern regions of Korea. Recently, Sohn et al. [15] and Yoon et al. [17] surveyed the prevalence of CsMc in freshwater fishes from the water systems of Seomjin-gang and Tamjin-gang. Sohn et al. [16] also reported the prevalence and intensity of CsMc in freshwater fish from a highly endemic site, Wicheon (a branch stream of Nakdonggang) (cheon means stream), in Gunwi-gun, Gyeongsangbuk-do, Korea.

Gyeongho-gang is one of the branch stream of Nakdonggang, which rise from a mountainous area (Namdeokyu-san) (san means mountain) of Seosang-myeon (myeon = township) in Hamyang-gun (gun = county), flows via Hamyang-gun and Sancheong-gun and into Jinyang-ho (ho means lake) in Jinju-si (si = city), Gyeongsangnam-do. Yangcheon is one of the branch stream of Gyeongho-gang located in the Saengbiryang-myeon, Sancheong-gun [18]. The riverside area of Yangcheon has been reported as a high endemic area of clonorchiasis [7-9]. However, the infection status with CsMc in fish from this area has not been widely and systematically examined yet. Therefore, we intended to investigate the infection status with CsMc in fishes from Yangcheon for 7 years (2011-2017).

## MATERIALS AND METHODS

### Fish collection site and freshwater fishes examined

We collected total 2,201 freshwater fishes in 26 species in Yangcheon (a branch stream of Gyeongho-gang), which is located in Saengbiryang-myeon, Sancheong-gun (Latitude: 35.37015; Longitude: 128.08025), Gyeongsangnam-do, Korea. The numbers and species of fish by the year examined were as follows.

Total 201 freshwater fish in 14 species were examined in 2011. Fish species (No. of fish) examined were *Zacco temminckii* (27), *Coreoperca herzi* (10), *Odontobutis platycephala* (14), and *Carassius auratus* (1) including 10 ones with CsMc in Table 1. Total 221 freshwater fish in 11 species were examined in 2012. Fish species (No. of fish) examined were *Zacco platypus* (29), *C. herzi* (4), and *C. auratus* (1) including 8 ones with CsMc in Table 1. Total 644 freshwater fish in 20 species were examined in 2013. Fish species (No. of fish) examined were *C. auratus* (20), *C. herzi* (19), *Liobagrus mediadiposalis* (3), *O. platycephala* (3), *Acheilognathus yamatsutae* (1), and *Z. temminckii* (1) including 14 ones with CsMc in Table 1.

Total 291 freshwater fish in 14 species were examined in

2014. Fish species (No. of fish) examined were *O. platycephala* (13) and *Siniperca scherzeri* (1) including 12 ones with CsMc in Table 1. Total 183 freshwater fish in 15 species were examined in 2015. Fish species (No. of fish) examined were *Z. temminckii* (40), *C. herzi* (2), *O. platycephala* (1), *Micropterus salmoides* (1), *C. auratus* (1), and *A. macropterus* (1) including 9 ones with CsMc in Table 1.

Total 253 freshwater fish in 16 species were examined in 2016. Fish species (No. of fish) examined were *C. auratus* (22), *Z. temminckii* (21), *Z. koreanus* (20), *C. herzi* (10), *O. platycephala* (1), and *A. macropterus* (1) including 10 ones with CsMc in Table 1. Total 408 fish in 22 species were examined in 2017. Fish species (No. of fish) examined were *M. salmoides* (36), *C. auratus* (28), *Z. koreanus* (20), *C. herzi* (18), and *O. platycephala* (4) including 17 ones with CsMc in Table 1.

### Examination methods

All collected fishes were transferred to the laboratory of the Department of Parasitology and Tropical Medicine, Gyeongsang National University College of Medicine, Jinju, Korea. After the identification of fish species, they were individually ground with a mortar or grinder. Each ground fish meat was mixed with artificial gastric juice and the mixture was incubated at 36°C for about 2 hr. The digested material was filtered with 1 × 1 mm of mesh and washed with 0.85% saline until the supernatant is clear. The sediment was carefully examined under a stereomicroscope. The metacercariae of *C. sinensis* (CsMc) were separately collected by the general feature [19,20], and they were counted to get hold of infection rates (No. of fish with CsMc/No. of fish examined × 100) and densities (No. of CsMc/a fish infected) by fish species. The susceptibility indices of CsMc in each fish species were calculated by the formula, prevalence/100 × mean metacercarial density per fish infected.

## RESULTS

### Infection status with CsMc in overall fishes

The metacercariae of *C. sinensis* were detected in 1,171 (53.2%) out of 2,201 fishes in 26 species examined, and their average density was 85 per fish infected. In CsMc positive fish species, the number of fish examined was 1,827 (83.0%) in 21 species (80.8%), and the positive rate was 64.1%. The infection status by the fish species and surveyed years was detailedly shown in Table 1.

**Table 1.** Infection status of *Clonorchis sinensis* metacercariae in fishes from Yangcheon (a stream of Gyeongho-gang) in Sancheong-gun, Gyeongsangnam-do

Year and fish sp.	No. of fish examined	No. of fish infected (%)	No. of CsMc detected	
			Range	Average
<b>2011</b>				
<i>Pungtungia herzi</i>	95	95 (100)	4-930	157.2
<i>Zacco platypus</i>	16	1 (6.3)	-	1.0
<i>Squalidus chankaensis</i>	13	13 (100)	14-322	95.1
<i>Squalidus japonicus coreanus</i>	12	12 (100)	7-69	23.3
<i>Pseudogobio esocinus</i>	4	4 (100)	11-167	57.3
<i>Hemibarbus longirostris</i>	4	3 (75.0)	-	1.0
<i>Sarcocheilichthys nigripinnis</i>	2	2 (100)	43-96	69.5
<i>Hemibarbus labeo</i>	1	1 (100)	-	6.0
<i>Sarcocheilichthys variegatus</i>	1	1 (100)	-	512.0
<i>Acheilognathus majusculus</i>	1	1 (100)	-	2.0
Subtotal	149	133 (89.3)	1-930	130.4
<b>2012</b>				
<i>Pungtungia herzi</i>	106	106 (100)	2-1,157	168.0
<i>Zacco temminckii</i>	35	3 (8.6)	1-2	1.3
<i>Hemibarbus longirostris</i>	21	5 (23.8)	1-2	1.4
<i>Sarcocheilichthys nigripinnis</i>	8	8 (100)	7-795	205.1
<i>Acheilognathus majusculus</i>	7	2 (28.6)	7-12	9.5
<i>Squalidus chankaensis</i>	5	5 (100)	18-58	42.6
<i>Acanthorhodeus macropterus</i>	4	1 (25.0)	-	1.0
<i>Squalidus gracilis majimae</i>	1	1 (100)	-	7.0
Subtotal	187	131 (70.1)	1-1,157	150.4
<b>2013</b>				
<i>Pungtungia herzi</i>	142	141 (99.3)	11-742	152.4
<i>Zacco koreanus</i>	94	2 (2.1)	-	1.0
<i>Acheilognathus majusculus</i>	85	40 (47.1)	1-10	2.5
<i>Zacco platypus</i>	55	19 (34.5)	1-13	2.3
<i>Squalidus chankaensis</i>	51	51 (100)	7-183	61.3
<i>Hemibarbus longirostris</i>	34	19 (55.9)	1-3	1.6
<i>Acheilognathus koreensis</i>	31	25 (80.6)	1-20	5.3
<i>Squalidus japonicus coreanus</i>	30	30 (100)	1-134	33.7
<i>Pseudogobio esocinus</i>	20	18 (90.0)	2-23	7.8
<i>Acanthorhodeus gracilis</i>	20	3 (15.0)	-	1.0
<i>Acanthorhodeus macropterus</i>	17	3 (17.6)	-	1.0
<i>Sarcocheilichthys variegatus</i>	8	8 (100)	56-530	227.9
<i>Acheilognathus rhombeus</i>	8	5 (62.5)	1-5	2.6
<i>Sarcocheilichthys nigripinnis</i>	2	2 (100)	21-153	87.0
Subtotal	597	366 (61.3)	1-742	76.8
<b>2014</b>				
<i>Pungtungia herzi</i>	65	65 (100)	1-443	114.3
<i>Acheilognathus majusculus</i>	40	9 (22.5)	1-8	2.9
<i>Zacco koreanus</i>	40	1 (2.5)	-	1.0
<i>Coreoperca herzi</i>	30	1 (3.3)	-	1.0
<i>Zacco platypus</i>	25	13 (52.0)	1-10	2.8
<i>Squalidus japonicus coreanus</i>	25	25 (100)	9-195	86.7
<i>Pseudogobio esocinus</i>	16	16 (100)	1-84	36.8
<i>Acheilognathus koreensis</i>	13	9 (69.2)	1-7	3.2
<i>Acheilognathus yamatsutae</i>	10	6 (60.0)	1-7	2.7
<i>Hemibarbus longirostris</i>	7	5 (71.4)	1-12	5.8
<i>Sarcocheilichthys variegatus</i>	5	5 (100)	2-13	8.0
<i>Sarcocheilichthys nigripinnis</i>	1	1 (100)	-	29.0
Subtotal	277	156 (56.3)	1-443	66.6

(Continued to the next page)

Infection status with CsMc in index fish, *Pungtungia herzi*  
CsMc were detected in 532 (99.6%) out of 534 *P. herzi* examined and their average density was 147 per fish infected.

The densities were most high in 2015 (179), and followed by 2012 (168), 2011 (157), 2013 (152), 2016 (145), 2014 (114), and 2017 (89). The infection status with CsMc in *P. herzi* by

Table 1. Continued

Year and fish sp.	No. of fish examined	No. of fish infected (%)	No. of CsMc detected	
			Range	Average
2015				
<i>Pungtungia herzi</i>	40	40 (100)	9-434	178.8
<i>Zacco platypus</i>	40	18 (45.0)	1-10	2.6
<i>Hemibarbus longirostris</i>	25	2 (8.0)	-	1.0
<i>Sarcocheilichthys variegatus</i>	10	10 (100)	42-362	224.3
<i>Acheilognathus majusculus</i>	9	3 (33.3)	2-6	3.3
<i>Acheilognathus koreensis</i>	7	6 (85.7)	1-5	2.0
<i>Acheilognathus yamatsutae</i>	3	3 (100)	1-5	2.3
<i>Squalidus japonicus coreanus</i>	2	2 (100)	98-194	146.0
<i>Pseudogobio esocinus</i>	1	1 (100)	-	12.0
Subtotal	137	85 (62.0)	1-434	115.0
2016				
<i>Acheilognathus koreensis</i>	40	25 (62.5)	1-9	2.9
<i>Pungtungia herzi</i>	33	33 (100)	23-341	144.8
<i>Zacco platypus</i>	32	14 (43.8)	1-7	2.8
<i>Acheilognathus majusculus</i>	23	13 (56.5)	1-16	2.8
<i>Acheilognathus rhombeus</i>	20	19 (95.0)	6-51	23.1
<i>Acheilognathus yamatsutae</i>	13	9 (69.2)	1-5	1.4
<i>Channa argus</i>	6	1 (16.7)	-	1.0
<i>Squalidus japonicus coreanus</i>	5	5 (100)	25-162	97.0
<i>Hemibarbus longirostris</i>	4	3 (75.0)	1-8	3.7
<i>Pseudogobio esocinus</i>	2	2 (100)	28-35	31.5
Subtotal	277	156 (56.3)	1-443	66.6
2017				
<i>Pungtungia herzi</i>	53	52 (98.1)	6-451	89.0
<i>Zacco platypus</i>	38	5 (13.2)	1-4	2.2
<i>Acheilognathus majusculus</i>	37	18 (48.6)	1-10	2.4
<i>Acheilognathus yamatsutae</i>	31	21 (67.7)	1-12	3.3
<i>Zacco temminckii</i>	30	1 (3.3)	-	1.0
<i>Lepomis macrochirus</i>	25	1 (4.0)	-	1.0
<i>Acanthorhodeus gracilis</i>	21	18 (85.7)	1-59	17.1
<i>Pseudogobio esocinus</i>	18	18 (100)	1-124	34.2
<i>Squalidus japonicus coreanus</i>	15	15 (100)	15-364	133.7
<i>Hemibarbus labeo</i>	8	6 (75.0)	1-6	2.3
<i>Hemibarbus longirostris</i>	7	5 (71.4)	1-8	2.4
<i>Acheilognathus rhombeus</i>	7	6 (85.7)	1-34	11.8
<i>Acheilognathus koreensis</i>	5	3 (60.0)	1-3	2.3
<i>Squalidus gracilis majimae</i>	3	3 (100)	32-195	86.7
<i>Acanthorhodeus macropterus</i>	2	2 (100)	4-7	5.5
<i>Sarcocheilichthys nigripinnis</i>	1	1 (100)	-	172.0
<i>Sarcocheilichthys variegatus</i>	1	1 (100)	-	208.0
Subtotal	302	176 (58.3)	1-451	48.0
Total	1,827	1,171 (64.1)	1-1,157	85.1

the year examined was detailedly revealed in Table 2.

#### Infection status with CsMc in gobioninid fish group

CsMc were detected in 841 (92.7%) out of 907 fishes in 9 species including *P. herzi* and their average density was 117 per fish infected. The densities were most high in *Sarcocheilichthys* spp. (179), and followed by *P. herzi* (147), *Squalidus* spp. (68), *P. esocinus* (28), *H. labeo* (3), and *H. longirostris* (2). The infection status by the fish species in this group was detailedly shown in Table 3.

#### Infection status with CsMc in acheilognathinid fish group

CsMc were detected in 250 (54.7%) out of 457 fishes examined and their average density was 5.8 per fish infected. The infection status with CsMc by the fish species, i.e., *A. majusculus*, *A. koreensis*, *A. yamatsutae*, *A. gracilis*, *A. rhombeus*, and *A. macropterus*, was detailedly revealed in Table 4.

#### Infection status with CsMc in rasborinid fish group

CsMc were detected in 77 (13.7%) out of 563 fishes examined and their average density was 2.4 per fish infected. The in-

**Table 2.** Infection status of *Clonorchis sinensis* metacercariae in index fish (striped shinner), *Pungtungia herzi*, from Yangcheon in Sancheong-gun, Gyeongsangnam-do

No. of examined	No. of fish examined	No. of fish infected (%)	No. of CsMc detected	
			Range	Average
2011	95	95 (100)	4-930	157.2
2012	106	106 (100)	2-1,157	168.0
2013	142	141 (99.3)	11-742	152.4
2014	65	65 (100)	1-443	114.3
2015	40	40 (100)	9-434	178.8
2016	33	33 (100)	23-341	144.8
2017	53	52 (98.1)	6-451	89.0
Total	534	532 (99.6)	1-1,157	147.0

**Table 3.** Infection status of *Clonorchis sinensis* metacercariae by the fish species in the gobioninid group (subfamily Gobioninae)

Species of fish	No. of fish examined	No. of fish infected (%)	No. of CsMc detected	
			Range	Average
<i>Pungtungia herzi</i>	534	532 (99.6)	1-1,157	147.0
<i>Sarcocheilichthys</i> spp.	39	39 (100)	2-795	179.0
<i>S. variegatus wakiyae</i>	25	25 (100)	2-530	193.0
<i>S. nigripinnis morii</i>	14	14 (100)	7-795	153.9
<i>Squalidus</i> spp.	162	162 (100)	1-364	68.4
<i>S. japonicus coreanus</i>	89	89 (100)	1-364	70.1
<i>S. chankaensis tsuchigae</i>	69	69 (100)	7-322	66.3
<i>S. gracilis majimae</i>	4	4 (100)	7-195	66.8
<i>Pseudogobio esocinus</i>	61	59 (96.7)	1-124	27.9
<i>Hemibarbus longirostris</i>	102	42 (41.2)	1-12	2.2
<i>Hemibarbus labeo</i>	9	7 (77.8)	1-6	2.9
Total	907	841 (92.7)	1-1,157	116.6

**Table 4.** Infection status of *Clonorchis sinensis* metacercariae by the fish species in the acheilognathinid group (subfamily Acheilognathinae)

Species of fish	No. of fish examined	No. of fish infected (%)	No. of CsMc detected	
			Range	Average
<i>Acheilognathus majusculus</i>	202	86 (42.6)	1-16	2.8
<i>Acheilognathus koreensis</i>	96	68 (70.8)	1-20	3.7
<i>Acheilognathus yamatsutae</i>	58	39 (67.2)	1-12	2.7
<i>Acanthorhodeus gracilis</i>	41	21 (51.2)	1-59	14.8
<i>Acheilognathus rhombeus</i>	35	30 (85.7)	1-51	17.4
<i>Acanthorhodeus macropterus</i>	25	6 (24.0)	1-7	2.5
Total	457	250 (54.7)	1-59	5.8

**Table 5.** Infection status of *Clonorchis sinensis* metacercariae by the fish species in the rasborinid group (subfamily Rasborinae)

Species of fish	No. of fish examined	No. of fish infected (%)	No. of CsMc detected	
			Range	Average
<i>Zacco platypus</i>	235	70 (29.8)	1-13	2.5
<i>Zacco koreanus</i>	174	3 (1.7)	-	1.0
<i>Zacco temminckii</i>	154	4 (2.6)	1-2	1.3
Total	563	77 (13.7)	1-13	2.4

fection status with CsMc by the fish species, i.e., *Z. platypus*, *Z. koreanus*, and *Z. temminckii*, was detailedly shown in Table 5.

#### Susceptibility index of CsMc by the fish groups

The susceptibility indices of CsMc were 54.6 in the overall



positive fish group, 108.1 in the gobioninid group, 3.2 in the acheilognathinid, and 0.3 in the rasborinid fish group respectively.

## DISCUSSION

By the present study, it was confirmed that CsMc are moderately prevalent in fishes from Yangcheon, in Sancheong-gun, Gyeongsangnam-do, Korea. The infection status was showed with a certain tendency by the subfamily groups, i.e., Gobioninae, Acheilognathinae and Rasborinae, in the cyprinid fish (Family Cyprinidae) hosts of *C. sinensis* like Sohn et al. [16]. The prevalences were 92.7%, 54.7%, and 13.7%, and metacercarial densities were 116.6, 5.8, and 2.4 per fish infected in 3 fish groups respectively. When we compared with those of Sohn et al. [16], prevalences (100%, 79.7%, and 35.5%) and metacercarial densities (1,310, 50, and 15) were much lower in this study. However, we also knew that the endemicity of CsMc is closely related with the subfamily groups in the cyprinid fish hosts of *C. sinensis* from a moderately endemic area, Yangcheon, in Sancheong-gun, Gyeongsangnam-do, Korea.

The water ecosystem of Yancheon is more or less healthy but the ecological conditions for fish is not so good. Total 2,201 freshwater fishes in 26 species were collected through 7 years (2011-2017) in the same site of Saengbiryang-myeon in Sancheong-gun, Gyeongsangnam-do, in this study. Among them, striped shinner (534 *P. herzi*: 24.3%), was the most dominant fish species like Yoon et al. [17] in Tamjingang, and followed by pale chub (235 *Z. platypus*: 10.7%), large striped bitterling (202 *A. majusculus*: 9.2%), Korean chub (174 *Z. koreanus*: 7.9%), dark chub (154 *Z. temminckii*: 7.0%), long nose barbel (102 *H. longirostris*: 4.6%), and oily bitterling (96 *A. koreensis*: 4.4%). The number of fish examined was 1,497 (68.0%) in major 7 species and 704 (32.0%) in remain 19 species. The disproportion of fish number examined is suggested that the ecological conditions for fish was relatively not so good, and because of all fishes were collected only by the netting in the nighttime. If we used together with other methods for fish catching like a casting net, the more various species of fish including diurnal ones were able to collect.

Total 51 fish species (in 36 genera 9 families) have been reported as the second intermediate hosts of *C. sinensis* in Korea [11,12,15-17,19]. In the present study, CsMc were found in 21 fish species, i.e., *P. herzi*, *S. variegatus wakiyae*, *S. nigripinnis morii*, *S. japonicus coreanus*, *S. chankaensis tsuchigae*, *S. gracilis maji-*

*mae*, *P. esocinus*, *H. longirostris*, *H. labeo*, *A. majusculus*, *A. koreensis*, *A. yamatsutae*, *A. gracilis*, *A. rhombeus*, *A. macropterus*, *Z. platypus*, *Z. koreanus*, *Z. temminckii*, *C. herzi*, *Channa argus*, and *Lepomis macrochirus*. Among them, 2 fish species, i.e., *L. macrochirus* (Centrarchidae) and *C. argus* (Channidae), are to be newly added in the list of the second intermediate hosts of *C. sinensis* in Korea. Accordingly, total 53 fish species in 38 genera (10 families) are to be the second intermediate hosts of *C. sinensis* in Korea.

The blue gill, *L. macrochirus*, was imported from Japan in 1969 and stocked in the Korean ecosystems in 1975 as the edible fish species. However, this predatory fish was widely spread in the water systems of whole country, and then specified as an agitating fish species of ecosystem in Korea [21]. In this study, no CsMc were detected in 5 (19.2%) out of 26 fish species examined, i.e., *Carassius auratus* (n=73), *Micropterus salmoides* (37), *Odontobutis platycephala* (36), *Liobagrus mediadiposalis* (3), *Siniperca scherzeri* (1), and only 1 CsMc was found in 1 (4.0%) out of 25 blue gills examined. Infections with CsMc in 2 exotic fish species, *L. macrochirus* and *M. salmoides* [16], are meaningful, although the number of fish examined and metacercariae detected were not so many. On the other hand, Choe et al. [22] did not found any other zoonotic trematode metacercariae including *C. sinensis* in 107 large mouth bass (*M. salmoides*) and 244 blue gills (*L. macrochirus*) from 2 sites, Daechong-ho and Musimcheon, of Chungcheongbuk-do, Korea. However, we should pay attention to the exotic fish species, such as large mouth bass and blue gill, in the metacercarial survey for the fishborne zoonotic trematodes.

Fish species edible in the raw, i.e., Mandarin fish (*S. scherzeri*), Korean aucha perch (*C. herzi*), dark sleeper (*Odontobutis* spp.), common carp (*Cyprinus carpio*), and crucian carp (*C. auratus*), practically act as the infection source of clonorchiasis in Korea. Fortunately, these fish species are less prevalent with CsMc. In even such a highly endemic area, Wicheon, 3 (8.8%) out of 34 *C. auratus* were infected with total 3 CsMc, 2 (11.1%) *C. herzi* were retained with a total of 4 CsMc and only 1 (33.3%) *S. scherzeri* were infected with total 6 CsMc [16]. No CsMc were detected in 4 fish species, i.e., *C. herzi* (n=57), *C. auratus* (42), *S. scherzeri* (11), and *C. carpio* (2), from the water systems of Seomjingang [18]. In this study, only one CsMc were found in only 1 (1.1%) out of 93 *C. herzi* and no CsMc were detected from 73 *C. auratus*, 36 *O. platycephala*, and 1 *S. scherzeri*.

So, the striped shinner, *P. herzi*, broadly live in the water systems of river in Korea and is highly susceptible with CsMc, it is

appropriate to evaluate the endemicities of *C. sinensis* infection as the index fish species. The number of this fish species examined were 197 (12.3%) in Seomjin-gang [15], 222 (13.5%) in Tamjin-gang [17], 169 (14.5%) in Wicheon [16], and 534 (24.3%) in this study. In the present study, 532 (99.6%) *P. herzi* were infected with 147 CsMc in average. Therefore, the endemicity (susceptibility index) was 146.4 (87.3-178.8 by the year examined). This value is higher than those in Tamjin-gang (103.2) [17] and Seomjin-gang (34.8) [15]. However, it is very low when we compared with that (1,550) in Wicheon [16]. Accordingly, based on the endemicity of CsMc in the index fish, *P. herzi*, we can know that Wicheon is highly endemic, Yangcheon in this study and Tamjin-gang are moderately endemic and Seomjin-gang is more or less low endemic areas.

In a highly endemic area, Wicheon in Gunwi-gun, Gyeongsangbuk-do, the infection tendency with CsMc was observed in positive fish species by the subfamily groups, i.e., Gobioninae, Acheilognathinae and Rasborinae, in the Cyprinidae fish hosts [16]. The endemicities in each group were 1,310.0, 39.9, and 5.3. Meanwhile, in this study performed in the moderately endemic area, they were 130.9, 3.2, and 0.3. The difference of endemicity between 2 regions was about 10 times in gobioninid group, 12.5 times in acheilognathinid and 17.7 times in rasborinid fish group. These findings suggest that the endemicity of CsMc is closely related with the fish groups in the cyprinid fish hosts and the endemicity difference in each fish groups from 2 regions is smaller in the susceptible gobioninid fish group, which is strongly related with host-parasite relationship, such as the infectivity of cercariae and susceptibility of host fish.

Conclusively, it is confirmed that the endemicity of CsMc is the moderate level in fishes from Yangcheon and the infection tendency of CsMc is obviously showed by the subfamily groups, i.e., Gobioninae, Acheilognathinae and Rasborinae, in the family Cyprinidae fish hosts from a moderately endemic site.

## ACKNOWLEDGMENTS

This study was supported by an anti-communicable diseases control program, 2014-E54002-00 (Investigation of fish-borne parasites and acquisition of their biological resources in the southern regions of Korea) of National Institute of Health (NIH), Korea Centers for Disease Control and Prevention (KCDCP), and a grant from Korea Association of Health Promotion (KAHP-2017: Survey on the effects of migratory bird

in the infection status of trematode metacercariae in fish from Junam Reservoir). We thank Jung-A Kim and Hee-Joo Kim (Department of Parasitology and Tropical Medicine, Gyeongsang National University College of Medicine, Jinju, Korea), for their help in the examination of fish.

## CONFLICT OF INTEREST

The authors have no conflicts of interest concerning the work reported in this paper.

## REFERENCES

1. Korea National Institute of Health, Korea Centers for Disease Control and Prevention. National survey of the prevalence of intestinal parasitic infections in Korea, 2012-The 8th Report. Osong Chungcheongbuk-do, Korea, 2013.
2. Seo BS, Lee SH, Cho SY, Chai JY, Hong ST, Han IS, Sohn JS, Cho BH, Ahn SR, Lee SK, Chung SC, Kang KS, Shim HS, Hwang IS. An epidemiologic study on clonorchiasis and metagonimiasis in riverside areas in Korea. *Korean J Parasitol* 1981; 19: 137-150.
3. Rim HJ. The current pathobiology and chemotherapy of clonorchiasis. *Korean J Parasitol* 1986; 24 (suppl): 1-141.
4. Cho SH, Lee KY, Lee BC, Cho PY, Cheun HI, Hong ST, Sohn WM, Kim TS. Prevalence of clonorchiasis in southern endemic areas of Korea in 2006. *Korean J Parasitol* 2008; 46: 133-137.
5. Kim HK, Cheun HI, Chung BS, Lee KY, Kim TS, Lee SE, Lee WJ, Cho SH. Prevalence of *Clonorchis sinensis* infections along the five major rivers in Republic of Korea, 2007. *Osong Public Health Res Perspect* 2010; 1: 43-49.
6. Jeong YI, Shin HE, Lee SE, Cheun HI, Ju JW, Kim JY, Park MY, Cho SH. Prevalence of *Clonorchis sinensis* infection among residents along 5 major rivers in the Republic of Korea. *Korean J Parasitol* 2016; 54: 215-219.
7. Bae KH, Ahn YK, Soh CT, Tsutsumi H. Epidemiological studies on *Clonorchis sinensis* infection along the Nam-river in Gyeongnam Province, Korea. *Korean J Parasitol* 1983; 21: 167-186 (In Korean).
8. Lee JS, Lee WJ, Kim TS, In TS, Kim WS, Kim SK. Current status and the changing pattern of the prevalence of clonorchiasis in the inhabitants in Sanchon-gun, Kyongsangnam-do, Korea. *Korean J Parasitol* 1993; 31: 207-213 (In Korean).
9. Hong SJ, Lee YH, Chung MH, Lee DH, Woo HC. Egg positive rates of *Clonorchis sinensis* and intestinal helminths among residents in Kagye-ri, Saengbiryang-myon, Sanchong-gun, Kyongsangnam-do. *Korean J Parasitol* 1994; 32: 271-273.
10. Kim BJ, Ock MS, Kim IS, Yeo UB. Infection status of *Clonorchis sinensis* in residents of Hamyang-gun, Gyeongsangnam-do, Korea. *Korean J Parasitol* 2002; 40: 191-193.
11. Kim EM, Kim JL, Choi SY, Kim JW, Kim S, Choi MH, Bae YM, Lee SH, Hong ST. Infection status of freshwater fish with meta-

- cercariae of *Clonorchis sinensis* in Korea. *Korean J Parasitol* 2008; 46: 247-251.
12. Cho SH, Sohn WM, Na BK, Kim TS, Kong Y, Eom K, Seok WS, Lee T. Prevalence of *Clonorchis sinensis* metacercariae in freshwater fish from three latitudinal regions of the Korean Peninsula. *Korean J Parasitol* 2011; 49: 385-398.
  13. Cho SH, Lee WJ, Kim TS, Seok WS, Lee TJ, Jeong KJ, Na BK, Sohn WM. Prevalence of zoonotic trematode metacercariae in freshwater fish from Gangwon-do, Korea. *Korean J Parasitol* 2014; 52: 399-412.
  14. Sohn WM, Na BK, Cho SH, Lee SW, Choi SB, Seok WS. Trematode metacercariae in freshwater fish from water systems of Hantangang and Imjingang in Republic of Korea. *Korean J Parasitol* 2015; 53: 289-298.
  15. Sohn WM, Na BK, Cho SH, Park MY, Kim CH, Hwang MA, No KW, Yoon KB, Lim HC. Prevalence of *Clonorchis sinensis* metacercariae in fish from water systems of Seomjin-gang (River). *Korean J Parasitol* 2017; 55: 305-312.
  16. Sohn WM, Na BK, Cho SH, Ju JW, Son DC. Prevalence and intensity of *Clonorchis sinensis* metacercariae in freshwater fish from Wicheon Stream in Gunwi-gun, Gyeongsangbuk-do, Korea. *Korean J Parasitol* 2018; 56: 43-50.
  17. Yoon KB, Lim HC, Jeon DY, Park S, Cho SH, Ju JW, Shin SS, Na BK, Sohn WM. Infection status with *Clonorchis sinensis* metacercariae in fish from Tamjin-gang (River) in Jeollanam-do, Republic of Korea. *Korean J Parasitol* 2018; 56: 183-188.
  18. Gyeongho-gang in Gyeongnam Province (South Korea) in Wikipedia - The free encyclopedia; <http://en.wikipedia.org>.
  19. Sohn WM. Fish-borne zoonotic trematode metacercariae in the Republic of Korea. *Korean J Parasitol* 2009; 47 (suppl): 103-113.
  20. Sohn WM. Invertebrate Founa of Korea Vol. 6, No. 1. Trematodes. Incheon, Korea. The National Institute of Biological Resources. 2013, pp 1-125.
  21. Kim IS, Kang EJ. Coloured Fishes of Korea. Seoul, Korea. Academy Publishing Company. 1993, pp 1-477 (in Korean).
  22. Choe S, Park H, Lee D, Kang Y, Jeon HK, Eom KS. Infections with digenean trematode metacercariae in two invasive alien fish, *Micropterus salmoides* and *Lepomis macrochirus*, in two rivers in Chungcheongbuk-do, Republic of Korea. *Korean J Parasitol* 2018; 56: 509-513.