



# Key Considerations for Radiologists When Diagnosing the Novel Coronavirus Disease (COVID-19)

Pinggui Lei, MD, PhD<sup>1</sup>, Jujiang Mao, MD<sup>1</sup>,  
 Zhaoshu Huang, MD<sup>1</sup>, Guoli Liu, MD<sup>1</sup>,  
 Pingxian Wang, MD<sup>2</sup>, Wen Song, MD<sup>1</sup>

Departments of <sup>1</sup>Radiology and <sup>2</sup>Medical Insurance, The Affiliated Hospital of Guizhou Medical University, Guiyang, China

Dear Editor,

We have read the publication titled “Chest Radiographic and CT Findings of the 2019 Novel Coronavirus Disease (COVID-19): Analysis of Nine Patients Treated in Korea” with great interest (1). The study suggested that the typical computed tomography (CT) features of COVID-19 pneumonia were ill-defined pure ground-glass opacities (GGO) or mixed GGO and consolidation. The statistical analysis revealed that there was a significant difference in the distribution of lesions ( $p < 0.05$ ) (1). We would like to share our opinion regarding diagnosing COVID-19 using radiological findings. Firstly, even though there are some typical radiological features of COVID-19 pneumonia, these features are not specific signs for different kinds of virus-related pneumonia or COVID-19 infection stages (2, 3). CT is a useful tool for the surveillance of pneumonic lesions and is more likely

Received February 28, 2020; accepted after revision February 28, 2020.

**Corresponding author:** Pinggui Lei, MD, PhD, Department of Radiology, The Affiliated Hospital of Guizhou Medical University, No.28, Guiyi Street, Yunyan District, Guiyang 550004, China.

• Tel: (86) 187 8611 8165 • Fax: (86) 851-86855119  
 • E-mail: pingguilei@foxmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

to detect early and/or mild lesions in the virus lifecycle (2, 4). Therefore, CT examinations play an important role in detecting or monitoring pulmonary parenchyma in patients suspected with COVID-19 pneumonia. Secondly, not all patients with COVID-19 have abnormal CT features (5). The real-time reverse transcriptase-polymerase chain reaction amplification of the viral DNA is considered the “gold standard”. Therefore, a CT examination and nuclear acid test should be used together for detecting COVID-19. Thirdly, fever and cough have been the most frequent initial symptoms (3), and radiologists should pay attention to these chief symptoms. Lastly, the history of exposure to other patients with COVID-19 or the epidemic area is critical to know. In conclusion, radiologists should make a comprehensive analysis when diagnosing COVID-19 infection that is beyond just assessing the radiological features.

## ORCID iD

Pinggui Lei

<https://orcid.org/0000-0001-7610-0292>

## REFERENCES

1. Yoon SH, Lee KH, Kim JY, Lee YK, Ko H, Kim KH, et al. Chest radiographic and CT findings of the 2019 novel coronavirus disease (COVID-19): analysis of nine patients treated in Korea. *Korean J Radiol* 2020 Feb 26 [Epub]. <https://doi.org/10.3348/kjr.2020.0132>
2. Koo HJ, Lim S, Choe J, Choi SH, Sung H, Do KH. Radiographic and CT features of viral pneumonia. *Radiographics* 2018;38:719-739
3. Pan F, Ye T, Sun P, Gui S, Liang B, Li L, et al. Time course of lung changes on chest CT during recovery from 2019 novel coronavirus (COVID-19) pneumonia. *Radiology* 2020 Feb 13 [Epub]. <https://doi.org/10.1148/radiol.2020200370>
4. Paul NS, Roberts H, Butany J, Chung T, Gold W, Mehta S, et al. Radiologic pattern of disease in patients with severe acute respiratory syndrome: the Toronto experience. *Radiographics* 2004;24:553-563
5. Fang Y, Zhang H, Xie J, Lin M, Ying L, Pang P, et al. Sensitivity of chest CT for COVID-19: comparison to RT-PCR. *Radiology* 2020 Feb 19 [Epub]. <https://doi.org/10.1148/radiol.2020200432>

## Response

Soon Ho Yoon, MD, PhD

Department of Radiology, Seoul National University College of Medicine,  
Seoul National University Hospital, Seoul, Korea

To the Editor,

We are grateful for you sharing your opinion regarding our publication. We agree that computed tomography (CT) is an important imaging modality for managing pneumonia related to the 2019 novel coronavirus disease (COVID-19). Early investigations, primarily based on CT images, expanded the understanding of radiologic manifestations and the temporal evolution of COVID-19 pneumonia (1-3). Nevertheless, unless COVID-19 is highly prevalent in an area, similar to Wuhan, we believe that chest CT should not be recommended as a primary surveillance tool in Korea. This is because the degree of incremental diagnostic yield of adding chest CT to reverse transcriptase-polymerase chain reaction was not sufficiently validated in a low-prevalence setting. Furthermore, the introduction of routine chest CT scanning requires extra resources, strict preventative measures against transmission, and increases the patient's radiation exposure. It is well-known that 15% to 20% of patients with coronavirus pneumonia have normal CT findings by referring to cases of Severe Acute Respiratory Syndrome and Middle East Respiratory Syndrome (4). Examining the presence and duration of patient symptoms, including fever and cough, is important when interpreting chest radiographic and CT examinations in suspected COVID-19 cases. Nevertheless, around 50% of patients with COVID-19 do not have a fever

at admission (5). The history of exposure is fundamental for making accurate COVID-19 pneumonia diagnoses, particularly for active case finding during the early phase of the outbreak, which requires collaborative image interpretation by clinicians and radiologists.

### ORCID iD

Soon Ho Yoon

<https://orcid.org/0000-0002-3700-0165>

### REFERENCES

1. Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, et al. CT imaging features of 2019 novel coronavirus (2019-nCoV). *Radiology* 2020 Feb 4 [Epub]. <https://doi.org/10.1148/radiol.2020200230>
2. Song F, Shi N, Shan F, Zhang Z, Shen J, Lu H, et al. Emerging 2019 novel coronavirus (2019-nCoV) pneumonia. *Radiology* 2020 Feb 6 [Epub]. <https://doi.org/10.1148/radiol.2020200274>
3. Pan F, Ye T, Sun P, Gui S, Liang B, Li L, et al. Time course of lung changes on chest CT during recovery from 2019 novel coronavirus (COVID-19) pneumonia. *Radiology* 2020 Feb 13 [Epub]. <https://doi.org/10.1148/radiol.2020200370>
4. Hosseiny M, Kooraki S, Gholamrezanezhad A, Reddy S, Myers L. Radiology perspective of coronavirus disease 2019 (COVID-19): lessons from severe acute respiratory syndrome and Middle East respiratory syndrome. *AJR Am J Roentgenol* 2020 Feb 28 [Epub]. <https://doi.org/10.2214/AJR.20.22969>
5. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al.; China Medical Treatment Expert Group for Covid-19. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020 Feb 28 [Epub]. <https://doi.org/10.1056/NEJMoa2002032>