

Commentary on "Autoimmune encephalitis and epilepsy: evolving definition and clinical spectrum"

Jieun Choi, MD, PhD

Department of Pediatrics, Seoul Metropolitan Government Seoul National University Boramae Medical Center, Seoul National University College of Medicine, Seoul, Korea

Acute encephalitis is neurological syndrome with diverse etiologies that occurs mostly in children, impacting their physical and cognitive development.¹⁾ Its estimated annual incidence in children is a minimum of 10.5/100,000.²⁾ The etiology of acute encephalitis remains unknown in more than half of cases. Herpes simplex virus (HSV) is the main virus found in the cerebrospinal fluid and blood; therefore, most cases of encephalitis might be infectious diseases.³⁾ However, recent advanced molecular technologies revealed that the presence of antibodies against neuronal proteins is associated with acute encephalitis syndromes, so-called autoimmune encephalitis (AIE). AIE reportedly occurred in many children presenting with seizures and behavioral and psychiatric changes that did not respond to conventional encephalitis antiviral or antibacterial medications. However, the overall prevalence of AIE remains unknown and its diagnostic algorithms remain undetermined. The mean incidence of antibody-mediated AIE was recently reported as 1.54 children/million in the Netherlands.⁴⁾

In a review article, Seo et al.⁵⁾ described the most relevant data of the immunologic and clinical characteristics of AIE recognized to date with the aim of assisting clinicians with the differential diagnosis and favoring early and effective treatment.

Autoantibodies found in AIE syndromes are classified into 3 types depending on the locations of their antigens, such as intracellular proteins, plasma membrane and synaptic receptors, and ion channels and/or other cell surface proteins. Anti-N-methyl-D-aspartate receptor (NMDAR) encephalitis, the most common AIE, is predominant in young women and children.⁶⁾ Anti-NMDAR encephalitis may be associated with HSV, *Mycoplasma pneumoniae* infection, measles, mumps, influenza A/H1N1, and group A hemolytic *Streptococcus* infection.⁷⁾ Autoimmune responses to the NR1 subunit of NMDAR were demonstrated to be triggered by various infectious agents in children. Movement disorders and seizures are significantly more common than psychiatric symptoms in children with AIE than in affected adults.

Gamma-aminobutyric acid A receptors (GABA_AR) are ligand-

gated ion channels. Patients with anti-GABA_AR encephalitis develop a rapidly progressive encephalitis with refractory seizures, status epilepticus, and/or epilepsy partialis continua. Nearly half of the reported cases occurred in children.⁸⁾ In the pediatric population, anti-GABA_AR encephalitis may progress as a postviral encephalitis and coexist with NMDAR antibodies.

Patients with suspected AIE should be thoroughly investigated using neuroimaging, electroencephalography, lumbar puncture, and serologic testing for appropriate biomarkers to confirm the diagnosis and exclude alternative etiologies. AIE might be detected in children presenting with multifaceted seizures and unexpected behavioral changes. The spectrum of neuropsychiatric manifestations is less clear in the pediatric than adult population with AIE. Screening for malignancy is mandatory in children to rule out paraneoplastic syndrome. The detection of specific autoantibodies confirms the diagnosis of AIE. Thus, clinical suspicion is important for the diagnosis and early intervention because the earlier immunomodulatory therapy is administered, the better the outcome. However, not all patients with AIE have antibodies; thus, the absence of antibodies does not rule out underlying autoimmune mechanism. Therefore, AIE must be considered a possible cause in patients with progressive sudden-onset sensorimotor deficits and movement disorders of unknown etiology.

The use of immunosuppressive therapy should not be delayed until the confirmation of a cancer diagnosis or the presence of antibodies to neuronal proteins and no definite contraindications for immunosuppressive therapy. Overall, outcomes in children are usually good, although this varies by the underlying tumor type and stage as well as the severity of the initial neurological symptoms and timing of the initial treatments. Abnormal magnetic resonance imaging findings, a clinical presentation with sensorimotor deficits, and a treatment delay exceeding 4 weeks were associated with worse clinical outcomes in 38 children with anti-NMDAR encephalitis.⁹⁾

The central nervous system (CNS) is considered an immune-privileged organ, but recent research revealed that it is an active

environment that interacts with diverse immune cells in both the disease state and the healthy brain. Immune processes play a vital role in CNS homeostasis, resilience, and brain reserve, and highly sensitive narrow ranges of inflammatory balance seem to be required for health. Crosstalk between the immune and nervous systems has been revealed in a wide spectrum of neurological diseases.¹⁰ Increased levels of proinflammatory cytokines are found in patients with epilepsy, developmental disorders, neurodegenerative diseases, and psychiatric disorders.¹¹⁻¹³ Even antibodies against fetal brain tissues, such as contactin-associated protein-like 2, were detected in the sera of pregnant mothers whose babies were later diagnosed with autism spectrum disorders but not in mothers of healthy children.¹⁴ Therefore, many neurologic and neurodevelopmental disorders in children might be re-evaluated and investigated in terms of immunologic interactions including autoantibodies, cytokines, and microRNAs.¹⁵

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

See the article “Autoimmune encephalitis and epilepsy: evolving definition and clinical spectrum” via <https://doi.org/10.3345/kjp.2019.00598>.

References

1. Venkatesan A. Epidemiology and outcomes of acute encephalitis. *Curr Opin Neurol* 2015;28:277-82.
2. Jmor F, Emsley HC, Fischer M, Solomon T, Lewthwaite P. The incidence of acute encephalitis syndrome in Western industrialised and tropical countries. *Virology* 2008;5:134.
3. Rozenberg F. Acute viral encephalitis. *Handb Clin Neurol* 2013;112:

- 1171-81.
4. de Bruijn M, Bruijstens AL, Bastiaansen AEM, van Sonderen A, Schreurs MWJ, Sillevius Smitt PAE, et al. Pediatric autoimmune encephalitis: recognition and diagnosis. *Neurol Neuroimmunol Neuroinflamm* 2020; 7:e682.
5. Seo JH, Lee YJ, Lee KH, Gireesh E, Skinner H, Westerveld M. Autoimmune encephalitis and epilepsy: evolving definition and clinical spectrum. *Clin Exp Pediatr* 2020;63:291-300.
6. Titulaer MJ, McCracken L, Gabilondo I, Armangue T, Glaser C, Iizuka T, et al. Treatment and prognostic factors for long-term outcome in patients with anti-NMDA receptor encephalitis: an observational cohort study. *Lancet Neurol* 2013;12:157-65.
7. Armangue T, Titulaer MJ, Malaga I, Bataller L, Gabilondo I, Graus F, et al. Pediatric anti-N-methyl-D-aspartate receptor encephalitis-clinical analysis and novel findings in a series of 20 patients. *J Pediatr* 2013;162:850-6.e2.
8. Spatola M, Petit-Pedrol M, Simabukuro MM, Armangue T, Castro FJ, Barcelo Artigues MI, et al. Investigations in GABAA receptor antibody-associated encephalitis. *Neurology* 2017;88:1012-20.
9. Bartels F, Krohn S, Nikolaus M, Johannsen J, Wickstrom R, Schimmel M, et al. Clinical and magnetic resonance imaging outcome predictors in pediatric anti-N-Methyl-D-aspartate receptor encephalitis. *Ann Neurol* 2020;88:148-59.
10. Choi J, Koh S. Role of brain inflammation in epileptogenesis. *Yonsei Med J* 2008;49:1-18.
11. Bergink V, Gibney SM, Drexhage HA. Autoimmunity, inflammation, and psychosis: a search for peripheral markers. *Biol Psychiatry* 2014;75:324-31.
12. Choi J, Nordli DR Jr, Alden TD, DiPatri A Jr, Laux L, Kelley K, et al. Cellular injury and neuroinflammation in children with chronic intractable epilepsy. *J Neuroinflammation* 2009;6:38. <https://doi.org/10.1186/1742-2094-6-38>.
13. Choi J, Min HJ, Shin JS. Increased levels of HMGB1 and pro-inflammatory cytokines in children with febrile seizures. *J Neuroinflammation* 2011;8:135.
14. Brimberg L, Mader S, Jeganathan V, Berlin R, Coleman TR, Gregersen PK, et al. Caspr2-reactive antibody cloned from a mother of an ASD child mediates an ASD-like phenotype in mice. *Mol Psychiatry* 2016;21:1663-71.
15. Hicks SD, Carpenter RL, Wagner KE, Pauley R, Barros M, Tierney-Aves C, et al. Saliva microRNA differentiates children with autism from peers with typical and atypical development. *J Am Acad Child Adolesc Psychiatry* 2020;59:296-308.